arXiv:0706.0066v1 [math.RT] 1 Jun 2007

THE $(\mathfrak{g}, K)$-MODULE STRUCTURES OF PRINCIPAL SERIES REPRESENTATIONS OF $Sp(3, \mathbf{R})$.

TADASHI MIYAZAKI

Abstract. We describe explicitly the whole structures of the $(\mathfrak{g}, K)$-modules of principal series representations of $Sp(3, \mathbf{R})$. We apply this result to determine the holonomic system characterizing those Whittaker functions.

1. Introduction

In the investigation of a representation π of a reductive Lie group G, it is a standard method to pass from the original π to its associated $(\mathfrak{g}, K)$-module. Also in many applications, it is important to understand various 'good' realizations of π in some function spaces. These functions can be sometimes described as solutions of some differential equations, say, the Casimir equations derived from the $(\mathfrak{g}, K)$-module structure of the given representation π.

For some 'small' semisimple Lie groups G, the $(\mathfrak{g}, K)$-module structures of the standard representations are completely described. For example, the description of them for $SL(2, \mathbf{R})$ is found in standard textbooks, and there are rather complete results for some groups of real rank 1, e.g. $SU(n, 1)$ in [7] and $Spin(1, 2n)$ in [17]. But, for Lie groups of higher rank there are few references as far as the author knows. It seems to be difficult to describe the whole $(\mathfrak{g}, K)$-module structures even for standard representations of classical groups of higher rank. However, in this paper we consider the case of the real symplectic group $Sp(3, \mathbf{R})$ of rank 3, and solve this problem generalizing the result of the paper [15] for $Sp(2, \mathbf{R})$ of rank 2.

Before describing our situation for $Sp(3, \mathbf{R})$, let us explain the problem in a more precise form. Let G be a real semisimple Lie group and $\mathfrak{g}$ the Lie algebra of G. Fix a maximal compact subgroup K of G. Since any standard $(\mathfrak{g}, K)$-modules are realized as subspaces of $L^2(K)$ as K-modules, we investigate the K-module structure of standard $(\mathfrak{g}, K)$-modules by the Peter-Weyl theorem. Because of the Cartan decomposition $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$, in order to describe the action of $\mathfrak{g}$ or $\mathfrak{g}_{\mathbf{C}} = \mathfrak{g} \otimes_{\mathbf{R}} \mathbf{C}$ it suffices to investigate the action of $\mathfrak{p}$ or $\mathfrak{p}_{\mathbf{C}}$. Therefore, the investigation of the action of $\mathfrak{p}$ or $\mathfrak{p}_{\mathbf{C}}$ is essential to give the explicit $(\mathfrak{g}, K)$-module structure of a standard representation. To investigate the action of $\mathfrak{p}_{\mathbf{C}}$, we compute the linear map $\Gamma_{\tau,i}$ defined as follows. Let (π, H_π) be a standard representation of G with the subspace $H_{\pi,K}$ of K-finite vectors. For a K-type (τ, V_τ) of π, and a nonzero K-homomorphism $\eta\colon V_\lambda \to H_{\pi,K}$, we define a linear map $\tilde{\eta}\colon \mathfrak{p}_{\mathbf{C}} \otimes_{\mathbf{C}} V_\lambda \to H_{\pi,K}$ by $X \otimes v \mapsto X \cdot \eta(v)$. Then $\tilde{\eta}$ is a K-homomorphism with $\mathfrak{p}_{\mathbf{C}}$ endowed with the adjoint action Ad of K. Let $V_\tau \otimes_{\mathbf{C}} \mathfrak{p}_{\mathbf{C}} \simeq \bigoplus_{i \in I} V_{\tau_i}$ be the decomposition into a direct sum of irreducible K-modules and ι_i an injective K-homomorphism from V_{τ_i} to $V_\tau \otimes_{\mathbf{C}} \mathfrak{p}_{\mathbf{C}}$ for each i. We define a linear map $\Gamma_{\tau,i}\colon \mathrm{Hom}_K(V_\tau, H_{\pi,K}) \to \mathrm{Hom}_K(V_{\tau_i}, H_{\pi,K})$ by $\eta \mapsto \tilde{\eta} \circ \iota_i$. These linear maps $\Gamma_{\tau,i}$ $(i \in I)$ characterize the action of $\mathfrak{p}_{\mathbf{C}}$. Our purpose of this paper is to give the explicit expressions of $\Gamma_{\tau,i}$ when π is a principal series representation of $G = Sp(3, \mathbf{R})$. This is described in Theorem 5.6.

As an application of the explicit expressions of $\Gamma_{\tau,i}$, we get a system of differential equations satisfied by some types of spherical functions. Here we consider only the case of the Whittaker functions. In order to introduce this application, let us recall the general setting of the theory of the spherical functions. Fix a closed subgroup R of G. Take a character ξ of R and consider its C^∞-induction $C^\infty \mathrm{Ind}_R^G(\xi)$. For an irreducible admissible representation (π, H_π) of G with the subspace $H_{\pi,K}$ of K-finite vectors, we consider the subspace $\mathrm{Hom}_{(\mathfrak{g}_{\mathbf{C}}, K)}(H_{\pi,K}, C^\infty \mathrm{Ind}_R^G(\xi))$

of intertwining operators. Consider the restriction of elements in this space to specific K-type as follows. Let (τ, V_τ) be a multiplicity one K-type of π and let $\iota\colon V_\tau \to H_\pi$ be a nonzero K-homomorphism. For $\Phi \in \mathrm{Hom}_{(\mathfrak{g}_{\mathbf{C}},K)}(H_{\pi,K}, C^\infty \mathrm{Ind}_R^G(\xi))$, we can define the function ϕ_{π,τ^*} contained in the space $C^\infty_{\xi,\tau^*}(R\backslash G/K)$ of V_τ^*-valued smooth functions on G satisfying $f(rgk) = \xi(r)\tau^*(k)^{-1}f(g)$ for all $(r,g,k) \in R \times G \times K$ by $\Phi(\iota(v))(g) = \langle v, \phi_{\pi,\tau^*}\rangle$ $(g \in G,\ v^* \in V_\tau)$. Here τ^* means the contragradient representation of τ and $\langle , \rangle$ is the canonical pairing of $V_\tau \times V_{\tau^*}$. When R is a maximal unipotent subgroup of G and ξ is a unitary character of R, the space $\mathrm{Hom}_{(\mathfrak{g}_{\mathbf{C}},K)}(H_{\pi,K}, C^\infty \mathrm{Ind}_R^G(\xi))$ is called the space of Whittaker functionals and ϕ_{π,τ^*} is called a Whittaker function.

In the case of $G = Sp(2,\mathbf{R})$, the explicit formulas of Whittaker functions for various standard representations are given in the papers [3], [5], [12], [13], [14] as well as the generalized Whittaker functions in the papers [2], [11]. In these papers, the explicit formulas of spherical functions are given as solutions of the system of differential equations which are obtained from shift operators and the Casimir element. Since the Casimir element is represented by a composite of the shift operators, the utilization of the shift operator is essential. Our operator $\Gamma_{\tau,i}$ is compatible with the shift operator and we give the holonomic systems of differential equations characterizing Whittaker functions in Theorem 6.12. Therefore, we can compute explicit formula of Whittaker functions of principal series representations of $G = Sp(3,\mathbf{R})$ by using the result of this paper. We hope that this interesting possibility will be considered in future work.

We give the contents of this paper. In Section 2, we recall the structure of $Sp(3,\mathbf{R})$ and define a principal series representation, that is, a standard representation obtained by a parabolic induction with respect to the minimal parabolic subgroup $P_{\min}$. In Section 3, we introduce the monomial basis of a finite dimensional irreducible representation of $\mathfrak{k}_{\mathbf{C}} \simeq \mathfrak{gl}(3,\mathbf{C})$ and investigate adjoint representation of K on $\mathfrak{p}_{\mathbf{C}}$. Here the monomial basis is an alias of Gelfand Zelevinsky basis, Which is twisted dual of the crystal basis of Kashiwara or the canonical basis of Lusztig. In Section 4, we give the explicit expressions of $\iota_i\colon V_{\tau_i} \to V_\tau \otimes_{\mathbf{C}} \mathfrak{p}_{\mathbf{C}}$. Section 5 is the main body of this paper and we give the matrix representation of $\Gamma_{\tau,i}$ with respect to the induced basis from the monomial basis in Theorem 5.6. In Section 6, we introduce some examples of $\Gamma_{\tau,i}$ and give the holonomic systems of differential equations characterizing Whittaker functions in Theorem 6.12.

This is an enhanced version of the Master's thesis [10]. The author would like to express his gratitude to Takayuki Oda for valuable advice on this work. He also would like to thank Miki Hirano to provide a reference for the computation of Clebsch-Gordan coefficients.

2. PRELIMINARIES

2.1. Groups and algebras.

We denote by $\mathbf{Z}$,$\mathbf{R}$ and $\mathbf{C}$ the ring of rational integers, the real number field and the complex number field, respectively. Let 1_n (resp. O_n) be the unit (resp. the zero) matrix in the space $M_n(\mathbf{R})$ of real matrices of size n.

The real symplectic group $G = Sp(3,\mathbf{R})$ of degree three is defined by

$$G = Sp(3,\mathbf{R}) = \{g \in SL(6,\mathbf{R}) \mid {}^t g J_3 g = J_3\}, \quad J_3 = \begin{pmatrix} O_3 & 1_3 \\ -1_3 & O_3 \end{pmatrix},$$

which is connected, semisimple, and split over $\mathbf{R}$. Here ${}^t g$ and g^{-1} mean the transpose and the inverse of g, respectively. Let $\theta\colon G \ni g \mapsto {}^t g^{-1} \in G$ be a Cartan involution of G. Then

$$\begin{aligned} K &= \{g \in G \mid \theta(g) = g\} \\ &= \left\{ g = \begin{pmatrix} A & B \\ -B & A \end{pmatrix} \in G \;\middle|\; A + \sqrt{-1}B \in U(3) \right\} \end{aligned}$$

is a maximal compact subgroup of G which is isomorphic to the unitary group $U(3)$ of degree three. The Lie algebra of G is given by

$$\mathfrak{g} = \mathfrak{sp}(3, \mathbf{R}) = \{X \in M_6(\mathbf{R}) \mid J_3 X + {}^t X J_3 = 0\}.$$

If we denote the differential of θ again by θ, then we have $\theta(X) = -{}^t X$ for $X \in \mathfrak{g}$. Let $\mathfrak{k}$ and $\mathfrak{p}$ be the $+1$ and the -1 eigenspaces of $\theta \in \mathfrak{g}$, respectively, that is,

$$\mathfrak{k} = \left\{ X = \begin{pmatrix} A & B \\ -B & A \end{pmatrix} \in \mathfrak{g} \;\middle|\; A, B \in M_3(\mathbf{R}), {}^t A = -A, {}^t B = B \right\},$$
$$\mathfrak{p} = \left\{ X = \begin{pmatrix} A & B \\ B & -A \end{pmatrix} \in \mathfrak{g} \;\middle|\; A, B \in M_3(\mathbf{R}), {}^t A = A, {}^t B = B \right\}.$$

Then $\mathfrak{k}$ is the Lie algebra of K which is isomorphic to the unitary algebra

$$\mathfrak{u}(3) = \{A + \sqrt{-1}B \in M_3(\mathbf{C}) \mid A, B \in M_3(\mathbf{R}), {}^t A = -A, {}^t B = B\}$$

of degree three, and $\mathfrak{g}$ has the Cartan decomposition $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$. We fix an isomorphism $\kappa \colon \mathfrak{u}(3) \to \mathfrak{k}$ which is given by the inverse of

$$\mathfrak{k} \ni \begin{pmatrix} A & B \\ -B & A \end{pmatrix} \mapsto A + \sqrt{-1}B \in \mathfrak{u}(3).$$

For a Lie algebra $\mathfrak{l}$, we denote by $\mathfrak{l}_{\mathbf{C}} = \mathfrak{l} \otimes_{\mathbf{R}} \mathbf{C}$ the complexification of $\mathfrak{l}$. For $1 \le i, j \le 3$, we denote by E_{ij} the matrix unit in $M_3(\mathbf{R})$ with entry 1 at (i, j)-th component and 0 at other entries. Take a compact Cartan subalgebra $\mathfrak{h} = \bigoplus_{i=1}^{3} \mathbf{R} T_i$ where $T_i = \kappa(\sqrt{-1}E_{ii}) \in \mathfrak{k}$. For $1 \le i \le 3$, define a linear form β_i on $\mathfrak{h}_{\mathbf{C}}$ by $\beta_i(T_j) = \sqrt{-1}\delta_{ij}$, $1 \le j \le 3$. Here δ_{ij} is the Kronecker's delta. Then the set Δ of the roots for $(\mathfrak{h}_{\mathbf{C}}, \mathfrak{g}_{\mathbf{C}})$ is given by

$$\Delta = \Delta(\mathfrak{h}_{\mathbf{C}}, \mathfrak{g}_{\mathbf{C}}) = \{\pm 2\beta_i \ (1 \le i \le 3),\ \pm\beta_j \pm \beta_k \ (1 \le j < k \le 3)\},$$

and the subset $\Delta^+ = \{2\beta_i \ (1 \le i \le 3),\ \beta_j \pm \beta_k \ (1 \le j < k \le 3)\}$ form a positive root system. Let

$$\Delta_c^+ = \{\beta_j - \beta_k \ (1 \le j < k \le 3)\},$$
$$\Delta_n^+ = \{2\beta_i \ (1 \le i \le 3),\ \beta_j + \beta_k \ (1 \le j < k \le 3)\}$$

be the set of compact and non-compact positive roots, respectively. If we denote the root space for $\beta \in \Delta$ by $\mathfrak{g}_\beta$, then $\mathfrak{k}_{\mathbf{C}} \simeq \mathfrak{gl}(3, \mathbf{C})$ and $\mathfrak{p}_{\mathbf{C}}$ have the decompositions

$$\mathfrak{k}_{\mathbf{C}} = \mathfrak{h}_{\mathbf{C}} \oplus \bigoplus_{\beta \in \Delta_c} \mathfrak{g}_\beta, \quad \Delta_c = \Delta_c^+ \cup (-\Delta_c^+),$$
$$\mathfrak{p}_{\mathbf{C}} = \mathfrak{p}_+ \oplus \mathfrak{p}_-, \quad \mathfrak{p}_\pm = \bigoplus_{\beta \in \Delta_n^+} \mathfrak{g}_{\pm\beta}.$$

Now we take a basis of $\mathfrak{k}_{\mathbf{C}}$ and $\mathfrak{p}_\pm$ consisting of root vectors. If we denote the extension of the isomorphism κ to their complexifications again by κ, then we have $\kappa(E_{ij}) \in \mathfrak{g}_{\beta_i - \beta_j}$ for each $1 \le i \ne j \le 3$ and thus the set $\{\kappa(E_{ij}) \mid 1 \le i, j \le 3\}$ forms a basis of $\mathfrak{k}_{\mathbf{C}}$. On the other hand, if we define a map

$$p_\pm \colon \{X \in M_3(\mathbf{C}) \mid X = {}^t X\} \ni X \mapsto \begin{pmatrix} X & \pm\sqrt{-1}X \\ \pm\sqrt{-1}X & -X \end{pmatrix} \in \mathfrak{p}_\pm,$$

then the element $X_{\pm ij} = p_\pm((E_{ij} + E_{ji})/2)$ is a root vector in $\mathfrak{g}_{\pm(\beta_i + \beta_j)}$ for $1 \le i \le j \le 3$ and the set $\{X_{\pm ij} \mid 1 \le i \le j \le 3\}$ gives a basis of $\mathfrak{p}_\pm$.

Put $\mathfrak{a} = \bigoplus_{i=1}^{3} \mathbf{R} H_i$ with $H_1 = \mathrm{diag}(1, 0, 0, -1, 0, 0)$, $H_2 = \mathrm{diag}(0, 1, 0, 0, -1, 0)$, $H_3 = \mathrm{diag}(0, 0, 1, 0, 0, -1)$. Then $\mathfrak{a}$ is a maximal abelian subalgebra of $\mathfrak{p}$. For each $1 \le i \le 3$, we define a linear form e_i on $\mathfrak{a}$ by $e_i(H_j) = \delta_{ij}$ for $1 \le j \le 3$. The set Σ of the restricted roots for $(\mathfrak{a}, \mathfrak{g})$ is given by

$$\Sigma = \Sigma(\mathfrak{a}, \mathfrak{g}) = \{\pm 2e_i \ (1 \le i \le 3), \ \pm e_j \pm e_k \ (1 \le j < k \le 3)\},$$

and the subset $\Sigma^+ = \{2e_i \ (1 \le i \le 3), \ e_j \pm e_k \ (1 \le j < k \le 3)\}$ forms a positive root system. For each $\alpha \in \Sigma$, we denote the restricted root space by $\mathfrak{g}_\alpha$ and choose a restricted root vector E_α in $\mathfrak{g}_\alpha$ as follows.

$$E_{2e_i} = \begin{pmatrix} O_3 & E_{ii} \\ O_3 & O_3 \end{pmatrix}, \ 1 \le i \le 3,$$

$$E_{e_j+e_k} = \begin{pmatrix} O_3 & E_{jk} + E_{kj} \\ O_3 & O_3 \end{pmatrix}, \ E_{e_j-e_k} = \begin{pmatrix} E_{jk} & O_3 \\ O_3 & -E_{kj} \end{pmatrix}, 1 \le j < k \le 3,$$

and $E_{-\alpha} = \theta(E_\alpha)$ for $\alpha \in \Sigma^+$. If we put $\mathfrak{n} = \bigoplus_{\alpha \in \Sigma^+} \mathfrak{g}_\alpha$ then $\mathfrak{g}$ has an Iwasawa decomposition $\mathfrak{g} = \mathfrak{n} \oplus \mathfrak{a} \oplus \mathfrak{k}$. Also we have $G = N_{\min} A_{\min} K$, where $A_{\min} = \exp(\mathfrak{a})$ and $N_{\min} = \exp(\mathfrak{n})$.

2.2. **Definition of principal series representations of G.** Let $P_{\min} = M_{\min} A_{\min} N_{\min}$ be the minimal parabolic subgroup of G, where

$$M_{\min} = Z_K(A_{\min}) = \{\mathrm{diag}(\varepsilon_1, \varepsilon_2, \varepsilon_3, \varepsilon_1, \varepsilon_2, \varepsilon_3) \mid \varepsilon_i \in \{\pm 1\} \ (1 \le i \le 3)\} \simeq \{\pm 1\}^{\oplus 3}.$$

For $\nu \in \mathrm{Hom}_{\mathbf{R}}(\mathfrak{a}, \mathbf{C})$, we define a coordinate $(\nu_1, \nu_2, \nu_3) \in \mathbf{C}^3$ by $\nu_i = \nu(H_i)$, $1 \le i \le 3$. Then the half sum $\rho = \frac{1}{2}\left(\sum_{\alpha \in \Sigma_+} \alpha\right) = 3e_1 + 2e_2 + e_3$ of the positive roots has coordinate $(\rho_1, \rho_2, \rho_3) = (3, 2, 1)$. We define a quasicharacter $e^\nu \colon A_{\min} \to \mathbf{C}^\times$ by

$$e^\nu(\mathrm{diag}(a_1, a_2, a_3, a_1^{-1}, a_2^{-1}, a_3^{-1})) = a_1^{\nu_1} a_2^{\nu_2} a_3^{\nu_3}.$$

Moreover, we fix a character σ of $M_{\min}$. σ is realized by $(\sigma_1, \sigma_2, \sigma_3) \in \{0, 1\}^{\oplus 3}$ such that

$$\sigma(\mathrm{diag}(\varepsilon_1, \varepsilon_2, \varepsilon_3, \varepsilon_1, \varepsilon_2, \varepsilon_3)) = \varepsilon_1^{\sigma_1} \varepsilon_2^{\sigma_2} \varepsilon_3^{\sigma_3} \in \mathbf{C} \ \text{ for } \ \varepsilon_i \in \{\pm 1\}, \ 1 \le i \le 3.$$

With these data (σ, ν), the parabolic induction

$$\pi_{(\sigma,\nu)} = \mathrm{Ind}_{P_{\min}}^G (\sigma \otimes e^{\nu+\rho} \otimes 1_{N_{\min}})$$

is a Hilbert representation of G by the right regular action on the Hilbert space $H_{(\sigma,\nu)}$ which is the completion of

$$H^\infty_{(\sigma,\nu)} = \left\{ f \colon G \to \mathbf{C} \text{ smooth} \ \middle| \ \begin{array}{c} f(manx) = \sigma(m) e^{\nu+\rho}(a) f(x) \\ \text{for } \ m \in M_{\min}, \ a \in A_{\min}, \ n \in N_{\min}, \ x \in G \end{array} \right\}$$

with an inner product

$$(f_1, f_2) = \int_K f_1(k) \overline{f_2(k)} dk \quad \text{ for } \quad f_1, f_2 \in H^\infty_{(\sigma,\nu)}.$$

Here dk is a Haar measure of K.

3. The monomial basis for simple $\mathfrak{gl}(3, \mathbf{C})$-modules and the adjoint representation of $\mathfrak{k}_{\mathbf{C}} \simeq \mathfrak{gl}(3, \mathbf{C})$ on $\mathfrak{p}_\pm$

In this section, we recall some basic facts about the representations of $\mathfrak{k}_{\mathbf{C}} \simeq \mathfrak{gl}(3, \mathbf{C})$, and evaluate the adjoint representations of $\mathfrak{k}_{\mathbf{C}}$ on $\mathfrak{p}_\pm$.

3.1. **The highest weight theory for $\mathfrak{gl}(n,\mathbf{C})$.** We recall the highest weight theory for $\mathfrak{gl}(n,\mathbf{C})$. A *weight* of length n is an integral vector $\gamma = (\gamma_1, \gamma_2, \cdots, \gamma_n) \in \mathbf{Z}^n$. The weight γ is called *dominant* if $\gamma_1 \geq \gamma_2 \geq \cdots \geq \gamma_n$. It is well-known that every irreducible finite dimensional representation (τ, V) of $\mathfrak{gl}(n,\mathbf{C})$ has a *weight space decomposition*

$$V = \bigoplus_{\gamma} V(\gamma), \quad V(\gamma) = \{v \in V \mid E_{ii}v = \gamma_i v,\ 1 \leq i \leq n\}.$$

There is a dominant weight λ which satisfies $\lambda \geq \gamma$ in the lexicographical order for any weight γ such that $V(\gamma) \neq 0$. Such dominant weight is called *the highest weight* and the representation (τ, V) is labeled by the highest weight, i.e., $(\tau_\lambda, V_\lambda)$.

3.2. **Gelfand Tsetlin patterns.** Recall that a *Gelfand-Tsetlin pattern* (which simply we call *G-pattern*) of type $\mathbf{m}_3 = (m_{13}, m_{23}, m_{33})$ is a triangular array

$$M = \begin{pmatrix} \mathbf{m}_3 \\ \mathbf{m}_2 \\ \mathbf{m}_1 \end{pmatrix} = \begin{pmatrix} m_{13}\ m_{23}\ m_{33} \\ m_{12}\ m_{22} \\ m_{11} \end{pmatrix}$$

of integers satisfying the conditions

$$m_{13} \geq m_{12} \geq m_{23} \geq m_{22} \geq m_{33}, \quad m_{12} \geq m_{11} \geq m_{22}. \tag{3.1}$$

The weight $\gamma^M = (\gamma_1^M, \gamma_2^M, \gamma_3^M)$ of a G-pattern M is defined from the equations

$$\gamma_1^M = m_{11}, \quad \gamma_1^M + \gamma_2^M = m_{12} + m_{22}, \quad \gamma_1^M + \gamma_2^M + \gamma_3^M = m_{13} + m_{23} + m_{33}.$$

For a G-pattern M, we define

$$M\begin{pmatrix} i_{13}\ i_{23}\ i_{33} \\ i_{12}\ i_{22} \\ i_{11} \end{pmatrix} = \begin{pmatrix} m_{13}+i_{13} \quad m_{23}+i_{23} \quad m_{33}+i_{33} \\ m_{12}+i_{12} \quad m_{22}+i_{22} \\ m_{11}+i_{11} \end{pmatrix}.$$

If the vector (i_{13}, i_{23}, i_{33}) is zero, we omit the top row in the left hand side of the above defining equality. So the left hand side is written as

$$M\begin{pmatrix} i_{12}\ i_{22} \\ i_{11} \end{pmatrix}.$$

A convenient symbol is $M[k]$, which is defined by

$$M\begin{pmatrix} k \ \ -k \\ 0 \end{pmatrix}.$$

We note that G-patterns M_1 and M_2 have the same type and weight if and only if $M_1[k] = M_2$ for some $k \in \mathbf{Z}$.

We define some functions of G-patterns. We set

$$\delta(M) = \gamma_2^M - m_{23} = m_{12} + m_{22} - m_{11} - m_{23}.$$

Let $\chi_+(M)$ and $\chi_-(M)$ be the characteristic functions of the sets $\{M \mid \delta(M) > 0\}$ and $\{M \mid \delta(M) < 0\}$, respectively. More generally we introduce functions $\chi_\pm^{(i)}(M)$ by

$$\chi_+^{(i)}(M) = \begin{cases} 1, & \delta(M) > i \\ 0, & \delta(M) \leq i \end{cases}, \quad \chi_-^{(i)}(M) = \begin{cases} 1, & \delta(M) < -i \\ 0, & \delta(M) \geq -i \end{cases}.$$

Then we have $\chi_+(M) = \chi_+^{(0)}(M)$ and $\chi_-(M) = \chi_-^{(0)}(M)$.

We introduce 'piecewise-linear' functions $C_1(M)$, $\bar{C}_1(M)$ and $C_2(M)$ by

$$C_1(M) = \begin{cases} m_{11} - m_{22}, & \text{if } \delta(M) \geq 0 \\ m_{12} - m_{23}, & \text{if } \delta(M) \leq 0 \end{cases}, \quad \bar{C}_1(M) = \begin{cases} m_{23} - m_{22}, & \text{if } \delta(M) \geq 0 \\ m_{12} - m_{11}, & \text{if } \delta(M) \leq 0 \end{cases},$$

and

$$C_2(M) = C_1(M)\bar{C}_1(M).$$

Another expressions of $C_1(M)$ and $\bar{C}_1(M)$ are

$$C_1(M) = \min\{m_{11} - m_{22},\ m_{12} - m_{23}\}, \quad \bar{C}_1(M) = \min\{m_{23} - m_{22},\ m_{12} - m_{11}\}.$$

3.3. **The monomial basis in the sense of Gelfand-Zelevinsky.** We recall the definition of the monomial basis in the sense of Gelfand-Zelevinsky.

For a weight subspace $V_\lambda(\gamma)$ and a dominant weight $\nu = (\nu_1, \nu_2, \cdots, \nu_n)$, we set

$$V_\lambda(\gamma, \nu) = \{v \in V_\lambda(\gamma) \mid E_{i\,i+1}^{\nu_i - \nu_{i+1}+1} v = 0,\ 1 \le i \le n-1\}.$$

A basis B in V_λ is called *proper* if each of subspaces $V_\lambda(\gamma, \nu)$ (for all possible γ, ν) is spanned by its subset $B \cap V_\lambda(\gamma, \nu)$. It is known that the representation $(\tau_\lambda, V_\lambda)$ of $\mathfrak{gl}(3, \mathbf{C})$ has a proper basis, which is unique up to scalar multiple. Gelfand and Zelevinsky normalized the scalar factor somehow to get the formulas in Proposition 3.1. The normalized proper basis is called *the monomial basis* because it is twisted dual of the crystal basis of Kashiwara or the canonical basis of Lusztig. For the representation $(\tau_{\mathbf{m}_3}, V_{\mathbf{m}_3})$, the monomial basis in $V_{\mathbf{m}_3}(\gamma)$ is parameterized by the G-patterns of type $\mathbf{m}_3$ whose weights are γ. We denote the monomial basis of $V_{\mathbf{m}_3}$ by $\{f(M)\}_{M \in G(\mathbf{m}_3)}$. Here $G(\mathbf{m}_3)$ is the set of the G-patterns of type $\mathbf{m}_3$.

The action of $\mathfrak{k}_{\mathbf{C}} \simeq \mathfrak{gl}(3, \mathbf{C})$ is given by following formulas.

Proposition 3.1. *(Gelfand-Zelevinsky) The action of simple root vectors on the monomial basis* $\{f(M)\}_{M \in G(\mathbf{m}_3)}$ *of* $V_{\mathbf{m}_3}$ *are given as follows.*

$$\begin{aligned}
E_{12}f(M) &= (m_{12} - m_{11}) f\left(M\left(\begin{smallmatrix} 0\ 0 \\ 1 \end{smallmatrix}\right)\right) + (m_{23} - m_{22})\chi_+(M) f\left(M\left(\begin{smallmatrix} 0\ 0 \\ 1 \end{smallmatrix}\right)[-1]\right)\\
E_{21}f(M) &= (m_{11} - m_{22}) f\left(M\left(\begin{smallmatrix} 0\ 0 \\ -1 \end{smallmatrix}\right)\right) + (m_{12} - m_{23})\chi_-(M) f\left(M\left(\begin{smallmatrix} 0\ 0 \\ -1 \end{smallmatrix}\right)[-1]\right)\\
E_{23}f(M) &= (m_{13} - m_{12}) f\left(M\left(\begin{smallmatrix} 1\ 0 \\ 0 \end{smallmatrix}\right)\right) + \{m_{13} - m_{12} - \delta(M)\}\chi_-(M) f\left(M\left(\begin{smallmatrix} 1\ 0 \\ 0 \end{smallmatrix}\right)[-1]\right)\\
E_{32}f(M) &= (m_{22} - m_{33}) f\left(M\left(\begin{smallmatrix} 0\ -1 \\ 0 \end{smallmatrix}\right)\right) + \{m_{22} - m_{33} + \delta(M)\}\chi_+(M) f\left(M\left(\begin{smallmatrix} 0\ -1 \\ 0 \end{smallmatrix}\right)[-1]\right)
\end{aligned}$$

In the right hand side of above formulas, we put $f(M') = 0$ *if* M' *is a triangular array which does not satisfy the condition (3.1) of G-patterns.*

In the later section, we need the action of root vectors on the monomial basis of $V_{\mathbf{m}_3}$. So we compute the action of E_{13} and E_{31}. Since $E_{13} = [E_{12}, E_{23}]$ and $E_{31} = [E_{32}, E_{21}]$, we obtain

$$E_{13}f(M) = [E_{12}, E_{23}]f(M) = (m_{13} - m_{12}) f\left(M\left(\begin{smallmatrix} 1\ 0 \\ 1 \end{smallmatrix}\right)\right) - \bar{C}_1(M) f\left(M\left(\begin{smallmatrix} 1\ 0 \\ 1 \end{smallmatrix}\right)[-1]\right), \tag{3.2}$$

$$E_{31}f(M) = [E_{32}, E_{21}]f(M) = (m_{33} - m_{22}) f\left(M\left(\begin{smallmatrix} 0\ -1 \\ -1 \end{smallmatrix}\right)\right) + C_1(M) f\left(M\left(\begin{smallmatrix} 0\ -1 \\ -1 \end{smallmatrix}\right)[-1]\right). \tag{3.3}$$

Here $[,]$ is a Lie bracket, i.e., $[X, Y] = XY - YX$ for $X, Y \in \mathfrak{gl}(3, \mathbf{C})$.

The monomial basis has a interesting symmetry property. For each G-pattern M, we define the dual pattern $\hat{M}$ by

$$\hat{M} = \begin{pmatrix} -m_{33} \quad -m_{23} \quad -m_{13} \\ -m_{22} \quad -m_{12} \\ -m_{11} \end{pmatrix}.$$

If M is a G-pattern of type $\mathbf{m}_3$ and weight γ^M then $\hat{M}$ is a G-pattern of type $\hat{\mathbf{m}}_3 = (-m_{33}, -m_{23}, -m_{13})$ and weight $-\gamma^M = (-\gamma_1^M, -\gamma_2^M, -\gamma_3^M)$.

Proposition 3.2. *Let* ω *be the automorphism of* $\mathfrak{gl}(3, \mathbf{C})$ *defined by* $\omega(E_{ii}) = -E_{ii}$ *and* $\omega(E_{jk}) = E_{kj}$ *for* $i, j, k \in \{1, 2, 3\}$ *such that* $|j - k| = 1$. *Let* $T_{\mathbf{m}_3} \colon V_{\mathbf{m}_3} \to V_{\hat{\mathbf{m}}_3}$ *be the linear map defined by* $T_{\mathbf{m}_3}(f(M)) = f(\hat{M})$ *for* $M \in G(\mathbf{m}_3)$. *Then* $X \circ T_{\mathbf{m}_3} = T_{\mathbf{m}_3} \circ \omega(X)$.

For the existence, uniqueness and properties of the monomial basis, we refer to the paper [1].

In the later setions, we give the explicit expressions of various K-homomorphisms in terms of the monomial basis. However, the monomial basis have the ambiguity of scalar multiple. Thus, we have to fix the monomial basis for simple K-modules.

Definition 3.3. *A simple K-module V_λ equipped with the fixed monomial basis $\{f(M)\}_{M\in G(\lambda)}$ is called a simple K-module with the marking $\{f(M)\}_{M\in G(\lambda)}$.*

3.4. **The adjoint representations of $\mathfrak{k}_{\mathbf{C}}$ on $\mathfrak{p}_\pm$.** We denote by $\mathbf{e}_i$ the unit vector of degree three with its i-th component 1 and the remaining component 0. It is known that both of $\mathfrak{p}_\pm$ become K-modules via the adjoint action of K. Concerning this, we have the following lemma.

Lemma 3.4. *We have isomorphisms $i_{\mathfrak{p}_+} : \mathfrak{p}_+ \to V_{2\mathbf{e}_1}$ and $i_{\mathfrak{p}_-} : \mathfrak{p}_- \to V_{-2\mathbf{e}_3}$ by the correspondences between their basis*

$$\begin{aligned}&(X_{+11}, X_{+12}, X_{+13}, X_{+22}, X_{+23}, X_{+33})\\ &\leftrightarrow (f\left(\begin{smallmatrix}2\,0\,0\\2\,0\\2\end{smallmatrix}\right), f\left(\begin{smallmatrix}2\,0\,0\\2\,0\\1\end{smallmatrix}\right), f\left(\begin{smallmatrix}2\,0\,0\\1\,0\\1\end{smallmatrix}\right), f\left(\begin{smallmatrix}2\,0\,0\\2\,0\\0\end{smallmatrix}\right), f\left(\begin{smallmatrix}2\,0\,0\\1\,0\\0\end{smallmatrix}\right), f\left(\begin{smallmatrix}2\,0\,0\\0\,0\\0\end{smallmatrix}\right)),\end{aligned}$$

$$\begin{aligned}&(X_{-33}, -X_{-23}, X_{-22}, X_{-13}, -X_{-12}, X_{-11})\\ &\leftrightarrow (f\left(\begin{smallmatrix}0\,0\,-2\\0\,0\\0\end{smallmatrix}\right), f\left(\begin{smallmatrix}0\,0\,-2\\0\,-1\\0\end{smallmatrix}\right), f\left(\begin{smallmatrix}0\,0\,-2\\0\,-2\\0\end{smallmatrix}\right), f\left(\begin{smallmatrix}0\,0\,-2\\0\,-1\\-1\end{smallmatrix}\right), f\left(\begin{smallmatrix}0\,0\,-2\\0\,-2\\-1\end{smallmatrix}\right), f\left(\begin{smallmatrix}0\,0\,-2\\0\,-2\\-2\end{smallmatrix}\right))\end{aligned}$$

Proof. By direct computation, we have the following tables of the adjoint actions of Cartan subalgebra and simple root vectors of $\mathfrak{k}_{\mathbf{C}}$ on the basis $\{X_{\pm ij}\}_{1\leq i\leq j\leq 3}$ of $\mathfrak{p}_\pm$.

	$\kappa(E_{11})$	$\kappa(E_{22})$	$\kappa(E_{33})$	$\kappa(E_{12})$	$\kappa(E_{21})$	$\kappa(E_{23})$	$\kappa(E_{32})$	$\kappa(E_{13})$	$\kappa(E_{31})$
X_{+11}	$2X_{+11}$	0	0	0	$2X_{+12}$	0	0	0	$2X_{+13}$
X_{+12}	X_{+12}	X_{+12}	0	X_{+11}	X_{+22}	0	X_{+13}	0	X_{+23}
X_{+13}	X_{+13}	0	X_{+13}	0	X_{+23}	X_{+12}	0	0	0
X_{+22}	0	$2X_{+22}$	0	$2X_{+12}$	0	0	$2X_{+23}$	X_{+11}	X_{+33}
X_{+23}	0	X_{+23}	X_{+23}	X_{+13}	0	X_{+22}	X_{+33}	X_{+12}	0
X_{+33}	0	0	$2X_{+33}$	0	0	$2X_{+23}$	0	$2X_{+13}$	0

TABLE 1. The adjoint actions of $\mathfrak{k}_{\mathbf{C}}$ on the basis $\{X_{+ij}\}_{1\leq i\leq j\leq 3}$ of $\mathfrak{p}_+$

	$\kappa(E_{11})$	$\kappa(E_{22})$	$\kappa(E_{33})$	$\kappa(E_{12})$	$\kappa(E_{21})$	$\kappa(E_{23})$	$\kappa(E_{32})$	$\kappa(E_{13})$	$\kappa(E_{31})$
$-X_{-11}$	$2X_{-11}$	0	0	$2X_{-12}$	0	0	0	$2X_{-13}$	0
$-X_{-12}$	X_{-12}	X_{-12}	0	X_{-22}	X_{-11}	X_{-13}	0	X_{-23}	0
$-X_{-13}$	X_{-13}	0	X_{-13}	X_{-23}	0	0	X_{-12}	0	0
$-X_{-22}$	0	$2X_{-22}$	0	0	$2X_{-12}$	$2X_{-23}$	0	X_{-33}	X_{-11}
$-X_{-23}$	0	X_{-23}	X_{-23}	0	X_{-13}	X_{-33}	X_{-22}	0	X_{-12}
$-X_{-33}$	0	0	$2X_{-33}$	0	0	0	$2X_{-23}$	0	$2X_{-13}$

TABLE 2. The adjoint actions of $\mathfrak{k}_{\mathbf{C}}$ on the basis $\{X_{-ij}\}_{1\leq i\leq j\leq 3}$ of $\mathfrak{p}_-$

Comparing the actions in above tables with the actions in Proposition 3.1, we have the assertion. □

Remark 3.5. *The above lemma tells that $\mathfrak{p}_+$ and $\mathfrak{p}_-$ are simple K-modules with the markings $\{X_{+ij}\}_{1\leq i\leq j\leq 3}$ and $\{X_{-ij}\}_{1\leq i\leq j\leq 3}$, respectively. From now on we always take these markings for $\mathfrak{p}_\pm$.*

4. CLEBSCH-GORDAN COEFFICIENTS FOR THE REPRESENTATIONS OF $\mathfrak{gl}(3, \mathbf{C})$ WITH RESPECT TO THE MONOMIAL BASIS

In the later sections, we need irreducible decompositions of the tensor products $V \otimes_{\mathbf{C}} \mathfrak{p}_+$ and $V \otimes_{\mathbf{C}} \mathfrak{p}_-$ as K-modules for a K-type (τ, V) of $\pi_{(\sigma,\nu)}$. Since $\mathfrak{p}_+ \simeq V_{2\mathbf{e}_1}$, $\mathfrak{p}_- \simeq V_{-2\mathbf{e}_3}$ and $\mathfrak{k}_{\mathbf{C}} \simeq \mathfrak{gl}(3, \mathbf{C})$, it suffices to consider the irreducible decomposition of $V_\lambda \otimes_{\mathbf{C}} V_{2\mathbf{e}_1}$ and $V_\lambda \otimes_{\mathbf{C}} V_{-2\mathbf{e}_3}$ as $\mathfrak{gl}(3, \mathbf{C})$-modules for arbitrary dominant weight λ. In this section, we take the marking $\{f(M)\}_{M\in G(\lambda)}$ for a simple K-module V_λ.

4.1. **The irreducible decomposition of $V_\lambda \otimes_{\mathbf{C}} V_{\mathbf{e}_1}$.** Generically the tensor product $V_\lambda \otimes_{\mathbf{C}} V_{\mathbf{e}_1}$ has three irreducible components: $V_{\lambda+\mathbf{e}_1}$, $V_{\lambda+\mathbf{e}_2}$ and $V_{\lambda+\mathbf{e}_3}$. If $\lambda+\mathbf{e}_i$ $(i=2,3)$ is not dominant, the corresponding irreducible component does not occur.

For $1 \le i \le 3$, let $i^\lambda_{\mathbf{e}_i}$ be a non-zero generator of $\mathrm{Hom}_K(V_{\lambda+\mathbf{e}_i}, V_\lambda \otimes_{\mathbf{C}} V_{\mathbf{e}_1})$, which is unique up to scalar multiple if $V_{\lambda+\mathbf{e}_i}$ is non-zero. Our purpose of this subsection is to give explicit expressions of these injectors $i^\lambda_{\mathbf{e}_1}$, $i^\lambda_{\mathbf{e}_2}$ and $i^\lambda_{\mathbf{e}_3}$ in terms of the monomial basis. For this purpose, we prepare following equations of the functions of G-patterns.

Lemma 4.1. *(i) We have another expressions of $C_1(M)$, $\bar{C}_1(M)$ and $C_2(M)$, which is suitable for computation:*

$$\begin{aligned} C_1(M) &= m_{11} - m_{22} + \delta(M)\chi_-(M) \\ &= m_{12} - m_{23} - \delta(M)\chi_+(M), \end{aligned} \tag{4.1}$$

$$\begin{aligned} \bar{C}_1(M) &= m_{23} - m_{22} + \delta(M)\chi_-(M) \\ &= m_{12} - m_{11} - \delta(M)\chi_+(M), \end{aligned} \tag{4.2}$$

$$\begin{aligned} C_2(M) &= (m_{12} - m_{23})(m_{12} - m_{11}) - (m_{12} - m_{22})\delta(M)\chi_+(M) \\ &= (m_{11} - m_{22})(m_{23} - m_{22}) + (m_{12} - m_{22})\delta(M)\chi_-(M). \end{aligned} \tag{4.3}$$

(ii) We have relations of the values of the functions δ and $\chi^{(r)}_\pm$ for another G-patterns as follows:

$$\delta\left(M \begin{pmatrix} i_{13}\ i_{23}\ i_{33} \\ i_{12}\ i_{22} \\ i_{11} \end{pmatrix} [-l]\right) = \delta(M) + d, \tag{4.4}$$

$$\chi^{(r)}_+(M \begin{pmatrix} i_{13}\ i_{23}\ i_{33} \\ i_{12}\ i_{22} \\ i_{11} \end{pmatrix} [-l]) = \chi^{(r-d)}_+(M), \tag{4.5}$$

$$\chi^{(r)}_-(M \begin{pmatrix} i_{13}\ i_{23}\ i_{33} \\ i_{12}\ i_{22} \\ i_{11} \end{pmatrix} [-l]) = \chi^{(r+d)}_-(M). \tag{4.6}$$

Here $d = \delta \begin{pmatrix} i_{13}\ i_{23}\ i_{33} \\ i_{12}\ i_{22} \\ i_{11} \end{pmatrix}$.

(iii) We have shift relations of $\chi^{(r)}_\pm(M)$ as follows:

$$(\delta(M) - r)\chi^{(r)}_+(M) = (\delta(M) - r)\chi^{(r-1)}_+(M), \tag{4.7}$$

$$(\delta(M) + r)\chi^{(r)}_-(M) = (\delta(M) + r)\chi^{(r-1)}_-(M), \tag{4.8}$$

$$\chi^{(r)}_+(M) + \chi^{(-r-1)}_-(M) = 1, \tag{4.9}$$

$$\chi^{(r_1)}_+(M)\chi^{(r_2)}_-(M) = 0 \quad \text{if} \quad r_1 + r_2 > -2, \tag{4.10}$$

$$\chi^{(r_1)}_+(M)\chi^{(r_2)}_+(M) = \chi^{(r_1)}_+(M) \quad \text{if} \quad r_1 > r_2, \tag{4.11}$$

$$\chi^{(r_1)}_-(M)\chi^{(r_2)}_-(M) = \chi^{(r_1)}_-(M) \quad \text{if} \quad r_1 > r_2. \tag{4.12}$$

(iv) We have convenient relations of $C_1(M)\chi^{(r)}_\pm(M)$ and $\bar{C}_1(M)\chi^{(r)}_\pm(M)$ as follows:

$$C_1(M)\chi^{(r)}_+(M) = (m_{11} - m_{22})\chi^{(r)}_+(M) \qquad (r \ge -1), \tag{4.13}$$

$$C_1(M)\chi^{(r)}_-(M) = (m_{12} - m_{23})\chi^{(r)}_-(M) \qquad (r \ge -1), \tag{4.14}$$

$$\bar{C}_1(M)\chi^{(r)}_+(M) = (m_{23} - m_{22})\chi^{(r)}_+(M) \qquad (r \ge -1), \tag{4.15}$$

$$\bar{C}_1(M)\chi^{(r)}_-(M) = (m_{12} - m_{11})\chi^{(r)}_-(M) \qquad (r \ge -1). \tag{4.16}$$

Proof. We can easily check these equations by direct computation. □

The explicit expressions of the injectors $i^\lambda_{\mathbf{e}_1}$, $i^\lambda_{\mathbf{e}_2}$ and $i^\lambda_{\mathbf{e}_3}$ are given as follows.

Proposition 4.2. *For $1 \le i \le 3$, the image of the monomial basis by the injector $i^{\lambda}_{\mathbf{e}_i} \colon V_{\lambda+\mathbf{e}_i} \to V_{\lambda} \otimes_{\mathbf{C}} V_{\mathbf{e}_1}$ is given by the form*

$$i^{\lambda}_{\mathbf{e}_i}(f(M)) = \sum_{0 \le j \le k \le 1} \left\{ \sum_{l=0}^{r_{[i;jk]}} c^{\lambda}_{[i;jk;l]}(M)\, f\left(M \left(\begin{smallmatrix} -\mathbf{e}_i \\ 0\ -k \\ -j \end{smallmatrix} \right) [-l] \right) \right\} \otimes f \left(\begin{smallmatrix} 1\,0\,0 \\ k\,0 \\ j \end{smallmatrix} \right)$$

for a G-pattern M of type $\lambda + \mathbf{e}_i$. In the right hand side of the above equation, we put $f(M') = 0$ if M' is a triangular array which does not satisfy the condition (3.1) of G-patterns.

The explicit expressions of the coefficients are given by following formulas.

Formula 1: *The coefficients of the injector $i^{\lambda}_{\mathbf{e}_1} \colon V_{\lambda+\mathbf{e}_1} \to V_{\lambda} \otimes_{\mathbf{C}} V_{\mathbf{e}_1}$ are given as follows: $(r_{[1;11]}, r_{[1;01]}, r_{[1;00]}) = (1, 2, 1)$ and*

$$\begin{aligned}
&c^{\lambda}_{[1;11;0]}(M) = (m_{13} - m_{12})(m_{22} - m_{33}), && c^{\lambda}_{[1;11;1]}(M) = -\bar{E}(M), \\
&c^{\lambda}_{[1;01;0]}(M) = -(m_{13} - m_{12})(m_{22} - m_{33}), && c^{\lambda}_{[1;01;1]}(M) = \bar{F}(M), \\
&c^{\lambda}_{[1;01;2]}(M) = -C_2(M)\chi_{+}(M), && c^{\lambda}_{[1;00;0]}(M) = -(m_{13} - m_{12})(m_{13} - m_{22} + 1), \\
&c^{\lambda}_{[1;00;1]}(M) = C_2(M).
\end{aligned}$$

Here

$$\begin{aligned}
&\bar{E}(M) = C_1(M)\{m_{13} - m_{33} + 1 - \bar{C}_1(M)\}, \\
&\bar{F}(M) = -C_2(M) - \chi_{+}(M)\{(m_{13} - m_{12})(m_{22} - m_{33}) + (m_{13} - m_{33} + 1)\delta(M)\}.
\end{aligned}$$

Formula 2: *The coefficients of the injector $i^{\lambda}_{\mathbf{e}_2} \colon V_{\lambda+\mathbf{e}_2} \to V_{\lambda} \otimes_{\mathbf{C}} V_{\mathbf{e}_1}$ are given as follows: $(r_{[2;11]}, r_{[2;01]}, r_{[2;00]}) = (1, 1, 1)$ and*

$$\begin{aligned}
&c^{\lambda}_{[2;11;0]}(M) = m_{22} - m_{33}, && c^{\lambda}_{[2;11;1]}(M) = -\bar{D}(M)\chi_{-}(M), \\
&c^{\lambda}_{[2;01;0]}(M) = -(m_{22} - m_{33}), && c^{\lambda}_{[2;01;1]}(M) = \bar{C}_1(M), \\
&c^{\lambda}_{[2;00;0]}(M) = -(m_{23} - m_{22}), && c^{\lambda}_{[2;00;1]}(M) = -\bar{C}_1(M)\chi_{-}(M).
\end{aligned}$$

Here $\bar{D}(M) = -m_{22} + m_{33} + \delta(M)$.

Formula 3: *The coefficients of the injector $i^{\lambda}_{\mathbf{e}_3} \colon V_{\lambda+\mathbf{e}_3} \to V_{\lambda} \otimes_{\mathbf{C}} V_{\mathbf{e}_1}$ are given as follows: $(r_{[2;11]}, r_{[2;01]}, r_{[2;00]}) = (0, 1, 0)$ and*

$$c^{\lambda}_{[3;11;0]}(M) = 1, \qquad c^{\lambda}_{[3;01;0]}(M) = -1, \qquad c^{\lambda}_{[3;01;1]}(M) = -\chi_{+}(M), \qquad c^{\lambda}_{[3;00;0]}(M) = 1.$$

Proof. For $1 \le i \le 3$, let $i^{\lambda}_{\mathbf{e}_i}$ be a linear map which is defined by the equations in the statement of this proposition. In order to prove that $i^{\lambda}_{\mathbf{e}_i}$ is a K-homomorphism, it suffices to check the actions of the basis E_{mm} $(1 \le m \le 3)$ of Cartan subalgebra and simple root vectors $E_{n\,n+1},\ E_{n+1\,n}$ $(n = 1, 2)$ by direct computation.

Since

$$\begin{aligned}
E_{mm}(f(M_1) \otimes f(M_2)) &= (E_{mm} f(M_1)) \otimes f(M_2) + f(M_1) \otimes (E_{mm} f(M_2)) \\
&= (\gamma^{M_1}_{m} + \gamma^{M_2}_{m}) f(M_1) \otimes f(M_2),
\end{aligned}$$

we easily check $E_{mm} \circ i^{\lambda}_{\mathbf{e}_i}(f(M)) = i^{\lambda}_{\mathbf{e}_i} \circ E_{mm}(f(M))$ for $1 \le m \le 3$ and a G-pattern M of type $\lambda + e_i$.

Therefore the essential computation is those of simple root vectors. We have to confirm that $E_{mn} \circ i^{\lambda}_{\mathbf{e}_i} = i^{\lambda}_{\mathbf{e}_i} \circ E_{mn}$ for four simple root vectors E_{mn} $(|m - n| = 1)$ and each $i = 1, 2, 3$. We set $c^{\lambda}_{[i;jk;l]}(M) = 0$ if $l > r_{[i;jk]}$ or $l < 0$.

First, we compute the image of the monomial basis $f(M)$ by $i^{\lambda}_{\mathbf{e}_i} \circ E_{mn}$. By using the equations in Proposition 3.1, we have

$$i^{\lambda}_{\mathbf{e}_i}(E_{12} f(M)) = (m_{12} - m_{11})\, i^{\lambda}_{\mathbf{e}_i}\left(f\left(M \left(\begin{smallmatrix} 0\,0 \\ 1 \end{smallmatrix} \right)\right)\right) + (m_{23} - m_{22})\chi_{+}(M)\, i^{\lambda}_{\mathbf{e}_i}\left(f\left(M \left(\begin{smallmatrix} 0\,0 \\ 1 \end{smallmatrix} \right) [-1]\right)\right)$$

$$= (m_{12} - m_{11}) \sum_{0 \le j \le k \le 1} \left\{ \sum_{l=0}^{r_{[i;jk]}} c^{\lambda}_{[i;jk;l]} \left(M \left(\begin{smallmatrix} 0\ 0 \\ 1 \end{smallmatrix} \right) \right) f \left(M \left(\begin{smallmatrix} -\mathbf{e}_i \\ 0\ -k \\ 1-j \end{smallmatrix} \right) [-l] \right) \right\} \otimes f \left(\begin{smallmatrix} 1\ 0\ 0 \\ k\ 0 \\ j \end{smallmatrix} \right)$$

$$+ (m_{23} - m_{22}) \chi_+(M) \sum_{0 \le j \le k \le 1} \left\{ \sum_{l=0}^{r_{[i;jk]}} c^{\lambda}_{[i;jk;l]} \left(M \left(\begin{smallmatrix} 0\ 0 \\ 1 \end{smallmatrix} \right) [-1] \right) f \left(M \left(\begin{smallmatrix} -\mathbf{e}_i \\ 0\ -k \\ 1-j \end{smallmatrix} \right) [-l-1] \right) \right\}$$

$$\otimes f \left(\begin{smallmatrix} 1\ 0\ 0 \\ k\ 0 \\ j \end{smallmatrix} \right)$$

$$= \sum_{0 \le j \le k \le 1} \left\{ \sum_{l=0}^{r_{[i;jk]}+1} A^{\lambda}_{[12;i;jk;l]}(M) f \left(M \left(\begin{smallmatrix} -\mathbf{e}_i \\ 0\ -k \\ 1-j \end{smallmatrix} \right) [-l] \right) \right\} \otimes f \left(\begin{smallmatrix} 1\ 0\ 0 \\ k\ 0 \\ j \end{smallmatrix} \right),$$

where

$$\begin{aligned} A^{\lambda}_{[12;i;jk;l]}(M) = & (m_{12} - m_{11}) c^{\lambda}_{[i;jk;l]} \left(M \left(\begin{smallmatrix} 0\ 0 \\ 1 \end{smallmatrix} \right) \right) \\ & + (m_{23} - m_{22}) \chi_+(M) c^{\lambda}_{[i;jk;l-1]} \left(M \left(\begin{smallmatrix} 0\ 0 \\ 1 \end{smallmatrix} \right) [-1] \right). \end{aligned}$$

Similarly, we obtain the following equations:

$$i^{\lambda}_{\mathbf{e}_i}(E_{21} f(M)) = \sum_{0 \le j \le k \le 1} \left\{ \sum_{l=0}^{r_{[i;jk]}+1} A^{\lambda}_{[21;i;jk;l]}(M) f \left(M \left(\begin{smallmatrix} -\mathbf{e}_i \\ 0\ -k \\ -1-j \end{smallmatrix} \right) [-l] \right) \right\} \otimes f \left(\begin{smallmatrix} 1\ 0\ 0 \\ k\ 0 \\ j \end{smallmatrix} \right),$$

where

$$\begin{aligned} A^{\lambda}_{[21;i;jk;l]}(M) = & (m_{11} - m_{22}) c^{\lambda}_{[i;jk;l]} \left(M \left(\begin{smallmatrix} 0\ 0 \\ -1 \end{smallmatrix} \right) \right) \\ & + (m_{12} - m_{23}) \chi_-(M) c^{\lambda}_{[i;jk;l-1]} \left(M \left(\begin{smallmatrix} 0\ 0 \\ -1 \end{smallmatrix} \right) [-1] \right), \end{aligned}$$

$$i^{\lambda}_{\mathbf{e}_i}(E_{23} f(M)) = \sum_{0 \le j \le k \le 1} \left\{ \sum_{l=0}^{r_{[i;jk]}+1} A^{\lambda}_{[23;i;jk;l]}(M) f \left(M \left(\begin{smallmatrix} -\mathbf{e}_i \\ 1\ -k \\ -j \end{smallmatrix} \right) [-l] \right) \right\} \otimes f \left(\begin{smallmatrix} 1\ 0\ 0 \\ k\ 0 \\ j \end{smallmatrix} \right),$$

where

$$\begin{aligned} A^{\lambda}_{[23;i;jk;l]}(M) = & (m_{13} - m_{12}) c^{\lambda}_{[i;jk;l]} \left(M \left(\begin{smallmatrix} 1\ 0 \\ 0 \end{smallmatrix} \right) \right) \\ & + \{m_{13} - m_{12} - \delta(M)\} \chi_-(M) c^{\lambda}_{[i;jk;l-1]} \left(M \left(\begin{smallmatrix} 1\ 0 \\ 0 \end{smallmatrix} \right) [-1] \right), \end{aligned}$$

and

$$i^{\lambda}_{\mathbf{e}_i}(E_{32} f(M)) = \sum_{0 \le j \le k \le 1} \left\{ \sum_{l=0}^{r_{[i;jk]}+1} A^{\lambda}_{[32;i;jk;l]}(M) f \left(M \left(\begin{smallmatrix} -\mathbf{e}_i \\ 0\ -1-k \\ -j \end{smallmatrix} \right) [-l] \right) \right\} \otimes f \left(\begin{smallmatrix} 1\ 0\ 0 \\ k\ 0 \\ j \end{smallmatrix} \right),$$

where

$$\begin{aligned} A^{\lambda}_{[32;i;jk;l]}(M) = & (m_{22} - m_{33}) c^{\lambda}_{[i;jk;l]} \left(M \left(\begin{smallmatrix} 0\ -1 \\ 0 \end{smallmatrix} \right) \right) \\ & + \{m_{22} - m_{33} + \delta(M)\} \chi_+(M) c^{\lambda}_{[i;jk;l-1]} \left(M \left(\begin{smallmatrix} 0\ -1 \\ 0 \end{smallmatrix} \right) [-1] \right). \end{aligned}$$

Next, we compute the image of the the monomial basis $f(M)$ by $E_{mn} \circ i^{\lambda}_{\mathbf{e}_i}$ as follows:

$$E_{12} \circ i^{\lambda}_{\mathbf{e}_i}(f(M)) = \sum_{0 \le j \le k \le 1} \sum_{l=0}^{r_{[i;jk]}} c^{\lambda}_{[i;jk;l]}(M) E_{12} \left\{ f \left(M \left(\begin{smallmatrix} -\mathbf{e}_i \\ 0\ -k \\ -j \end{smallmatrix} \right) [-l] \right) \otimes f \left(\begin{smallmatrix} 1\ 0\ 0 \\ k\ 0 \\ j \end{smallmatrix} \right) \right\}$$

$$= \sum_{0 \le j \le k \le 1} \left\{ \sum_{l=0}^{r_{[i;jk]}} (m_{12} - m_{11} - l + j) c^{\lambda}_{[i;jk;l]}(M) f \left(M \left(\begin{smallmatrix} -\mathbf{e}_i \\ 0\ -k \\ 1-j \end{smallmatrix} \right) [-l] \right) \right.$$

$$+ (m_{23} - m_{22} + k - l - \delta_{2i}) \chi_+ \left(M \left(\begin{smallmatrix} -\mathbf{e}_i \\ 0\ -k \\ -j \end{smallmatrix} \right) [-l] \right)$$

$$
\begin{aligned}
&\qquad \times c^{\lambda}_{[i;jk;l]}(M)\, f\left(M\left(\begin{smallmatrix} -\mathbf{e}_i \\ 0\ -k \\ 1-j \end{smallmatrix}\right)[-l-1]\right)\Bigg\}\otimes f\left(\begin{smallmatrix} 1\,0\,0 \\ k\ 0 \\ j \end{smallmatrix}\right) \\
&\quad + \left\{\sum_{l=0}^{2} c^{\lambda}_{[i;01;l]}(M)\, f\left(M\left(\begin{smallmatrix} -\mathbf{e}_i \\ 0\ -1 \\ 0 \end{smallmatrix}\right)[-l]\right)\right\}\otimes f\left(\begin{smallmatrix} 1\,0\,0 \\ 1\ 0 \\ 1 \end{smallmatrix}\right) \\
&= \sum_{0\le j\le k\le 1}\left\{\sum_{l=0}^{r_{[i;jk]}+1} B^{\lambda}_{[12;i;jk;l]}(M)\, f\left(M\left(\begin{smallmatrix} -\mathbf{e}_i \\ 0\ -k \\ 1-j \end{smallmatrix}\right)[-l]\right)\right\}\otimes f\left(\begin{smallmatrix} 1\,0\,0 \\ k\ 0 \\ j \end{smallmatrix}\right),
\end{aligned}
$$

where

$$
\begin{aligned}
B^{\lambda}_{[12;i;jk;l]}(M) =&(m_{12}-m_{11}-l+j)c^{\lambda}_{[i;jk;l]}(M) \\
&+(m_{23}-m_{22}+k-l+1-\delta_{2i})\chi_{+}^{(k-j-\delta_{2i})}(M)c^{\lambda}_{[i;jk;l-1]}(M) \\
&+\begin{cases} c^{\lambda}_{[i;01;l]}(M) & \text{if } \ (j,k)=(1,1), \\ 0 & \text{otherwise.} \end{cases}
\end{aligned}
$$

Here we use the relation 4.5 in Lemma 4.1.

Similarly, we obtain the following equations:

$$
E_{21}\circ i^{\lambda}_{\mathbf{e}_i}(f(M)) = \sum_{0\le j\le k\le 1}\left\{\sum_{l=0}^{r_{[i;jk]}+1} B^{\lambda}_{[i;jk;l]}(M)\, f\left(M\left(\begin{smallmatrix} -\mathbf{e}_i \\ 0\ -k \\ -1-j \end{smallmatrix}\right)[-l]\right)\right\}\otimes f\left(\begin{smallmatrix} 1\,0\,0 \\ k\ 0 \\ j \end{smallmatrix}\right),
$$

where

$$
\begin{aligned}
B^{\lambda}_{[21;i;jk;l]}(M) =&(m_{11}-m_{22}+k-l-j)c^{\lambda}_{[i;jk;l]}(M) \\
&+(m_{12}-m_{23}-l+1+\delta_{2i})\chi_{-}^{(-k+j+\delta_{2i})}(M)c^{\lambda}_{[i;jk;l-1]}(M) \\
&+\begin{cases} c^{\lambda}_{[i;11;l]}(M) & \text{if } \ (j,k)=(0,1), \\ 0 & \text{otherwise,} \end{cases}
\end{aligned}
$$

$$
E_{23}\circ i^{\lambda}_{\mathbf{e}_i}(f(M)) = \sum_{0\le j\le k\le 1}\left\{\sum_{l=0}^{r_{[i;jk]}+1} B^{\lambda}_{[23;i;jk;l]}(M)\, f\left(M\left(\begin{smallmatrix} -\mathbf{e}_i \\ 1\ -k \\ -j \end{smallmatrix}\right)[-l]\right)\right\}\otimes f\left(\begin{smallmatrix} 1\,0\,0 \\ k\ 0 \\ j \end{smallmatrix}\right),
$$

where

$$
\begin{aligned}
B^{\lambda}_{[23;i;jk;l]}(M) =&(m_{13}-m_{12}+l-\delta_{1i})c^{\lambda}_{[i;jk;l]}(M) \\
&+\{m_{13}-m_{12}-\delta(M)+k-j+l-1-\delta_{1i}-\delta_{2i}\} \\
&\quad\times\chi_{-}^{(-k+j+\delta_{2i})}(M)c^{\lambda}_{[i;jk;l-1]}(M) \\
&+\begin{cases} c^{\lambda}_{[i;00;l-1]}(M) & \text{if } \ (j,k)=(0,1), \\ 0 & \text{otherwise,} \end{cases}
\end{aligned}
$$

and

$$
E_{32}\circ i^{\lambda}_{\mathbf{e}_i}(f(M)) = \sum_{0\le j\le k\le 1}\left\{\sum_{l=0}^{r_{[i;jk]}+1} B^{\lambda}_{[32;i;jk;l]}(M)\, f\left(M\left(\begin{smallmatrix} -\mathbf{e}_i \\ 0\quad -1-k \\ -j \end{smallmatrix}\right)[-l]\right)\right\}\otimes f\left(\begin{smallmatrix} 1\,0\,0 \\ k\ 0 \\ j \end{smallmatrix}\right),
$$

where

$$
\begin{aligned}
B^{\lambda}_{[32;i;jk;l]}(M) =&(m_{22}-m_{33}-k+l+\delta_{3i})c^{\lambda}_{[i;jk;l]}(M) \\
&+\{m_{22}-m_{33}+\delta(M)+j-2k+l-1+\delta_{2i}+\delta_{3i}\}
\end{aligned}
$$

$$\times\, \chi_+^{(k-j-\delta_{2i})}(M)c^{\lambda}_{[i;jk;l-1]}(M)$$
$$+\left\{\begin{array}{ll} c^{\lambda}_{[i;01;l]}(M) & \text{if } \ (j,k)=(0,0),\\ 0 & \text{otherwise.}\end{array}\right.$$

Here we use the relations in Lemma 4.1 (ii).

In order to complete the proof, we check the equations

$$A^{\lambda}_{[mn;i;jk;l]}(M) = B^{\lambda}_{[mn;i;jk;l]}(M) \tag{4.17}$$

by direct computation.

First, we check the equations (4.17) for $i=3$, that is, the case of formula 3.

- the proof of $E_{12}\circ i^{\lambda}_{\mathbf{e}_3} = i^{\lambda}_{\mathbf{e}_3}\circ E_{12}$.

We have

$$\begin{aligned}
A^{\lambda}_{[12;3;11;0]}(M) =& m_{12}-m_{11},\\
A^{\lambda}_{[12;3;11;1]}(M) =& (m_{23}-m_{22})\chi_+(M),\\
A^{\lambda}_{[12;3;01;0]}(M) =& -(m_{12}-m_{11}),\\
A^{\lambda}_{[12;3;01;1]}(M) =& -(m_{12}-m_{11})\chi_+\left(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)\right)-(m_{23}-m_{22})\chi_+(M),\\
A^{\lambda}_{[12;3;01;2]}(M) =& -(m_{23}-m_{22})\chi_+(M)\chi_+\left(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)[-1]\right),\\
A^{\lambda}_{[12;3;00;0]}(M) =& m_{12}-m_{11},\\
A^{\lambda}_{[12;3;00;1]}(M) =& (m_{23}-m_{22})\chi_+(M),
\end{aligned}$$

and

$$\begin{aligned}
B^{\lambda}_{[12;3;11;0]}(M) =& (m_{12}-m_{11}+1)-1,\\
B^{\lambda}_{[12;3;11;1]}(M) =& (m_{23}-m_{22}+1)\chi_+(M)-\chi_+(M),\\
B^{\lambda}_{[12;3;01;0]}(M) =& -(m_{12}-m_{11}),\\
B^{\lambda}_{[12;3;01;1]}(M) =& -(m_{12}-m_{11}-1)\chi_+(M)-(m_{23}-m_{22}+1)\chi_+^{(1)}(M),\\
B^{\lambda}_{[12;3;01;2]}(M) =& -(m_{23}-m_{22})\chi_+^{(1)}(M)\chi_+(M),\\
B^{\lambda}_{[12;3;00;0]}(M) =& m_{12}-m_{11},\\
B^{\lambda}_{[12;3;00;1]}(M) =& (m_{23}-m_{22})\chi_+(M).
\end{aligned}$$

By direct computation, we have

$$\begin{aligned}
A^{\lambda}_{[12;3;01;2]}(M)-B^{\lambda}_{[12;3;01;2]}(M) &= (m_{23}-m_{22})\left(\chi_+^{(1)}(M)\chi_+(M)-\chi_+(M)\chi_+\left(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)[-1]\right)\right)\\
&= (m_{23}-m_{22})(\chi_+^{(1)}(M)-\chi_+^{(1)}(M))=0,
\end{aligned}$$

$$\begin{aligned}
A^{\lambda}_{[12;3;01;1]}(M)-B^{\lambda}_{[12;3;01;1]}(M) &= -(m_{12}-m_{11})\chi_+\left(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)\right)-(m_{23}-m_{22})\chi_+(M)\\
&\quad\ +(m_{12}-m_{11}-1)\chi_+(M)+(m_{23}-m_{22}+1)\chi_+^{(1)}(M)\\
&= (\delta(M)-1)\chi_+(M)-(\delta(M)-1)\chi_+^{(1)}(M)=0.
\end{aligned}$$

Hence we obtain the equations (4.17) for $(j,k,l)=(0,1,1)$ and $(0,1,2)$. Here we use the relations (4.5) and (4.7).

It is trivial that the equations (4.17) hold for other (j,k,l).

- the proof of $E_{21}\circ i^{\lambda}_{\mathbf{e}_3} = i^{\lambda}_{\mathbf{e}_3}\circ E_{21}$.

We have

$$A^{\lambda}_{[21;3;11;0]}(M) = m_{11}-m_{22},$$

$$\begin{aligned}
A^{\lambda}_{[21;3;11;1]}(M) =&(m_{12}-m_{23})\chi_{-}(M),\\
A^{\lambda}_{[21;3;01;0]}(M) =&-(m_{11}-m_{22}),\\
A^{\lambda}_{[21;3;01;1]}(M) =&-(m_{11}-m_{22})\chi_{+}\left(M\left(\begin{smallmatrix}0\;0\\-1\end{smallmatrix}\right)\right)-(m_{12}-m_{23})\chi_{-}(M),\\
A^{\lambda}_{[21;3;01;2]}(M) =&-(m_{12}-m_{23})\chi_{-}(M)\chi_{+}\left(M\left(\begin{smallmatrix}0\;0\\-1\end{smallmatrix}\right)[-1]\right),\\
A^{\lambda}_{[21;3;00;0]}(M) =&m_{11}-m_{22},\\
A^{\lambda}_{[21;3;00;1]}(M) =&(m_{12}-m_{23})\chi_{-}(M),
\end{aligned}$$

and

$$\begin{aligned}
B^{\lambda}_{[21;3;11;0]}(M) =&m_{11}-m_{22},\\
B^{\lambda}_{[21;3;11;1]}(M) =&(m_{12}-m_{23})\chi_{-}(M),\\
B^{\lambda}_{[21;3;01;0]}(M) =&-(m_{11}-m_{22}+1)+1,\\
B^{\lambda}_{[21;3;01;1]}(M) =&-(m_{11}-m_{22})\chi_{+}(M)-(m_{12}-m_{23})\chi_{-}^{(-1)}(M),\\
B^{\lambda}_{[21;3;01;2]}(M) =&-(m_{12}-m_{23}-1)\chi_{-}^{(-1)}(M)\chi_{+}(M),\\
B^{\lambda}_{[21;3;00;0]}(M) =&m_{11}-m_{22},\\
B^{\lambda}_{[21;3;00;1]}(M) =&(m_{12}-m_{23})\chi_{-}(M).
\end{aligned}$$

We have

$$\begin{aligned}
A^{\lambda}_{[21;3;01;1]}(M) &= -(m_{11}-m_{22})\chi_{+}^{(-1)}(M)-(m_{12}-m_{23})(1-\chi_{+}^{(-1)}(M))\\
&= -m_{12}+m_{23}+\delta(M)\chi_{+}^{(-1)}(M)\\
&= -m_{12}+m_{23}+\delta(M)\chi_{+}(M),\\
B^{\lambda}_{[21;3;01;1]}(M) &= -(m_{11}-m_{22})\chi_{+}(M)-(m_{12}-m_{23})(1-\chi_{+}(M))\\
&= -m_{12}+m_{23}+\delta(M).
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[21;3;01;1]}(M) = B^{\lambda}_{[21;3;01;1]}(M)$. Here we use the relations (4.5), (4.7) and (4.9).

We have

$$\begin{aligned}
A^{\lambda}_{[21;3;01;2]}(M) &= -(m_{12}-m_{23})\chi_{-}(M)\chi_{+}^{(-1)}(M)=0,\\
B^{\lambda}_{[21;3;01;2]}(M) &= -(m_{12}-m_{23}-1)\chi_{-}^{(-1)}(M)\chi_{+}(M)=0.
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[21;3;01;2]}(M) = B^{\lambda}_{[21;3;01;2]}(M)$. Here we use the relations (4.5) and (4.10).

It is trivial that the equations (4.17) hold for other (j, k, l).

- the proof of $E_{23}\circ i^{\lambda}_{\mathbf{e}_3} = i^{\lambda}_{\mathbf{e}_3}\circ E_{23}$.

We have

$$\begin{aligned}
A^{\lambda}_{[23;3;11;0]}(M) =&(m_{13}-m_{12}),\\
A^{\lambda}_{[23;3;11;1]}(M) =&\{m_{13}-m_{12}-\delta(M)\}\chi_{-}(M),\\
A^{\lambda}_{[23;3;01;0]}(M) =&-(m_{13}-m_{12}),\\
A^{\lambda}_{[23;3;01;1]}(M) =&-(m_{13}-m_{12})\chi_{+}\left(M\left(\begin{smallmatrix}1\;0\\0\end{smallmatrix}\right)\right)-\{m_{13}-m_{12}-\delta(M)\}\chi_{-}(M),\\
A^{\lambda}_{[23;3;01;2]}(M) =&-\{m_{13}-m_{12}-\delta(M)\}\chi_{-}(M)\chi_{+}\left(M\left(\begin{smallmatrix}1\;0\\0\end{smallmatrix}\right)[-1]\right),\\
A^{\lambda}_{[23;3;00;0]}(M) =&m_{13}-m_{12},\\
A^{\lambda}_{[23;3;00;1]}(M) =&\{m_{13}-m_{12}-\delta(M)\}\chi_{-}(M),
\end{aligned}$$

and

$$\begin{aligned}
B^{\lambda}_{[23;3;11;0]}(M) &= m_{13} - m_{12},\\
B^{\lambda}_{[23;3;11;1]}(M) &= \{m_{13} - m_{12} - \delta(M)\}\chi_-(M),\\
B^{\lambda}_{[23;3;01;0]}(M) &= -(m_{13} - m_{12}),\\
B^{\lambda}_{[23;3;01;1]}(M) &= -(m_{13} - m_{12} + 1)\chi_+(M) - \{m_{13} - m_{12} - \delta(M) + 1\}\chi_-^{(-1)}(M) + 1,\\
B^{\lambda}_{[23;3;01;2]}(M) &= -\{m_{13} - m_{12} - \delta(M) + 2\}\chi_-^{(-1)}(M)\chi_+(M),\\
B^{\lambda}_{[23;3;00;0]}(M) &= m_{13} - m_{12},\\
B^{\lambda}_{[23;3;00;1]}(M) &= \{m_{13} - m_{12} - \delta(M)\}\chi_-(M).
\end{aligned}$$

We have

$$\begin{aligned}
A^{\lambda}_{[23;3;01;1]}(M) &= -(m_{13} - m_{12})(\chi_+^{(-1)}(M) + \chi_-(M)) + \delta(M)\chi_-(M)\\
&= -m_{13} + m_{12} + \delta(M)\chi_-(M),\\
B^{\lambda}_{[23;3;01;1]}(M) &= -(m_{13} - m_{12} + 1)(\chi_+(M) + \chi_-^{(-1)}(M)) + 1 + \delta(M)\chi_-^{(-1)}(M)\\
&= -m_{13} + m_{12} + \delta(M)\chi_-(M).
\end{aligned}$$

Hence $A^{\lambda}_{[23;3;01;1]}(M) = B^{\lambda}_{[23;3;01;1]}(M)$. Here we use the relations (4.5) and (4.9).

We have

$$\begin{aligned}
A^{\lambda}_{[23;3;01;2]}(M) &= -\{m_{13} - m_{12} - \delta(M)\}\chi_-(M)\chi_+^{(-1)}(M) = 0,\\
B^{\lambda}_{[23;3;01;2]}(M) &= -\{m_{13} - m_{12} - \delta(M) + 2\}\chi_-^{(-1)}(M)\chi_+(M) = 0.
\end{aligned}$$

Hence $A^{\lambda}_{[23;3;01;2]}(M) = B^{\lambda}_{[23;3;01;2]}(M)$. Here we use the relations (4.5) and (4.10).

It is trivial that the equations (4.17) hold for other (j, k, l).

- the proof of $E_{32} \circ i^{\lambda}_{\mathbf{e}_3} = i^{\lambda}_{\mathbf{e}_3} \circ E_{32}$.

$$\begin{aligned}
A^{\lambda}_{[32;3;11;0]}(M) &= m_{22} - m_{33},\\
A^{\lambda}_{[32;3;11;1]}(M) &= \{m_{22} - m_{33} + \delta(M)\}\chi_+(M),\\
A^{\lambda}_{[32;3;01;0]}(M) &= -(m_{22} - m_{33}),\\
A^{\lambda}_{[32;3;01;1]}(M) &= -(m_{22} - m_{33})\chi_+\left(M\left(\begin{smallmatrix}0\ -1\\0\end{smallmatrix}\right)\right) - \{m_{22} - m_{33} + \delta(M)\}\chi_+(M),\\
A^{\lambda}_{[32;3;01;2]}(M) &= -\{m_{22} - m_{33} + \delta(M)\}\chi_+(M)\chi_+\left(M\left(\begin{smallmatrix}0\ -1\\0\end{smallmatrix}\right)[-1]\right),\\
A^{\lambda}_{[32;3;00;0]}(M) &= (m_{22} - m_{33}),\\
A^{\lambda}_{[32;3;00;1]}(M) &= \{m_{22} - m_{33} + \delta(M)\}\chi_+(M),
\end{aligned}$$

and

$$\begin{aligned}
B^{\lambda}_{[32;3;11;0]}(M) &= m_{22} - m_{33},\\
B^{\lambda}_{[32;3;11;1]}(M) &= \{m_{22} - m_{33} + \delta(M)\}\chi_+(M),\\
B^{\lambda}_{[32;3;01;0]}(M) &= -(m_{22} - m_{33}),\\
B^{\lambda}_{[32;3;01;1]}(M) &= -(m_{22} - m_{33} + 1)\chi_+(M) - \{m_{22} - m_{33} + \delta(M) - 1\}\chi_+^{(1)}(M),\\
B^{\lambda}_{[32;3;01;2]}(M) &= -\{m_{22} - m_{33} + \delta(M)\}\chi_+^{(1)}(M)\chi_+(M),\\
B^{\lambda}_{[32;3;00;0]}(M) &= (m_{22} - m_{33} + 1) - 1,\\
B^{\lambda}_{[32;3;00;1]}(M) &= \{m_{22} - m_{33} + \delta(M) + 1\}\chi_+(M) - \chi_+(M).
\end{aligned}$$

We have

$$\begin{aligned}
&A^{\lambda}_{[32;3;01;1]}(M) - B^{\lambda}_{[32;3;01;1]}(M)\\
&= -(m_{22} - m_{33})\chi_{+}^{(1)}(M) - \{m_{22} - m_{33} + \delta(M)\}\chi_{+}(M)\\
&\quad + (m_{22} - m_{33} + 1)\chi_{+}(M) + \{m_{22} - m_{33} + \delta(M) - 1\}\chi_{+}^{(1)}(M)\\
&= -(\delta(M) - 1)\chi_{+}(M) + (\delta(M) - 1)\chi_{+}^{(1)}(M) = 0.
\end{aligned}$$

Hence we obtain the equation (4.17) for $(j, k, l) = (0, 1, 1)$. Here we use the relations (4.5) and (4.7).

We have

$$\begin{aligned}
&A^{\lambda}_{[32;3;01;2]}(M) - B^{\lambda}_{[32;3;01;2]}(M)\\
&= -\{m_{22} - m_{33} + \delta(M)\}\left(\chi_{+}(M)\chi_{+}\left(M\left(\begin{smallmatrix}0&&-1\\&0&\end{smallmatrix}\right)[-1]\right) - \chi_{+}^{(1)}(M)\chi_{+}(M)\right)\\
&= -\{m_{22} - m_{33} + \delta(M)\}\left(\chi_{+}^{(1)}(M) - \chi_{+}^{(1)}(M)\right) = 0.
\end{aligned}$$

Hence we obtain the equation (4.17) for $(j, k, l) = (0, 1, 2)$. Here we use the relations (4.5) and (4.11).

It is trivial that the equations (4.17) hold for other (j, k, l).

In these computations, we use the relations in Lemma 4.1, frequently. So we use these relations without notice in the proof of formula 1 and formula 2. Next, we check the equations 4.17 for $i = 2$, that is, the case of formula 2.

- the proof of $E_{12} \circ i^{\lambda}_{\mathbf{e}_2} = i^{\lambda}_{\mathbf{e}_2} \circ E_{12}$.

We have

$$\begin{aligned}
A^{\lambda}_{[12;2;11;0]}(M) =&(m_{12} - m_{11})(m_{22} - m_{33}),\\
A^{\lambda}_{[12;2;11;1]}(M) =&-(m_{12} - m_{11})\bar{D}\left(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)\right)\chi_{-}\left(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)\right)\\
&+ (m_{23} - m_{22})\chi_{+}(M)(m_{22} - m_{33} + 1),\\
A^{\lambda}_{[12;2;11;2]}(M) =&-(m_{23} - m_{22})\chi_{+}(M)\bar{D}\left(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)[-1]\right)\chi_{-}\left(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)[-1]\right),\\
A^{\lambda}_{[12;2;01;0]}(M) =&-(m_{12} - m_{11})(m_{22} - m_{33}),\\
A^{\lambda}_{[12;2;01;1]}(M) =&(m_{12} - m_{11})\bar{C}_1\left(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)\right) - (m_{23} - m_{22})\chi_{+}(M)(m_{22} - m_{33} + 1),\\
A^{\lambda}_{[12;2;01;2]}(M) =&(m_{23} - m_{22})\chi_{+}(M)\bar{C}_1\left(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)[-1]\right),\\
A^{\lambda}_{[12;2;00;0]}(M) =&-(m_{12} - m_{11})(m_{23} - m_{22}),\\
A^{\lambda}_{[12;2;00;1]}(M) =&-(m_{12} - m_{11})\bar{C}_1\left(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)\right)\chi_{-}\left(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)\right)\\
&- (m_{23} - m_{22})\chi_{+}(M)(m_{23} - m_{22} - 1),\\
A^{\lambda}_{[12;2;00;2]}(M) =&-(m_{23} - m_{22})\chi_{+}(M)\bar{C}_1\left(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)[-1]\right)\chi_{-}\left(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)[-1]\right),
\end{aligned}$$

and

$$\begin{aligned}
B^{\lambda}_{[12;2;11;0]}(M) =&(m_{12} - m_{11} + 1)(m_{22} - m_{33}) - (m_{22} - m_{33}),\\
B^{\lambda}_{[12;2;11;1]}(M) =&-(m_{12} - m_{11})\bar{D}(M)\chi_{-}(M)\\
&+ (m_{23} - m_{22})\chi_{+}^{(-1)}(M)(m_{22} - m_{33}) + \bar{C}_1(M),\\
B^{\lambda}_{[12;2;11;2]}(M) =&-(m_{23} - m_{22} - 1)\chi_{+}^{(-1)}(M)\bar{D}(M)\chi_{-}(M),\\
B^{\lambda}_{[12;2;01;0]}(M) =&-(m_{12} - m_{11})(m_{22} - m_{33}),\\
B^{\lambda}_{[12;2;01;1]}(M) =&(m_{12} - m_{11} - 1)\bar{C}_1(M) - (m_{23} - m_{22})\chi_{+}(M)(m_{22} - m_{33}),
\end{aligned}$$

$$\begin{aligned}
B^{\lambda}_{[12;2;01;2]}(M) =& (m_{23}-m_{22}-1)\chi_+(M)\bar{C}_1(M),\\
B^{\lambda}_{[12;2;00;0]}(M) =& -(m_{12}-m_{11})(m_{23}-m_{22}),\\
B^{\lambda}_{[12;2;00;1]}(M) =& -(m_{12}-m_{11}-1)\bar{C}_1(M)\chi_-(M)\\
& -(m_{23}-m_{22}-1)\chi_+^{(-1)}(M)(m_{23}-m_{22}),\\
B^{\lambda}_{[12;2;00;2]}(M) =& -(m_{23}-m_{22}-2)\chi_+^{(-1)}(M)\bar{C}_1(M)\chi_-(M).
\end{aligned}$$

We have

$$\begin{aligned}
A^{\lambda}_{[12;2;11;1]}(M) =& -(m_{12}-m_{11})(-m_{22}+m_{33}+\delta(M)-1)(1-\chi_+(M))\\
& +(m_{23}-m_{22})\chi_+(M)(m_{22}-m_{33}+1)\\
=& -(m_{12}-m_{11})(\bar{D}(M)-1)-(m_{12}-m_{11}+m_{22}-m_{33}+1)\delta(M)\chi_+(M),
\end{aligned}$$

$$\begin{aligned}
B^{\lambda}_{[12;2;11;1]}(M) =& -(m_{12}-m_{11})(-m_{22}+m_{33}+\delta(M))(1-\chi_+^{(-1)}(M))\\
& +(m_{23}-m_{22})\chi_+^{(-1)}(M)(m_{22}-m_{33})+(m_{12}-m_{11}-\delta(M)\chi_+(M))\\
=& -(m_{12}-m_{11})(\bar{D}(M)-1)-(m_{12}-m_{11}+m_{22}-m_{33})\delta(M)\chi_+^{(-1)}(M)\\
& -\delta(M)\chi_+(M)\\
=& -(m_{12}-m_{11})(\bar{D}(M)-1)-(m_{12}-m_{11}+m_{22}-m_{33}+1)\delta(M)\chi_+^{(-1)}(M).
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[12;2;11;1]}(M)=B^{\lambda}_{[12;2;11;1]}(M)$.

We have

$$\begin{aligned}
A^{\lambda}_{[12;2;11;2]}(M) =& -(m_{23}-m_{22})\bar{D}\left(M\left(\begin{smallmatrix}0\ 0\\ 1\end{smallmatrix}\right)[-1]\right)\chi_+(M)\chi_-^{(-1)}(M)=0,\\
B^{\lambda}_{[12;2;11;2]}(M) =& -(m_{23}-m_{22}-1)\bar{D}(M)\chi_+^{(-1)}(M)\chi_-(M)=0.
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[12;2;11;2]}(M)=B^{\lambda}_{[12;2;11;2]}(M)$.

We have

$$\begin{aligned}
A^{\lambda}_{[12;2;01;1]}(M) =& (m_{12}-m_{11})\{m_{12}-m_{11}-1-(\delta(M)-1)\chi_+(M)\}\\
& -(m_{23}-m_{22})\chi_+(M)(m_{22}-m_{33}+1)\\
=& (m_{12}-m_{11})(m_{12}-m_{11}-1)\\
& -\chi_+(M)\{(m_{23}-m_{22})(m_{22}-m_{33}+1)+(m_{12}-m_{11})(\delta(M)-1)\},
\end{aligned}$$

$$\begin{aligned}
B^{\lambda}_{[12;2;01;1]}(M) =& (m_{12}-m_{11}-1)(m_{12}-m_{11}-\delta(M)\chi_+(M))\\
& -(m_{23}-m_{22})\chi_+(M)(m_{22}-m_{33})\\
=& (m_{12}-m_{11})(m_{12}-m_{11}-1)\\
& -\chi_+(M)((m_{23}-m_{22})(m_{22}-m_{33})+(m_{12}-m_{11}-1)\delta(M)).
\end{aligned}$$

Therefore

$$A^{\lambda}_{[12;2;01;1]}(M)-B^{\lambda}_{[12;2;01;1]}(M)=-\chi_+(M)\{(m_{23}-m_{22})-(m_{12}-m_{11})+\delta(M)\}=0.$$

Hence we obtain $A^{\lambda}_{[12;2;01;1]}(M)=B^{\lambda}_{[12;2;01;1]}(M)$.

We have

$$\begin{aligned}
A^{\lambda}_{[12;2;01;2]}(M) =& (m_{23}-m_{22})\chi_+(M)\{m_{23}-m_{22}-1+(\delta(M)-1)\chi_-^{(-1)}(M)\}\\
=& (m_{23}-m_{22})(m_{23}-m_{22}-1)\chi_+(M),\\
B^{\lambda}_{[12;2;01;2]}(M) =& (m_{23}-m_{22}-1)(m_{23}-m_{22})\chi_+(M).
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[12;2;01;2]}(M)=B^{\lambda}_{[12;2;01;2]}(M)$.

We have

$$\begin{aligned} A^{\lambda}_{[12;2;00;1]}(M) =& -(m_{12}-m_{11})(m_{12}-m_{11}-1)\chi_{-}^{(-1)}(M) \\ & -(m_{23}-m_{22})(m_{23}-m_{22}-1)\chi_{+}(M) \\ =& -(\bar{C}_1(M))(\bar{C}_1(M)-1)(\chi_{-}^{(-1)}(M)+\chi_{+}(M)) \\ =& -(\bar{C}_1(M))(\bar{C}_1(M)-1), \end{aligned}$$

$$\begin{aligned} B^{\lambda}_{[12;2;00;1]}(M) =& -(m_{12}-m_{11}-1)(m_{12}-m_{11})\chi_{-}(M) \\ & -(m_{23}-m_{22})(m_{23}-m_{22}-1)\chi_{+}^{(-1)}(M) \\ =& -(\bar{C}_1(M))(\bar{C}_1(M)-1)(\chi_{-}(M)+\chi_{+}^{(-1)}(M)) \\ =& -(\bar{C}_1(M))(\bar{C}_1(M)-1). \end{aligned}$$

Hence we obtain $A^{\lambda}_{[12;2;00;1]}(M) = B^{\lambda}_{[12;2;00;1]}(M)$.

We have

$$\begin{aligned} A^{\lambda}_{[12;2;00;2]}(M) =& -(m_{23}-m_{22})\bar{C}_1\left(M\left(\begin{smallmatrix}0\,0\\1\end{smallmatrix}\right)[-1]\right)\chi_{+}(M)\chi_{-}^{(-1)}(M) = 0, \\ B^{\lambda}_{[12;2;00;2]}(M) =& -(m_{23}-m_{22}-2)\bar{C}_1(M)\chi_{+}^{(-1)}(M)\chi_{-}(M) = 0. \end{aligned}$$

Hence we obtain $A^{\lambda}_{[12;2;00;2]}(M) = B^{\lambda}_{[12;2;00;2]}(M)$.

It is trivial that the equations (4.17) hold for $(j,k,l) = (1,1,0),\ (0,1,0)$ and $(0,0,0)$.

- the proof of $E_{21} \circ i^{\lambda}_{\mathbf{e}_2} = i^{\lambda}_{\mathbf{e}_2} \circ E_{21}$.

We have

$$\begin{aligned} A^{\lambda}_{[21;2;11;0]}(M) =&(m_{11}-m_{22})(m_{22}-m_{33}) \\ A^{\lambda}_{[21;2;11;1]}(M) =& -(m_{11}-m_{22})\bar{D}\left(M\left(\begin{smallmatrix}0\,0\\-1\end{smallmatrix}\right)\right)\chi_{-}\left(M\left(\begin{smallmatrix}0\,0\\-1\end{smallmatrix}\right)\right) \\ & +(m_{12}-m_{23})\chi_{-}(M)(m_{22}-m_{33}+1), \\ A^{\lambda}_{[21;2;11;2]}(M) =& -(m_{12}-m_{23})\chi_{-}(M)\bar{D}\left(M\left(\begin{smallmatrix}0\,0\\-1\end{smallmatrix}\right)[-1]\right)\chi_{-}\left(M\left(\begin{smallmatrix}0\,0\\-1\end{smallmatrix}\right)[-1]\right), \\ A^{\lambda}_{[21;2;01;0]}(M) =& -(m_{11}-m_{22})(m_{22}-m_{33}), \\ A^{\lambda}_{[21;2;01;1]}(M) =&(m_{11}-m_{22})\bar{C}_1\left(M\left(\begin{smallmatrix}0\,0\\-1\end{smallmatrix}\right)\right)-(m_{12}-m_{23})\chi_{-}(M)(m_{22}-m_{33}+1), \\ A^{\lambda}_{[21;2;01;2]}(M) =&(m_{12}-m_{23})\chi_{-}(M)\bar{C}_1\left(M\left(\begin{smallmatrix}0\,0\\-1\end{smallmatrix}\right)[-1]\right), \\ A^{\lambda}_{[21;2;00;0]}(M) =& -(m_{11}-m_{22})(m_{23}-m_{22}), \\ A^{\lambda}_{[21;2;00;1]}(M) =& -(m_{11}-m_{22})\bar{C}_1\left(M\left(\begin{smallmatrix}0\,0\\-1\end{smallmatrix}\right)\right)\chi_{-}\left(M\left(\begin{smallmatrix}0\,0\\-1\end{smallmatrix}\right)\right) \\ & -(m_{12}-m_{23})\chi_{-}(M)(m_{23}-m_{22}-1), \\ A^{\lambda}_{[21;2;00;2]}(M) =& -(m_{12}-m_{23})\chi_{-}(M)\bar{C}_1\left(M\left(\begin{smallmatrix}0\,0\\-1\end{smallmatrix}\right)[-1]\right)\chi_{-}\left(M\left(\begin{smallmatrix}0\,0\\-1\end{smallmatrix}\right)[-1]\right), \end{aligned}$$

and

$$\begin{aligned} B^{\lambda}_{[21;2;11;0]}(M) =&(m_{11}-m_{22})(m_{22}-m_{33}), \\ B^{\lambda}_{[21;2;11;1]}(M) =& -(m_{11}-m_{22}-1)\bar{D}(M)\chi_{-}(M) \\ & +(m_{12}-m_{23}+1)\chi_{-}^{(1)}(M)(m_{22}-m_{33}), \\ B^{\lambda}_{[21;2;11;2]}(M) =& -(m_{12}-m_{23})\chi_{-}^{(1)}(M)\bar{D}(M)\chi_{-}(M), \\ B^{\lambda}_{[21;2;01;0]}(M) =& -(m_{11}-m_{22}+1)(m_{22}-m_{33})+(m_{22}-m_{33}), \\ B^{\lambda}_{[21;2;01;1]}(M) =&(m_{11}-m_{22})\bar{C}_1(M) \end{aligned}$$

$$\begin{aligned}
&\qquad -(m_{12}-m_{23}+1)\chi_-(M)(m_{22}-m_{33})-\bar{D}(M)\chi_-(M),\\
B^{\lambda}_{[21;2;01;2]}(M) =&(m_{12}-m_{23})\chi_-(M)\bar{C}_1(M),\\
B^{\lambda}_{[21;2;00;0]}(M) =&-(m_{11}-m_{22})(m_{23}-m_{22}),\\
B^{\lambda}_{[21;2;00;1]}(M) =&-(m_{11}-m_{22}-1)\bar{C}_1(M)\chi_-(M)\\
&-(m_{12}-m_{23}+1)\chi_-^{(1)}(M)(m_{23}-m_{22}),\\
B^{\lambda}_{[21;2;00;2]}(M) =&-(m_{12}-m_{23})\chi_-^{(1)}(M)\bar{C}_1(M)\chi_-(M).
\end{aligned}$$

We have

$$\begin{aligned}
A^{\lambda}_{[21;2;11;1]}(M) =&-(m_{11}-m_{22})(-m_{22}+m_{33}+\delta(M)+1)\chi_-^{(1)}(M)\\
&+(m_{12}-m_{23})\chi_-(M)(m_{22}-m_{33}+1)\\
=&-(m_{11}-m_{22})(\delta(M)+1)\chi_-^{(1)}(M)+(m_{12}-m_{23})\chi_-(M)\\
&+(m_{22}-m_{33})\{(m_{11}-m_{22})\chi_-^{(1)}(M)+(m_{12}-m_{23})\chi_-(M)\}\\
=&-(m_{11}-m_{22}-1)\delta(M)\chi_-(M)\\
&+(m_{22}-m_{33})\{(m_{11}-m_{22})\chi_-^{(1)}(M)+(m_{12}-m_{23})\chi_-(M)\},
\end{aligned}$$

$$\begin{aligned}
B^{\lambda}_{[21;2;11;1]}(M) =&-(m_{11}-m_{22}-1)(-m_{22}+m_{33}+\delta(M))\chi_-(M)\\
&+(m_{12}-m_{23}+1)\chi_-^{(1)}(M)(m_{22}-m_{33})\\
=&-(m_{11}-m_{22}-1)\delta(M)\chi_-(M)\\
&+(m_{22}-m_{33})\{(m_{11}-m_{22}-1)\chi_-(M)+(m_{12}-m_{23}+1)\chi_-^{(1)}(M)\}.
\end{aligned}$$

Therefore

$$\begin{aligned}
&A^{\lambda}_{[21;2;11;1]}(M)-B^{\lambda}_{[21;2;11;1]}(M)\\
&=(m_{22}-m_{33})\{(\delta(M)+1)\chi_-(M)-(\delta(M)+1)\chi_-^{(1)}(M)\}=0.
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[21;2;11;1]}(M)=B^{\lambda}_{[21;2;11;1]}(M)$.

We have

$$\begin{aligned}
A^{\lambda}_{[21;2;11;2]}(M) =&-(m_{12}-m_{23})\chi_-(M)(-m_{22}+m_{33}+\delta(M))\chi_-^{(1)}(M)\\
=&-(m_{12}-m_{23})\bar{D}(M)\chi_-^{(1)}(M)\chi_-(M),
\end{aligned}$$

$$B^{\lambda}_{[21;2;11;2]}(M) =-(m_{12}-m_{23})\bar{D}(M)\chi_-^{(1)}(M)\chi_-(M).$$

Hence we obtain $A^{\lambda}_{[21;2;11;2]}(M)=B^{\lambda}_{[21;2;11;2]}(M)$.

We have

$$\begin{aligned}
A^{\lambda}_{[21;2;01;1]}(M) =&(m_{11}-m_{22})\{m_{23}-m_{22}+(\delta(M)+1)\chi_-(M)\}\\
&-(m_{12}-m_{23})\chi_-(M)(m_{22}-m_{33}+1),
\end{aligned}$$

$$\begin{aligned}
B^{\lambda}_{[21;2;01;1]}(M) =&(m_{11}-m_{22})(m_{23}-m_{22}+\delta(M)\chi_-(M))\\
&-(m_{12}-m_{23}+1)\chi_-(M)(m_{22}-m_{33})-\bar{D}(M)\chi_-(M).
\end{aligned}$$

Therefore

$$\begin{aligned}
&A^{\lambda}_{[21;2;01;1]}(M)-B^{\lambda}_{[21;2;01;1]}(M)\\
&=(m_{11}-m_{22})\chi_-(M)-(m_{12}-m_{23})\chi_-(M)+(m_{22}-m_{33})\chi_-(M)+\bar{D}(M)\chi_-(M)=0.
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[21;2;01;1]}(M)=B^{\lambda}_{[21;2;01;1]}(M)$.

We have

$$\begin{aligned} A^{\lambda}_{[21;2;01;2]}(M) =&(m_{12}-m_{23})\chi_-(M)\{m_{12}-m_{11}-(\delta(M)+1)\chi_+^{(-1)}(M)\} \\ =& \,(m_{12}-m_{23})(m_{12}-m_{11})\chi_-(M), \end{aligned}$$

$$B^{\lambda}_{[21;2;01;2]}(M) =(m_{12}-m_{23})(m_{12}-m_{11})\chi_-(M).$$

Hence we obtain $A^{\lambda}_{[21;2;01;2]}(M) = B^{\lambda}_{[21;2;01;2]}(M)$.

We have

$$\begin{aligned} &A^{\lambda}_{[21;2;00;1]}(M) \\ &= -(m_{11}-m_{22})\bar{C}_1\left(M\left(\begin{smallmatrix}0\,0\\-1\end{smallmatrix}\right)\right)\chi_-\left(M\left(\begin{smallmatrix}0\,0\\-1\end{smallmatrix}\right)\right) - (m_{12}-m_{23})\chi_-(M)(m_{23}-m_{22}-1) \\ &= \begin{cases} 0 & \text{if } \delta(M) > -1, \\ -(m_{12}-m_{23})(m_{23}-m_{22}-1) & \text{if } \delta(M) = -1, \\ -(m_{11}-m_{22})(m_{12}-m_{11}+1) - (m_{12}-m_{23})(m_{23}-m_{22}-1) & \text{if } \delta(M) < -1, \end{cases} \\ &B^{\lambda}_{[21;2;00;1]}(M) \\ &= -(m_{11}-m_{22}-1)\bar{C}_1(M)\chi_-(M) - (m_{12}-m_{23}+1)\chi_-^{(1)}(M)(m_{23}-m_{22}) \\ &= \begin{cases} 0 & \text{if } \delta(M) > -1, \\ -(m_{11}-m_{22}-1)(m_{12}-m_{11}) & \text{if } \delta(M) = -1, \\ -(m_{11}-m_{22}-1)(m_{12}-m_{11}) - (m_{12}-m_{23}+1)(m_{23}-m_{22}) & \text{if } \delta(M) < -1. \end{cases} \end{aligned}$$

Hence we obtain $A^{\lambda}_{[21;2;00;1]}(M) = B^{\lambda}_{[21;2;00;1]}(M)$.

We have

$$\begin{aligned} A^{\lambda}_{[21;2;00;2]}(M) =& -(m_{12}-m_{23})\chi_-(M)(m_{12}-m_{11})\chi_-\left(M\left(\begin{smallmatrix}0\,0\\-1\end{smallmatrix}\right)[-1]\right) \\ =& -(m_{12}-m_{23})(m_{12}-m_{11})\chi_-^{(1)}(M)\chi_-(M), \\ B^{\lambda}_{[21;2;00;2]}(M) =& -(m_{12}-m_{23})\chi_-^{(1)}(M)(m_{12}-m_{11})\chi_-(M). \end{aligned}$$

Hence we obtain $A^{\lambda}_{[21;2;00;2]}(M) = B^{\lambda}_{[21;2;00;2]}(M)$.

It is trivial that the equations (4.17) hold for $(j,k,l) = (1,1,0),\ (0,1,0)$ and $(0,0,0)$.

- the proof of $E_{23} \circ i^{\lambda}_{\mathbf{e}_2} = i^{\lambda}_{\mathbf{e}_2} \circ E_{23}$.

We have

$$\begin{aligned} A^{\lambda}_{[23;2;11;0]}(M) =&(m_{13}-m_{12})(m_{22}-m_{33}), \\ A^{\lambda}_{[23;2;11;1]}(M) =& -(m_{13}-m_{12})\bar{D}\left(M\left(\begin{smallmatrix}1\,0\\0\end{smallmatrix}\right)\right)\chi_-\left(M\left(\begin{smallmatrix}1\,0\\0\end{smallmatrix}\right)\right) \\ &+\{m_{13}-m_{12}-\delta(M)\}\chi_-(M)(m_{22}-m_{33}+1), \\ A^{\lambda}_{[23;2;11;2]}(M) =& -\{m_{13}-m_{12}-\delta(M)\}\chi_-(M)\bar{D}\left(M\left(\begin{smallmatrix}1\,0\\0\end{smallmatrix}\right)[-1]\right)\chi_-\left(M\left(\begin{smallmatrix}1\,0\\0\end{smallmatrix}\right)[-1]\right), \\ A^{\lambda}_{[23;2;01;0]}(M) =& -(m_{13}-m_{12})(m_{22}-m_{33}), \\ A^{\lambda}_{[23;2;01;1]}(M) =&(m_{13}-m_{12})\bar{C}_1\left(M\left(\begin{smallmatrix}1\,0\\0\end{smallmatrix}\right)\right) \\ &-\{m_{13}-m_{12}-\delta(M)\}\chi_-(M)(m_{22}-m_{33}+1), \\ A^{\lambda}_{[23;2;01;2]}(M) =&\{m_{13}-m_{12}-\delta(M)\}\chi_-(M)\bar{C}_1\left(M\left(\begin{smallmatrix}1\,0\\0\end{smallmatrix}\right)[-1]\right), \\ A^{\lambda}_{[23;2;00;0]}(M) =& -(m_{13}-m_{12})(m_{23}-m_{22}), \\ A^{\lambda}_{[23;2;00;1]}(M) =& -(m_{13}-m_{12})\bar{C}_1\left(M\left(\begin{smallmatrix}1\,0\\0\end{smallmatrix}\right)\right)\chi_-\left(M\left(\begin{smallmatrix}1\,0\\0\end{smallmatrix}\right)\right) \\ &-\{m_{13}-m_{12}-\delta(M)\}\chi_-(M)(m_{23}-m_{22}-1), \\ A^{\lambda}_{[23;2;00;2]}(M) =& -\{m_{13}-m_{12}-\delta(M)\}\chi_-(M)\bar{C}_1\left(M\left(\begin{smallmatrix}1\,0\\0\end{smallmatrix}\right)[-1]\right)\chi_-\left(M\left(\begin{smallmatrix}1\,0\\0\end{smallmatrix}\right)[-1]\right), \end{aligned}$$

and

$$\begin{aligned}
B^{\lambda}_{[23;2;11;0]}(M) =&(m_{13}-m_{12})(m_{22}-m_{33}),\\
B^{\lambda}_{[23;2;11;1]}(M) =&-(m_{13}-m_{12}+1)\bar{D}(M)\chi_-(M)\\
&+\{m_{13}-m_{12}-\delta(M)-1\}\chi_-^{(1)}(M)(m_{22}-m_{33}),\\
B^{\lambda}_{[23;2;11;2]}(M) =&-\{m_{13}-m_{12}-\delta(M)\}\chi_-^{(1)}(M)\bar{D}(M)\chi_-(M),\\
B^{\lambda}_{[23;2;01;0]}(M) =&-(m_{13}-m_{12})(m_{22}-m_{33}),\\
B^{\lambda}_{[23;2;01;1]}(M) =&(m_{13}-m_{12}+1)\bar{C}_1(M)\\
&-\{m_{13}-m_{12}-\delta(M)\}\chi_-(M)(m_{22}-m_{33})-(m_{23}-m_{22}),\\
B^{\lambda}_{[23;2;01;2]}(M) =&\{m_{13}-m_{12}-\delta(M)+1\}\chi_-(M)\bar{C}_1(M)-\bar{C}_1(M)\chi_-(M),\\
B^{\lambda}_{[23;2;00;0]}(M) =&-(m_{13}-m_{12})(m_{23}-m_{22}),\\
B^{\lambda}_{[23;2;00;1]}(M) =&-(m_{13}-m_{12}+1)\bar{C}_1(M)\chi_-(M)\\
&-\{m_{13}-m_{12}-\delta(M)-1\}\chi_-^{(1)}(M)(m_{23}-m_{22}),\\
B^{\lambda}_{[23;2;00;2]}(M) =&-\{m_{13}-m_{12}-\delta(M)\}\chi_-^{(1)}(M)\bar{C}_1(M)\chi_-(M).
\end{aligned}$$

We have

$$\begin{aligned}
A^{\lambda}_{[23;2;11;1]}(M) =&-(m_{13}-m_{12})(-m_{22}+m_{33}+\delta(M)+1)\chi_-^{(1)}(M)\\
&+\{m_{13}-m_{12}-\delta(M)\}\chi_-(M)(m_{22}-m_{33}+1),\\
B^{\lambda}_{[23;2;11;1]}(M) =&-(m_{13}-m_{12}+1)(-m_{22}+m_{33}+\delta(M))\chi_-(M)\\
&+\{m_{13}-m_{12}-\delta(M)-1\}\chi_-^{(1)}(M)(m_{22}-m_{33}).
\end{aligned}$$

Therefore

$$\begin{aligned}
&A^{\lambda}_{[23;2;11;1]}(M)-B^{\lambda}_{[23;2;11;1]}(M)\\
&=(m_{13}-m_{12}-m_{22}+m_{33})(\delta(M)+1)(\chi_-(M)-\chi_-^{(1)}(M))=0.
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[23;2;11;1]}(M)=B^{\lambda}_{[23;2;11;1]}(M)$.

We have

$$\begin{aligned}
A^{\lambda}_{[23;2;11;2]}(M) =&-\{m_{13}-m_{12}-\delta(M)\}\chi_-(M)(-m_{22}+m_{33}+\delta(M))\chi_-^{(1)}(M)\\
=&-\{m_{13}-m_{12}-\delta(M)\}\bar{D}(M)\chi_-^{(1)}(M)\chi_-(M),
\end{aligned}$$

$$B^{\lambda}_{[23;2;11;2]}(M) =-\{m_{13}-m_{12}-\delta(M)\}\bar{D}(M)\chi_-^{(1)}(M)\chi_-(M).$$

Hence we obtain $A^{\lambda}_{[23;2;11;2]}(M)=B^{\lambda}_{[23;2;11;2]}(M)$.

We have

$$\begin{aligned}
A^{\lambda}_{[23;2;01;1]}(M) =&(m_{13}-m_{12})\{m_{23}-m_{22}+(\delta(M)+1)\chi_-(M)\}\\
&-\{m_{13}-m_{12}-\delta(M)\}\chi_-(M)(m_{22}-m_{33}+1),
\end{aligned}$$

$$\begin{aligned}
B^{\lambda}_{[23;2;01;1]}(M) =&(m_{13}-m_{12}+1)(m_{23}-m_{22}+\delta(M)\chi_-(M))\\
&-\{m_{13}-m_{12}-\delta(M)\}\chi_-(M)(m_{22}-m_{33})-(m_{23}-m_{22})\\
=&(m_{13}-m_{12})\{m_{23}-m_{22}+(\delta(M)+1)\chi_-(M)\}\\
&-\{m_{13}-m_{12}-\delta(M)\}\chi_-(M)(m_{22}-m_{33}+1).
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[23;2;01;1]}(M)=B^{\lambda}_{[23;2;01;1]}(M)$.

We have

$$\begin{aligned} A^{\lambda}_{[23;2;01;2]}(M) =& \{m_{13}-m_{12}-\delta(M)\}\chi_{-}(M)\{m_{12}-m_{11}-(\delta(M)+1)\chi_{+}^{(-1)}(M)\} \\ =& \{m_{13}-m_{12}-\delta(M)\}(m_{12}-m_{11})\chi_{-}(M), \end{aligned}$$

$$\begin{aligned} B^{\lambda}_{[23;2;01;2]}(M) =& \{m_{13}-m_{12}-\delta(M)\}\bar{C}_1(M)\chi_{-}(M) \\ =& \{m_{13}-m_{12}-\delta(M)\}(m_{12}-m_{11})\chi_{-}(M). \end{aligned}$$

Hence we obtain $A^{\lambda}_{[23;2;01;2]}(M) = B^{\lambda}_{[23;2;01;2]}(M)$.

We have

$$\begin{aligned} A^{\lambda}_{[23;2;00;1]}(M) =& -(m_{13}-m_{12})(m_{12}-m_{11}+1)\chi_{-}^{(1)}(M) \\ & -\{m_{13}-m_{12}-\delta(M)\}\chi_{-}(M)(m_{23}-m_{22}-1), \end{aligned}$$

$$\begin{aligned} B^{\lambda}_{[23;2;00;1]}(M) =& -(m_{13}-m_{12}+1)(m_{12}-m_{11})\chi_{-}(M) \\ & -\{m_{13}-m_{12}-\delta(M)-1\}\chi_{-}^{(1)}(M)(m_{23}-m_{22}). \end{aligned}$$

Therefore

$$\begin{aligned} & A^{\lambda}_{[23;2;00;1]}(M) - B^{\lambda}_{[23;2;00;1]}(M) \\ & = (m_{13}-m_{12}+m_{23}-m_{22})(\delta(M)+1)(\chi_{-}(M)-\chi_{-}^{(1)}(M)) = 0. \end{aligned}$$

Hence we obtain $A^{\lambda}_{[23;2;00;1]}(M) = B^{\lambda}_{[23;2;00;1]}(M)$.

We have

$$\begin{aligned} A^{\lambda}_{[23;2;00;2]}(M) =& -\{m_{13}-m_{12}-\delta(M)\}\chi_{-}(M)(m_{12}-m_{11})\chi_{-}\left(M\left(\begin{smallmatrix}1 & & 0\\ & 0 & \end{smallmatrix}\right)[-1]\right) \\ =& -\{m_{13}-m_{12}-\delta(M)\}\chi_{-}(M)(m_{12}-m_{11})\chi_{-}^{(1)}(M), \end{aligned}$$

$$B^{\lambda}_{[23;2;00;2]}(M) = -\{m_{13}-m_{12}-\delta(M)\}\chi_{-}^{(1)}(M)(m_{12}-m_{11})\chi_{-}(M).$$

Hence $A^{\lambda}_{[23;2;00;2]}(M) = B^{\lambda}_{[23;2;00;2]}(M)$.

It is trivial that the equations (4.17) hold for $(j,k,l) = (1,1,0),\ (0,1,0)$ and $(0,0,0)$.

- the proof of $E_{32}\circ i^{\lambda}_{\mathbf{e}_2} = i^{\lambda}_{\mathbf{e}_2}\circ E_{32}$.

We have

$$\begin{aligned} A^{\lambda}_{[32;2;11;0]}(M) =& (m_{22}-m_{33})(m_{22}-m_{33}-1), \\ A^{\lambda}_{[32;2;11;1]}(M) =& -(m_{22}-m_{33})\bar{D}\left(M\left(\begin{smallmatrix}0 & & -1\\ & 0 & \end{smallmatrix}\right)\right)\chi_{-}\left(M\left(\begin{smallmatrix}0 & & -1\\ & 0 & \end{smallmatrix}\right)\right) \\ & +\{m_{22}-m_{33}+\delta(M)\}\chi_{+}(M)(m_{22}-m_{33}), \\ A^{\lambda}_{[32;2;11;2]}(M) =& -\{m_{22}-m_{33}+\delta(M)\}\chi_{+}(M)\bar{D}\left(M\left(\begin{smallmatrix}0 & & -1\\ & 0 & \end{smallmatrix}\right)[-1]\right)\chi_{-}\left(M\left(\begin{smallmatrix}0 & & -1\\ & 0 & \end{smallmatrix}\right)[-1]\right), \\ A^{\lambda}_{[32;2;01;0]}(M) =& -(m_{22}-m_{33})(m_{22}-m_{33}-1), \\ A^{\lambda}_{[32;2;01;1]}(M) =& (m_{22}-m_{33})\bar{C}_1\left(M\left(\begin{smallmatrix}0 & & -1\\ & 0 & \end{smallmatrix}\right)\right) \\ & -\{m_{22}-m_{33}+\delta(M)\}\chi_{+}(M)(m_{22}-m_{33}), \\ A^{\lambda}_{[32;2;01;2]}(M) =& \{m_{22}-m_{33}+\delta(M)\}\chi_{+}(M)\bar{C}_1\left(M\left(\begin{smallmatrix}0 & & -1\\ & 0 & \end{smallmatrix}\right)[-1]\right), \\ A^{\lambda}_{[32;2;00;0]}(M) =& -(m_{22}-m_{33})(m_{23}-m_{22}+1), \\ A^{\lambda}_{[32;2;00;1]}(M) =& -(m_{22}-m_{33})\bar{C}_1\left(M\left(\begin{smallmatrix}0 & & -1\\ & 0 & \end{smallmatrix}\right)\right)\chi_{-}\left(M\left(\begin{smallmatrix}0 & & -1\\ & 0 & \end{smallmatrix}\right)\right) \\ & -\{m_{22}-m_{33}+\delta(M)\}\chi_{+}(M)(m_{23}-m_{22}), \\ A^{\lambda}_{[32;2;00;2]}(M) =& -\{m_{22}-m_{33}+\delta(M)\}\chi_{+}(M)\bar{C}_1\left(M\left(\begin{smallmatrix}0 & & -1\\ & 0 & \end{smallmatrix}\right)[-1]\right)\chi_{-}\left(M\left(\begin{smallmatrix}0 & & -1\\ & 0 & \end{smallmatrix}\right)[-1]\right), \end{aligned}$$

and

$$\begin{aligned}
B^{\lambda}_{[32;2;11;0]}(M) =&(m_{22}-m_{33}-1)(m_{22}-m_{33}),\\
B^{\lambda}_{[32;2;11;1]}(M) =&-(m_{22}-m_{33})\bar{D}(M)\chi_{-}(M)\\
&+\{m_{22}-m_{33}+\delta(M)\}\chi_{+}^{(-1)}(M)(m_{22}-m_{33}),\\
B^{\lambda}_{[32;2;11;2]}(M) =&-\{m_{22}-m_{33}+\delta(M)+1\}\chi_{+}^{(-1)}(M)\bar{D}(M)\chi_{-}(M),\\
B^{\lambda}_{[32;2;01;0]}(M) =&-(m_{22}-m_{33}-1)(m_{22}-m_{33}),\\
B^{\lambda}_{[32;2;01;1]}(M) =&(m_{22}-m_{33})\bar{C}_1(M)\\
&-\{m_{22}-m_{33}+\delta(M)-1\}\chi_{+}(M)(m_{22}-m_{33}),\\
B^{\lambda}_{[32;2;01;2]}(M) =&\{m_{22}-m_{33}+\delta(M)\}\chi_{+}(M)\bar{C}_1(M),\\
B^{\lambda}_{[32;2;00;0]}(M) =&-(m_{22}-m_{33})(m_{23}-m_{22})-(m_{22}-m_{33}),\\
B^{\lambda}_{[32;2;00;1]}(M) =&-(m_{22}-m_{33}+1)\bar{C}_1(M)\chi_{-}(M)\\
&-\{m_{22}-m_{33}+\delta(M)+1\}\chi_{+}^{(-1)}(M)(m_{23}-m_{22})+\bar{C}_1(M),\\
B^{\lambda}_{[32;2;00;2]}(M) =&-\{m_{22}-m_{33}+\delta(M)+2\}\chi_{+}^{(-1)}(M)\bar{C}_1(M)\chi_{-}(M).
\end{aligned}$$

We have

$$\begin{aligned}
A^{\lambda}_{[32;2;11;1]}(M) =&-(m_{22}-m_{33})(-m_{22}+m_{33}+\delta(M))(1-\chi_{+}(M))\\
&+\{m_{22}-m_{33}+\delta(M)\}\chi_{+}(M)(m_{22}-m_{33})\\
=&(m_{22}-m_{33})\{-\bar{D}(M)+2\delta(M)\chi_{+}(M)\},
\end{aligned}$$

$$\begin{aligned}
B^{\lambda}_{[32;2;11;1]}(M) =&-(m_{22}-m_{33})(-m_{22}+m_{33}+\delta(M))(1-\chi_{+}^{(-1)}(M))\\
&+\{m_{22}-m_{33}+\delta(M)\}\chi_{+}^{(-1)}(M)(m_{22}-m_{33})\\
=&(m_{22}-m_{33})\{-\bar{D}(M)+2\delta(M)\chi_{+}^{(-1)}(M)\}\\
=&(m_{22}-m_{33})\{-\bar{D}(M)+2\delta(M)\chi_{+}(M)\}.
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[32;2;11;1]}(M)=B^{\lambda}_{[32;2;11;1]}(M)$.

We have

$$A^{\lambda}_{[32;2;11;2]}(M)=-\{m_{22}-m_{33}+\delta(M)\}\bar{D}\left(M\left(\begin{smallmatrix}0&-1\\&0\end{smallmatrix}\right)[-1]\right)\chi_{+}(M)\chi_{-}^{(-1)}(M)=0,$$

$$B^{\lambda}_{[32;2;11;2]}(M)=-\{m_{22}-m_{33}+\delta(M)+1\}\bar{D}(M)\chi_{+}^{(-1)}(M)\chi_{-}(M)=0.$$

Hence we obtain $A^{\lambda}_{[32;2;11;2]}(M)=B^{\lambda}_{[32;2;11;2]}(M)$.

We have

$$\begin{aligned}
A^{\lambda}_{[32;2;01;1]}(M) =&(m_{22}-m_{33})\{m_{12}-m_{11}-(\delta(M)-1)\chi_{+}(M)\}\\
&-\{m_{22}-m_{33}+\delta(M)\}\chi_{+}(M)(m_{22}-m_{33})\\
=&(m_{22}-m_{33})\{(m_{12}-m_{11})-\chi_{+}(M)(m_{22}-m_{33}+2\delta(M)-1)\},
\end{aligned}$$

$$\begin{aligned}
B^{\lambda}_{[32;2;01;1]}(M) =&(m_{22}-m_{33})\{m_{12}-m_{11}-\delta(M)\chi_{+}(M)\}\\
&-\{m_{22}-m_{33}+\delta(M)-1\}\chi_{+}(M)(m_{22}-m_{33})\\
=&(m_{22}-m_{33})\{(m_{12}-m_{11})-\chi_{+}(M)(m_{22}-m_{33}+2\delta(M)-1)\}.
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[32;2;01;1]}(M)=B^{\lambda}_{[32;2;01;1]}(M)$.

We have

$$\begin{aligned}
A^{\lambda}_{[32;2;01;2]}(M) =& \{m_{22} - m_{33} + \delta(M)\}\chi_{+}(M)(m_{23} - m_{22} + (\delta(M) - 1)\chi_{-}(M)) \\
=& \{m_{22} - m_{33} + \delta(M)\}(m_{23} - m_{22})\chi_{+}(M), \\
B^{\lambda}_{[32;2;01;2]}(M) =& \{m_{22} - m_{33} + \delta(M)\}(m_{23} - m_{22})\chi_{+}(M).
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[32;2;01;2]}(M) = B^{\lambda}_{[32;2;01;2]}(M)$.

We have

$$\begin{aligned}
A^{\lambda}_{[32;2;00;1]}(M) =& -(m_{22} - m_{33})(m_{12} - m_{11})\chi_{-}^{(-1)}(M) \\
& - \{m_{22} - m_{33} + \delta(M)\}(m_{23} - m_{22})\chi_{+}(M) \\
=& -(m_{22} - m_{33})\bar{C}_1(M)\chi_{-}^{(-1)}(M) - \{m_{22} - m_{33} + \delta(M)\}\bar{C}_1(M)\chi_{+}(M) \\
=& -(m_{22} - m_{33})\bar{C}_1(M)(\chi_{-}^{(-1)}(M) + \chi_{+}(M)) - \bar{C}_1(M)\delta(M)\chi_{+}(M) \\
=& -(m_{22} - m_{33})\bar{C}_1(M) - \bar{C}_1(M)\delta(M)\chi_{+}(M),
\end{aligned}$$

$$\begin{aligned}
B^{\lambda}_{[32;2;00;1]}(M) =& -(m_{22} - m_{33} + 1)\bar{C}_1(M)\chi_{-}(M) \\
& - \{m_{22} - m_{33} + \delta(M) + 1\}\bar{C}_1(M)\chi_{+}^{(-1)}(M) + \bar{C}_1(M) \\
=& -(m_{22} - m_{33} + 1)\bar{C}_1(M)(\chi_{-}(M) + \chi_{+}^{(-1)}(M)) + \bar{C}_1(M) \\
& - \bar{C}_1(M)\delta(M)\chi_{+}^{(-1)}(M) \\
=& -(m_{22} - m_{33})\bar{C}_1(M) - \bar{C}_1(M)\delta(M)\chi_{+}(M).
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[32;2;00;1]}(M) = B^{\lambda}_{[32;2;00;1]}(M)$.

We have

$$A^{\lambda}_{[32;2;00;2]}(M) = -\{m_{22} - m_{33} + \delta(M)\}\bar{C}_1\left(M\left(\begin{smallmatrix}0&&-1\\&0&\end{smallmatrix}\right)[-1]\right)\chi_{+}(M)\chi_{-}^{(-1)}(M) = 0,$$

$$B^{\lambda}_{[32;2;00;2]}(M) = -\{m_{22} - m_{33} + \delta(M) + 2\}\bar{C}_1(M)\chi_{+}^{(-1)}(M)\chi_{-}(M) = 0.$$

Hence we obtain $A^{\lambda}_{[32;2;00;2]}(M) = B^{\lambda}_{[32;2;00;2]}(M)$.

It is trivial that the equations (4.17) hold for $(j, k, l) = (1, 1, 0),\ (0, 1, 0)$ and $(0, 0, 0)$.

At last, we check the equations 4.17 for $i = 1$, that is, the case of formula 1 by direct computation.

- the proof of $E_{12} \circ i^{\lambda}_{\mathbf{e}_1} = i^{\lambda}_{\mathbf{e}_1} \circ E_{12}$.

We have

$$\begin{aligned}
A^{\lambda}_{[12;1;11;0]}(M) =& (m_{12} - m_{11})(m_{13} - m_{12})(m_{22} - m_{33}), \\
A^{\lambda}_{[12;1;11;1]}(M) =& -(m_{12} - m_{11})\bar{E}\left(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)\right) \\
& + (m_{23} - m_{22})\chi_{+}(M)(m_{13} - m_{12} + 1)(m_{22} - m_{33} + 1), \\
A^{\lambda}_{[12;1;11;2]}(M) =& -(m_{23} - m_{22})\chi_{+}(M)\bar{E}\left(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)[-1]\right), \\
A^{\lambda}_{[12;1;01;0]}(M) =& -(m_{12} - m_{11})(m_{13} - m_{12})(m_{22} - m_{33}), \\
A^{\lambda}_{[12;1;01;1]}(M) =& (m_{12} - m_{11})\bar{F}\left(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)\right) \\
& - (m_{23} - m_{22})\chi_{+}(M)(m_{13} - m_{12} + 1)(m_{22} - m_{33} + 1), \\
A^{\lambda}_{[12;1;01;2]}(M) =& -(m_{12} - m_{11})C_2\left(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)\right)\chi_{+}\left(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)\right) \\
& + (m_{23} - m_{22})\chi_{+}(M)\bar{F}\left(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)[-1]\right), \\
A^{\lambda}_{[12;1;01;3]}(M) =& -(m_{23} - m_{22})\chi_{+}(M)C_2\left(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)[-1]\right)\chi_{+}\left(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)[-1]\right),
\end{aligned}$$

$$\begin{aligned}
A^{\lambda}_{[12;1;00;0]}(M) =& -(m_{12}-m_{11})(m_{13}-m_{12})(m_{13}-m_{22}+1),\\
A^{\lambda}_{[12;1;00;1]}(M) =& (m_{12}-m_{11})C_2\left(M\left(\begin{smallmatrix}0 & & 0\\ & 1 & \end{smallmatrix}\right)\right)\\
&-(m_{23}-m_{22})\chi_+(M)(m_{13}-m_{12}+1)(m_{13}-m_{22}),\\
A^{\lambda}_{[12;1;00;2]}(M) =& (m_{23}-m_{22})\chi_+(M)C_2\left(M\left(\begin{smallmatrix}0 & & 0\\ & 1 & \end{smallmatrix}\right)[-1]\right),
\end{aligned}$$

and

$$\begin{aligned}
B^{\lambda}_{[12;1;11;0]}(M) =& (m_{12}-m_{11}+1)(m_{13}-m_{12})(m_{22}-m_{33})-(m_{13}-m_{12})(m_{22}-m_{33}),\\
B^{\lambda}_{[12;1;11;1]}(M) =& -(m_{12}-m_{11})\bar{E}(M)\\
&+(m_{23}-m_{22}+1)\chi_+(M)(m_{13}-m_{12})(m_{22}-m_{33})+\bar{F}(M),\\
B^{\lambda}_{[12;1;11;2]}(M) =& -(m_{23}-m_{22})\chi_+(M)\bar{E}(M)-C_2(M)\chi_+(M),\\
B^{\lambda}_{[12;1;01;0]}(M) =& -(m_{12}-m_{11})(m_{13}-m_{12})(m_{22}-m_{33}),\\
B^{\lambda}_{[12;1;01;1]}(M) =& (m_{12}-m_{11}-1)\bar{F}(M)\\
&-(m_{23}-m_{22}+1)\chi_+^{(1)}(M)(m_{13}-m_{12})(m_{22}-m_{33}),\\
B^{\lambda}_{[12;1;01;2]}(M) =& -(m_{12}-m_{11}-2)C_2(M)\chi_+(M)+(m_{23}-m_{22})\chi_+^{(1)}(M)\bar{F}(M),\\
B^{\lambda}_{[12;1;01;3]}(M) =& -(m_{23}-m_{22}-1)\chi_+^{(1)}(M)C_2(M)\chi_+(M),\\
B^{\lambda}_{[12;1;00;0]}(M) =& -(m_{12}-m_{11})(m_{13}-m_{12})(m_{13}-m_{22}+1),\\
B^{\lambda}_{[12;1;00;1]}(M) =& (m_{12}-m_{11}-1)C_2(M)\\
&-(m_{23}-m_{22})\chi_+(M)(m_{13}-m_{12})(m_{13}-m_{22}+1),\\
B^{\lambda}_{[12;1;00;2]}(M) =& (m_{23}-m_{22}-1)\chi_+(M)C_2(M).
\end{aligned}$$

We have

$$\begin{aligned}
&A^{\lambda}_{[12;1;11;1]}(M)-B^{\lambda}_{[12;1;11;1]}(M)\\
&=-(m_{12}-m_{11})\{(m_{12}-m_{23})+(m_{13}-m_{33}-m_{12}+m_{22}+1)\chi_+(M)\}\\
&\quad+(m_{13}-m_{33}-m_{12}+m_{22}+1)(m_{23}-m_{22})\chi_+(M)\\
&\quad-(m_{13}-m_{12})(m_{22}-m_{33})\chi_+(M)-\bar{F}(M)\\
&=-(m_{13}-m_{33}-m_{12}+m_{22}+1)\delta(M)\chi_+(M)\\
&\quad-(m_{13}-m_{12})(m_{22}-m_{33})\chi_+(M)-(m_{12}-m_{11})(m_{12}-m_{23})-\bar{F}(M)\\
&=-\{(m_{12}-m_{11})(m_{12}-m_{23})-(m_{12}-m_{22})\delta(M)\chi_+(M)\}\\
&\quad-\chi_+(M)\{(m_{13}-m_{12})(m_{22}-m_{33})+(m_{13}-m_{33}+1)\delta(M)\}-\bar{F}(M)\\
&=0.
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[12;1;11;1]}(M)=B^{\lambda}_{[12;1;11;1]}(M)$. Here we use the relations

$$\bar{E}\left(M\left(\begin{smallmatrix}0 & & 0\\ & 1 & \end{smallmatrix}\right)\right)=\bar{E}(M)+(m_{12}-m_{23})+(m_{13}-m_{33}-m_{12}+m_{22}+1)\chi_+(M). \tag{4.18}$$

We have

$$\begin{aligned}
A^{\lambda}_{[12;1;11;2]}(M) =& -(m_{23}-m_{22})(\bar{E}(M)+C_1(M))\chi_+(M)\\
=& -(m_{23}-m_{22})\chi_+(M)\bar{E}(M)-C_1(M)(m_{23}-m_{22})\chi_+(M)\\
=& B^{\lambda}_{[12;1;11;2]}(M).
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[12;1;11;2]}(M)=B^{\lambda}_{[12;1;11;2]}(M)$. Here we use the relation

$$\bar{E}\left(M\left(\begin{smallmatrix}0 & & 0\\ & 1 & \end{smallmatrix}\right)[-1]\right)\chi_+(M)=(\bar{E}(M)+C_1(M))\chi_+(M). \tag{4.19}$$

We have

$$\begin{aligned}
&A^{\lambda}_{[12;1;01;1]}(M) - B^{\lambda}_{[12;1;01;1]}(M) \\
&= (m_{12} - m_{11})\{(m_{12} - m_{23}) - (m_{13} - m_{12})(m_{22} - m_{33})\chi^{(1)}_{+}(M) \\
&\quad + (m_{13} - m_{12} + 1)(m_{22} - m_{33} + 1)\chi_{+}(M)\} + \bar{F}(M) \\
&\quad - (m_{23} - m_{22})\chi_{+}(M)(m_{13} - m_{12} + 1)(m_{22} - m_{33} + 1) \\
&\quad + (m_{23} - m_{22} + 1)\chi^{(1)}_{+}(M)(m_{13} - m_{12})(m_{22} - m_{33}) \\
&= (m_{12} - m_{11})(m_{12} - m_{23}) - (m_{13} - m_{12})(m_{22} - m_{33})(\delta(M) - 1)\chi^{(1)}_{+}(M) \\
&\quad + (m_{13} - m_{12} + 1)(m_{22} - m_{33} + 1)\delta(M)\chi_{+}(M) + \bar{F}(M) \\
&= \{(m_{12} - m_{11})(m_{12} - m_{23}) - (m_{12} - m_{22})\delta(M)\chi_{+}(M)\} \\
&\quad + \chi_{+}(M)\{(m_{13} - m_{12})(m_{22} - m_{33}) + (m_{13} - m_{33} + 1)\delta(M)\} + \bar{F}(M) \\
&= 0.
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[12;1;01;1]}(M) = B^{\lambda}_{[12;1;01;1]}(M)$. Here we use the relations

$$\begin{aligned}
\bar{F}\left(M\left(\begin{smallmatrix} 0 & & 0 \\ & 1 & \end{smallmatrix}\right)\right) =& \bar{F}(M) + (m_{12} - m_{23}) - (m_{13} - m_{12})(m_{22} - m_{33})\chi^{(1)}_{+}(M) \\
&+ (m_{13} - m_{12} + 1)(m_{22} - m_{33} + 1)\chi_{+}(M).
\end{aligned} \tag{4.20}$$

We have

$$\begin{aligned}
A^{\lambda}_{[12;1;01;2]}(M) =& (m_{23} - m_{22})\{-(m_{12} - m_{11})(m_{11} - m_{22} + 1)\chi^{(1)}_{+}(M) \\
&+ \bar{F}\left(M\left(\begin{smallmatrix} 0 & & 0 \\ & 1 & \end{smallmatrix}\right)[-1]\right)\chi_{+}(M)\}, \\
B^{\lambda}_{[12;1;01;2]}(M) =& (m_{23} - m_{22})\{-(m_{12} - m_{11} - 2)(m_{11} - m_{22})\chi_{+}(M) \\
&+ \bar{F}(M)\chi^{(1)}_{+}(M)\}.
\end{aligned}$$

By the relation

$$\begin{aligned}
\bar{F}\left(M\left(\begin{smallmatrix} 0 & & 0 \\ & 1 & \end{smallmatrix}\right)[-1]\right)\chi_{+}(M) =& \bar{F}(M)\chi^{(1)}_{+}(M) + (m_{12} - m_{11})(m_{11} - m_{22} + 1)\chi^{(1)}_{+}(M) \\
&- (m_{12} - m_{11} - 2)(m_{11} - m_{22})\chi_{+}(M),
\end{aligned} \tag{4.21}$$

we have $A^{\lambda}_{[12;1;01;2]}(M) - B^{\lambda}_{[12;1;01;2]}(M) = 0$. Hence we obtain $A^{\lambda}_{[12;1;01;2]}(M) = B^{\lambda}_{[12;1;01;2]}(M)$.

We have

$$\begin{aligned}
A^{\lambda}_{[12;1;01;3]}(M) =& - (m_{23} - m_{22})\chi_{+}(M)C_1(M)(m_{23} - m_{22} - 1)\chi_{+}\left(M\left(\begin{smallmatrix} 0 & & 0 \\ & 1 & \end{smallmatrix}\right)[-1]\right) \\
=& - \bar{C}_1(M)\chi_{+}(M)C_1(M)(m_{23} - m_{22} - 1)\chi^{(1)}_{+}(M) \\
=& B^{\lambda}_{[12;1;01;3]}(M).
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[12;1;01;3]}(M) = B^{\lambda}_{[12;1;01;3]}(M)$.

We have

$$\begin{aligned}
&A^{\lambda}_{[12;1;00;1]}(M) - B^{\lambda}_{[12;1;00;1]}(M) \\
&= (m_{12} - m_{11})\{(m_{12} - m_{22})\chi_{+}(M) - (m_{12} - m_{23})\} + C_2(M) \\
&\quad - (m_{23} - m_{22})\chi_{+}(M)(m_{12} - m_{22}) \\
&= C_2(M) - (m_{12} - m_{23})(m_{12} - m_{11}) + (m_{12} - m_{22})\delta(M)\chi_{+}(M) \\
&= 0.
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[12;1;00;1]}(M) = B^{\lambda}_{[12;1;00;1]}(M)$. Here we use the relations

$$C_2\left(M\left(\begin{smallmatrix} 0 & & 0 \\ & 1 & \end{smallmatrix}\right)\right) = C_2(M) + (m_{12} - m_{22})\chi_{+}(M) - (m_{12} - m_{23}). \tag{4.22}$$

We have

$$\begin{aligned}&A^{\lambda}_{[12;1;00;2]}(M)\\&=(m_{11}-m_{22})(m_{23}-m_{22})(m_{23}-m_{22}-1)\chi_{+}(M)C_2(M)\\&=B^{\lambda}_{[12;1;00;2]}(M)\,.\end{aligned}$$

Hence we obtain $A^{\lambda}_{[12;1;00;2]}(M)=B^{\lambda}_{[12;1;00;2]}(M)$.

It is trivial that the equations (4.17) hold for $(j,k,l)=(1,1,0),\ (0,1,0)$ and $(0,0,0)$.

- the proof of $E_{21}\circ i^{\lambda}_{\mathbf{e}_1}=i^{\lambda}_{\mathbf{e}_1}\circ E_{21}$

We have

$$\begin{aligned}A^{\lambda}_{[21;1;11;0]}(M)=&(m_{11}-m_{22})(m_{13}-m_{12})(m_{22}-m_{33}),\\A^{\lambda}_{[21;1;11;1]}(M)=&-(m_{11}-m_{22})\bar{E}\left(M\left(\begin{smallmatrix}0\,0\\-1\end{smallmatrix}\right)\right)\\&+(m_{12}-m_{23})\chi_{-}(M)(m_{13}-m_{12}+1)(m_{22}-m_{33}+1),\\A^{\lambda}_{[21;1;11;2]}(M)=&-(m_{12}-m_{23})\chi_{-}(M)\bar{E}\left(M\left(\begin{smallmatrix}0\,0\\-1\end{smallmatrix}\right)[-1]\right),\\A^{\lambda}_{[21;1;01;0]}(M)=&-(m_{11}-m_{22})(m_{13}-m_{12})(m_{22}-m_{33}),\\A^{\lambda}_{[21;1;01;1]}(M)=&(m_{11}-m_{22})\bar{F}\left(M\left(\begin{smallmatrix}0\,0\\-1\end{smallmatrix}\right)\right)\\&-(m_{12}-m_{23})\chi_{-}(M)(m_{13}-m_{12}+1)(m_{22}-m_{33}+1),\\A^{\lambda}_{[21;1;01;2]}(M)=&-(m_{11}-m_{22})C_2\left(M\left(\begin{smallmatrix}0\,0\\-1\end{smallmatrix}\right)\right)\chi_{+}\left(M\left(\begin{smallmatrix}0\,0\\-1\end{smallmatrix}\right)\right)\\&+(m_{12}-m_{23})\chi_{-}(M)\bar{F}\left(M\left(\begin{smallmatrix}0\,0\\-1\end{smallmatrix}\right)[-1]\right),\\A^{\lambda}_{[21;1;01;3]}(M)=&-(m_{12}-m_{23})\chi_{-}(M)C_2\left(M\left(\begin{smallmatrix}0\,0\\-1\end{smallmatrix}\right)[-1]\right)\chi_{+}\left(M\left(\begin{smallmatrix}0\,0\\-1\end{smallmatrix}\right)[-1]\right),\\A^{\lambda}_{[21;1;00;0]}(M)=&-(m_{11}-m_{22})(m_{13}-m_{12})(m_{13}-m_{22}+1),\\A^{\lambda}_{[21;1;00;1]}(M)=&(m_{11}-m_{22})C_2\left(M\left(\begin{smallmatrix}0\,0\\-1\end{smallmatrix}\right)\right)\\&-(m_{12}-m_{23})\chi_{-}(M)(m_{13}-m_{12}+1)(m_{13}-m_{22}),\\A^{\lambda}_{[21;1;00;2]}(M)=&(m_{12}-m_{23})\chi_{-}(M)C_2\left(M\left(\begin{smallmatrix}0\,0\\-1\end{smallmatrix}\right)[-1]\right),\end{aligned}$$

and

$$\begin{aligned}B^{\lambda}_{[21;1;11;0]}(M)=&(m_{11}-m_{22})(m_{13}-m_{12})(m_{22}-m_{33}),\\B^{\lambda}_{[21;1;11;1]}(M)=&-(m_{11}-m_{22}-1)\bar{E}(M)\\&+(m_{12}-m_{23})\chi_{-}(M)(m_{13}-m_{12})(m_{22}-m_{33}),\\B^{\lambda}_{[21;1;11;2]}(M)=&-(m_{12}-m_{23}-1)\chi_{-}(M)\bar{E}(M),\\B^{\lambda}_{[21;1;01;0]}(M)=&-(m_{11}-m_{22}+1)(m_{13}-m_{12})(m_{22}-m_{33})+(m_{13}-m_{12})(m_{22}-m_{33}),\\B^{\lambda}_{[21;1;01;1]}(M)=&(m_{11}-m_{22})\bar{F}(M)\\&-(m_{12}-m_{23})\chi_{-}^{(-1)}(M)(m_{13}-m_{12})(m_{22}-m_{33})-\bar{E}(M),\\B^{\lambda}_{[21;1;01;2]}(M)=&-(m_{11}-m_{22}-1)C_2(M)\chi_{+}(M)\\&+(m_{12}-m_{23}-1)\chi_{-}^{(-1)}(M)\bar{F}(M),\\B^{\lambda}_{[21;1;01;3]}(M)=&-(m_{12}-m_{23}-2)\chi_{-}^{(-1)}(M)C_2(M)\chi_{+}(M),\\B^{\lambda}_{[21;1;00;0]}(M)=&-(m_{11}-m_{22})(m_{13}-m_{12})(m_{13}-m_{22}+1),\\B^{\lambda}_{[21;1;00;1]}(M)=&(m_{11}-m_{22}-1)C_2(M)\\&-(m_{12}-m_{23})\chi_{-}(M)(m_{13}-m_{12})(m_{13}-m_{22}+1),\end{aligned}$$

$B^{\lambda}_{[21;1;00;2]}(M) = (m_{12} - m_{23} - 1)\chi_-(M)C_2(M).$

We have

$$\begin{aligned}
&A^{\lambda}_{[21;1;11;1]}(M) - B^{\lambda}_{[21;1;11;1]}(M)\\
&= (m_{11} - m_{22})(m_{12} - m_{23})\chi_-(M) + \bar{E}(M)\chi_+^{(-1)}(M)\\
&\quad + (m_{13} - m_{33} - m_{12} + m_{22} + 1)(m_{12} - m_{23})\chi_-(M) - \bar{E}(M)\\
&= (m_{12} - m_{23})(m_{13} - m_{33} + 1 - m_{12} + m_{11})\chi_-(M) + \bar{E}(M)(\chi_+^{(-1)}(M) - 1)\\
&= \bar{E}(M)(\chi_-(M) + \chi_+^{(-1)}(M) - 1) = 0.
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[21;1;11;1]}(M) = B^{\lambda}_{[21;1;11;1]}(M)$. Here we use the relations

$$\begin{aligned}
(m_{11} - m_{22})\bar{E}\left(M\left(\begin{smallmatrix}0\,0\\ -1\end{smallmatrix}\right)\right) =&(m_{11} - m_{22})\{\bar{E}(M) - (m_{12} - m_{23})\chi_-(M)\}\\
&- \bar{E}(M)\chi_+^{(-1)}(M).
\end{aligned}\tag{4.23}$$

We have

$$\begin{aligned}
A^{\lambda}_{[21;1;11;2]}(M) =& - (m_{12} - m_{23})\chi_-(M)(m_{12} - m_{23} - 1)\{m_{13} - m_{33} + 1 + m_{12} - m_{11}\}\\
=&B^{\lambda}_{[21;1;11;2]}(M).
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[21;1;11;2]}(M) = B^{\lambda}_{[21;1;11;2]}(M)$.

We have

$$\begin{aligned}
&A^{\lambda}_{[21;1;01;1]}(M) - B^{\lambda}_{[21;1;01;1]}(M)\\
&= (m_{11} - m_{22})\{-(m_{13} - m_{12})(m_{22} - m_{33})\chi_-^{(-1)}(M)\\
&\quad + (m_{13} - m_{12} + 1)(m_{22} - m_{33} + 1)\chi_-(M) - (m_{13} - m_{23} - m_{33} + m_{22} + 1)\}\\
&\quad - (m_{12} - m_{23})\chi_-(M)(m_{13} - m_{12} + 1)(m_{22} - m_{33} + 1)\\
&\quad + (m_{12} - m_{23})\chi_-^{(-1)}(M)(m_{13} - m_{12})(m_{22} - m_{33}) + \bar{E}(M)\\
&= -\delta(M)\chi_-(M)(m_{13} - m_{12} + 1)(m_{22} - m_{33} + 1)\\
&\quad + \delta(M)\chi_-^{(-1)}(M)(m_{13} - m_{12})(m_{22} - m_{33})\\
&\quad - (m_{11} - m_{22})(m_{13} - m_{23} - m_{33} + m_{22} + 1) + \bar{E}(M)\\
&= -(m_{11} - m_{22})(m_{13} - m_{23} - m_{33} + m_{22} + 1)\\
&\quad - (m_{13} - m_{33} - m_{12} + m_{22} + 1)\delta(M)\chi_-(M) + \bar{E}(M)\\
&= 0.
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[21;1;01;1]}(M) = B^{\lambda}_{[21;1;01;1]}(M)$. Here we use the relations

$$\begin{aligned}
\bar{F}\left(M\left(\begin{smallmatrix}0\,0\\ -1\end{smallmatrix}\right)\right) =&\bar{F}(M) - (m_{13} - m_{12})(m_{22} - m_{33})\chi_-^{(-1)}(M)\\
&+ (m_{13} - m_{12} + 1)(m_{22} - m_{33} + 1)\chi_-(M)\\
&- (m_{13} - m_{23} - m_{33} + m_{22} + 1),
\end{aligned}\tag{4.24}$$

$$\begin{aligned}
\bar{E}(M) =&(m_{11} - m_{22})(m_{13} - m_{23} - m_{33} + m_{22} + 1)\\
&+ (m_{13} - m_{33} - m_{12} + m_{22} + 1)\delta(M)\chi_-(M).
\end{aligned}\tag{4.25}$$

We have

$$\begin{aligned}
A^{\lambda}_{[21;1;01;2]}(M) =& - (m_{11} - m_{22})(m_{11} - m_{22} - 1)(m_{23} - m_{22})\chi_+^{(-1)}(M)\\
&- (m_{12} - m_{23})(m_{12} - m_{23} - 1)(m_{12} - m_{11})\chi_-(M)\\
=& - C_2(M)(C_1(M) - 1)\chi_+^{(-1)}(M) - C_2(M)(C_1(M) - 1)\chi_-(M)\\
=& - C_2(M)(C_1(M) - 1),
\end{aligned}$$

$$\begin{aligned} B^{\lambda}_{[21;1;01;2]}(M) &= -(C_1(M)-1)C_2(M)\chi_+(M)-(C_1(M)-1)C_2(M)\chi_-^{(-1)}(M)\\ &= -C_2(M)(C_1(M)-1), \end{aligned}$$

Hence we obtain $A^{\lambda}_{[21;1;01;2]}(M)=B^{\lambda}_{[21;1;01;2]}(M)$. Here we use the relation

$$\bar{F}(M)\chi_-^{(r)}(M)=-C_2(M)\chi_-^{(r)}(M) \quad (r\geq -1). \tag{4.26}$$

We have

$$\begin{aligned} A^{\lambda}_{[21;1;01;3]}(M) &= -(m_{12}-m_{23})C_2\left(M\left(\begin{smallmatrix}0\;0\\-1\end{smallmatrix}\right)[-1]\right)\chi_-(M)\chi_+^{(-1)}(M)=0,\\ B^{\lambda}_{[21;1;01;3]}(M) &= -(m_{12}-m_{23}-2)C_2(M)\chi_-^{(-1)}(M)\chi_+(M)=0, \end{aligned}$$

Hence we obtain $A^{\lambda}_{[21;1;01;3]}(M)=B^{\lambda}_{[21;1;01;3]}(M)$.

We have

$$\begin{aligned} &A^{\lambda}_{[21;1;00;1]}(M)-B^{\lambda}_{[21;1;00;1]}(M)\\ &=(m_{11}-m_{22})\{(m_{12}-m_{22})\chi_-(M)-(m_{23}-m_{22})\}+C_2(M)\\ &\quad -(m_{12}-m_{23})(m_{12}-m_{22})\chi_-(M)\\ &=-(m_{11}-m_{22})(m_{23}-m_{22})-(m_{12}-m_{22})\delta(M)\chi_-(M)+C_2(M)\\ &=0. \end{aligned}$$

Hence we obtain $A^{\lambda}_{[21;1;00;1]}(M)=B^{\lambda}_{[21;1;00;1]}(M)$. Here we use the relations

$$C_2\left(M\left(\begin{smallmatrix}0\;0\\-1\end{smallmatrix}\right)\right)=C_2(M)+(m_{12}-m_{22})\chi_-(M)-(m_{23}-m_{22}). \tag{4.27}$$

We have

$$\begin{aligned} A^{\lambda}_{[21;1;00;2]}(M) =&(m_{12}-m_{23})(m_{12}-m_{23}-1)(m_{12}-m_{11})\chi_-(M)\\ =&B^{\lambda}_{[21;1;00;2]}(M). \end{aligned}$$

Hence we obtain $A^{\lambda}_{[21;1;00;2]}(M)=B^{\lambda}_{[21;1;00;2]}(M)$.

It is trivial that the equations (4.17) hold for $(j,k,l)=(1,1,0),\ (0,1,0)$ and $(0,0,0)$.

- the proof of $E_{23}\circ i^{\lambda}_{\mathbf{e}_1}=i^{\lambda}_{\mathbf{e}_1}\circ E_{23}$

We have

$$\begin{aligned} A^{\lambda}_{[23;1;11;0]}(M) =&(m_{13}-m_{12})(m_{13}-m_{12}-1)(m_{22}-m_{33}),\\ A^{\lambda}_{[23;1;11;1]}(M) =&-(m_{13}-m_{12})\bar{E}\left(M\left(\begin{smallmatrix}1\;0\\0\end{smallmatrix}\right)\right)\\ &+\{m_{13}-m_{12}-\delta(M)\}\chi_-(M)(m_{13}-m_{12})(m_{22}-m_{33}+1),\\ A^{\lambda}_{[23;1;11;2]}(M) =&-\{m_{13}-m_{12}-\delta(M)\}\chi_-(M)\bar{E}\left(M\left(\begin{smallmatrix}1\;0\\0\end{smallmatrix}\right)[-1]\right),\\ A^{\lambda}_{[23;1;01;0]}(M) =&-(m_{13}-m_{12})(m_{13}-m_{12}-1)(m_{22}-m_{33}),\\ A^{\lambda}_{[23;1;01;1]}(M) =&(m_{13}-m_{12})\bar{F}\left(M\left(\begin{smallmatrix}1\;0\\0\end{smallmatrix}\right)\right)\\ &-\{m_{13}-m_{12}-\delta(M)\}\chi_-(M)(m_{13}-m_{12})(m_{22}-m_{33}+1),\\ A^{\lambda}_{[23;1;01;2]}(M) =&-(m_{13}-m_{12})C_2\left(M\left(\begin{smallmatrix}1\;0\\0\end{smallmatrix}\right)\right)\chi_+\left(M\left(\begin{smallmatrix}1\;0\\0\end{smallmatrix}\right)\right)\\ &+\{m_{13}-m_{12}-\delta(M)\}\chi_-(M)\bar{F}\left(M\left(\begin{smallmatrix}1\;0\\0\end{smallmatrix}\right)[-1]\right),\\ A^{\lambda}_{[23;1;01;3]}(M) =&-\{m_{13}-m_{12}-\delta(M)\}\chi_-(M)C_2\left(M\left(\begin{smallmatrix}1\;0\\0\end{smallmatrix}\right)[-1]\right)\chi_+\left(M\left(\begin{smallmatrix}1\;0\\0\end{smallmatrix}\right)[-1]\right),\\ A^{\lambda}_{[23;1;00;0]}(M) =&-(m_{13}-m_{12})(m_{13}-m_{12}-1)(m_{13}-m_{22}+1),\\ A^{\lambda}_{[23;1;00;1]}(M) =&(m_{13}-m_{12})C_2\left(M\left(\begin{smallmatrix}1\;0\\0\end{smallmatrix}\right)\right)\\ &-\{m_{13}-m_{12}-\delta(M)\}\chi_-(M)(m_{13}-m_{12})(m_{13}-m_{22}), \end{aligned}$$

$$A^{\lambda}_{[23;1;00;2]}(M) = \{m_{13} - m_{12} - \delta(M)\}\chi_-(M)C_2\left(M\left(\begin{smallmatrix}1\,0\\0\end{smallmatrix}\right)[-1]\right),$$

and

$$\begin{aligned}
B^{\lambda}_{[23;1;11;0]}(M) =& (m_{13} - m_{12} - 1)(m_{13} - m_{12})(m_{22} - m_{33}),\\
B^{\lambda}_{[23;1;11;1]}(M) =& -(m_{13} - m_{12})\bar{E}(M)\\
&+ \{m_{13} - m_{12} - \delta(M) - 1\}\chi_-(M)(m_{13} - m_{12})(m_{22} - m_{33}),\\
B^{\lambda}_{[23;1;11;2]}(M) =& -\{m_{13} - m_{12} - \delta(M)\}\chi_-(M)\bar{E}(M),\\
B^{\lambda}_{[23;1;01;0]}(M) =& -(m_{13} - m_{12} - 1)(m_{13} - m_{12})(m_{22} - m_{33}),\\
B^{\lambda}_{[23;1;01;1]}(M) =& (m_{13} - m_{12})\bar{F}(M)\\
&- \{m_{13} - m_{12} - \delta(M)\}\chi_-^{(-1)}(M)(m_{13} - m_{12})(m_{22} - m_{33})\\
&- (m_{13} - m_{12})(m_{13} - m_{22} + 1),\\
B^{\lambda}_{[23;1;01;2]}(M) =& -(m_{13} - m_{12} + 1)C_2(M)\chi_+(M)\\
&+ \{m_{13} - m_{12} - \delta(M) + 1\}\chi_-^{(-1)}(M)\bar{F}(M) + C_2(M),\\
B^{\lambda}_{[23;1;01;3]}(M) =& -\{m_{13} - m_{12} - \delta(M) + 2\}\chi_-^{(-1)}(M)C_2(M)\chi_+(M),\\
B^{\lambda}_{[23;1;00;0]}(M) =& -(m_{13} - m_{12} - 1)(m_{13} - m_{12})(m_{13} - m_{22} + 1),\\
B^{\lambda}_{[23;1;00;1]}(M) =& (m_{13} - m_{12})C_2(M)\\
&- \{m_{13} - m_{12} - \delta(M) - 1\}\chi_-(M)(m_{13} - m_{12})(m_{13} - m_{22} + 1),\\
B^{\lambda}_{[23;1;00;2]}(M) =& \{m_{13} - m_{12} - \delta(M)\}\chi_-(M)C_2(M).
\end{aligned}$$

We have

$$\begin{aligned}
A^{\lambda}_{[23;1;11;1]}(M) - B^{\lambda}_{[23;1;11;1]}(M) =& -(m_{13} - m_{12})(m_{13} + m_{23} - m_{33} - 2m_{12} + m_{11})\chi_-(M)\\
&+ (m_{13} - m_{12})\chi_-(M)(m_{13} - m_{12} - \delta(M) + m_{22} - m_{33})\\
=& 0.
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[23;1;11;1]}(M) = B^{\lambda}_{[23;1;11;1]}(M)$. Here we use the relation

$$\bar{E}\left(M\left(\begin{smallmatrix}1\,0\\0\end{smallmatrix}\right)\right) = \bar{E}(M) + (m_{13} + m_{23} - m_{33} - 2m_{12} + m_{11})\chi_-(M). \tag{4.28}$$

By the relation

$$\bar{E}\left(M\left(\begin{smallmatrix}1\,0\\0\end{smallmatrix}\right)[-1]\right)\chi_-(M) = \bar{E}(M)\chi_-(M), \tag{4.29}$$

Hence we obtain

$$A^{\lambda}_{[23;1;11;2]}(M) = B^{\lambda}_{[23;1;11;2]}(M).$$

We have

$$\begin{aligned}
&A^{\lambda}_{[23;1;01;1]}(M) - B^{\lambda}_{[23;1;01;1]}(M)\\
&= (m_{13} - m_{12})\{(m_{13} - m_{12} - \delta(M))\chi_-(M)\\
&\quad - (m_{13} - m_{12})(m_{22} - m_{33})(\chi_-^{(-1)} - \chi_-(M)) + (m_{13} - m_{22} + 1)\}\\
&\quad - \{m_{13} - m_{12} - \delta(M)\}\chi_-(M)(m_{13} - m_{12})(m_{22} - m_{33} + 1)\\
&\quad + \{m_{13} - m_{12} - \delta(M)\}\chi_-^{(-1)}(M)(m_{13} - m_{12})(m_{22} - m_{33})\\
&\quad + (m_{13} - m_{12})(m_{13} - m_{22} + 1)\\
&= (m_{13} - m_{12})(m_{22} - m_{33})\delta(M)(\chi_-(M) - \chi_-^{(-1)}(M))\\
&= 0.
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[23;1;01;1]}(M)=B^{\lambda}_{[23;1;01;1]}(M)$. Here we use the relation

$$\begin{aligned}\bar{F}\left(M\left(\begin{smallmatrix}1&0\\&0\end{smallmatrix}\right)\right)=&\bar{F}(M)+(m_{13}-m_{12}-\delta(M))\chi_-(M)\\&-(m_{13}-m_{12})(m_{22}-m_{33})(\chi_-^{(-1)}-\chi_-(M))+(m_{13}-m_{22}+1).\end{aligned}\tag{4.30}$$

We have

$$\begin{aligned}A^{\lambda}_{[23;1;01;2]}(M)=&-(m_{13}-m_{12})(m_{11}-m_{22})(m_{23}-m_{22})\chi_+^{(-1)}(M)\\&-\{m_{13}-m_{12}-\delta(M)\}(m_{12}-m_{23})(m_{12}-m_{11})\chi_-(M)\\=&-(m_{13}-m_{12})C_2(M)\chi_+^{(-1)}(M)-\{m_{13}-m_{12}-\delta(M)\}C_2(M)\chi_-(M)\\=&-(m_{13}-m_{12})C_2(M)+\delta(M)C_2(M)\chi_-(M),\\B^{\lambda}_{[23;1;01;2]}(M)=&-(m_{13}-m_{12}+1)C_2(M)\chi_+(M)\\&-\{m_{13}-m_{12}-\delta(M)+1\}C_2(M)\chi_-^{(-1)}(M)+C_2(M)\\=&-(m_{13}-m_{12}+1)C_2(M)+C_2(M)\delta(M)\chi_-^{(-1)}(M)+C_2(M)\\=&-(m_{13}-m_{12})C_2(M)+\delta(M)C_2(M)\chi_-(M).\end{aligned}$$

Hence we obtain $A^{\lambda}_{[23;1;01;2]}(M)=B^{\lambda}_{[23;1;01;2]}(M)$. Here we use the relation (4.26).

We have

$$\begin{aligned}A^{\lambda}_{[23;1;01;3]}(M)&=-\{m_{13}-m_{12}-\delta(M)\}C_2\left(M\left(\begin{smallmatrix}1&0\\&0\end{smallmatrix}\right)[-1]\right)\chi_-(M)\chi_+^{(-1)}(M)=0,\\B^{\lambda}_{[23;1;01;3]}(M)&=-\{m_{13}-m_{12}-\delta(M)+2\}C_2(M)\chi_-^{(-1)}(M)\chi_+(M)=0.\end{aligned}$$

Hence we obtain $A^{\lambda}_{[23;1;01;3]}(M)=B^{\lambda}_{[23;1;01;3]}(M)$.

We have

$$\begin{aligned}&A^{\lambda}_{[23;1;00;1]}(M)-B^{\lambda}_{[23;1;00;1]}(M)\\&=(m_{13}-m_{12})(m_{12}-m_{22}+1+\delta(M))\chi_-(M)\\&\quad-\{m_{13}-m_{12}-\delta(M)\}\chi_-(M)(m_{13}-m_{12})(m_{13}-m_{22})\\&\quad+\{m_{13}-m_{12}-\delta(M)-1\}\chi_-(M)(m_{13}-m_{12})(m_{13}-m_{22}+1)\\&=0.\end{aligned}$$

Hence we obtain $A^{\lambda}_{[23;1;00;1]}(M)=B^{\lambda}_{[23;1;00;1]}(M)$. Here we use the relation

$$C_2\left(M\left(\begin{smallmatrix}1&0\\&0\end{smallmatrix}\right)\right)=C_2(M)+(m_{12}-m_{22}+1+\delta(M))\chi_-(M).\tag{4.31}$$

We have

$$\begin{aligned}&A^{\lambda}_{[23;1;00;2]}(M)\\&=\{m_{13}-m_{12}-\delta(M)\}\chi_-(M)C_2(M)\\&=B^{\lambda}_{[23;1;00;2]}(M).\end{aligned}$$

Hence we obtain $A^{\lambda}_{[23;1;00;2]}(M)=B^{\lambda}_{[23;1;00;2]}(M)$.

It is trivial that the equations (4.17) hold for $(j,k,l)=(1,1,0),\ (0,1,0)$ and $(0,0,0)$.

- the proof of $E_{32}\circ i^{\lambda}_{\mathbf{e}_1}=i^{\lambda}_{\mathbf{e}_1}\circ E_{32}$

We have

$$\begin{aligned}A^{\lambda}_{[32;1;11;0]}(M)=&(m_{22}-m_{33})(m_{13}-m_{12})(m_{22}-m_{33}-1),\\A^{\lambda}_{[32;1;11;1]}(M)=&-(m_{22}-m_{33})\bar{E}\left(M\left(\begin{smallmatrix}0&-1\\&0\end{smallmatrix}\right)\right)\\&+\{m_{22}-m_{33}+\delta(M)\}\chi_+(M)(m_{13}-m_{12}+1)(m_{22}-m_{33}),\\A^{\lambda}_{[32;1;11;2]}(M)=&-\{m_{22}-m_{33}+\delta(M)\}\chi_+(M)\bar{E}\left(M\left(\begin{smallmatrix}0&-1\\&0\end{smallmatrix}\right)[-1]\right),\end{aligned}$$

$$\begin{aligned}
A^{\lambda}_{[32;1;01;0]}(M) =& -(m_{22}-m_{33})(m_{13}-m_{12})(m_{22}-m_{33}-1),\\
A^{\lambda}_{[32;1;01;1]}(M) =& (m_{22}-m_{33})\bar{F}\left(M\tbinom{0\;-1}{0}\right)\\
&- \{m_{22}-m_{33}+\delta(M)\}\chi_{+}(M)(m_{13}-m_{12}+1)(m_{22}-m_{33}),\\
A^{\lambda}_{[32;1;01;2]}(M) =& -(m_{22}-m_{33})C_2\left(M\tbinom{0\;-1}{0}\right)\chi_{+}\left(M\tbinom{0\;-1}{0}\right)\\
&+ \{m_{22}-m_{33}+\delta(M)\}\chi_{+}(M)\bar{F}\left(M\tbinom{0\;-1}{0}[-1]\right),\\
A^{\lambda}_{[32;1;01;3]}(M) =& -\{m_{22}-m_{33}+\delta(M)\}\chi_{+}(M)C_2\left(M\tbinom{0\;-1}{0}[-1]\right)\chi_{+}\left(M\tbinom{0\;-1}{0}[-1]\right),\\
A^{\lambda}_{[32;1;00;0]}(M) =& -(m_{22}-m_{33})(m_{13}-m_{12})(m_{13}-m_{22}+2),\\
A^{\lambda}_{[32;1;00;1]}(M) =& (m_{22}-m_{33})C_2\left(M\tbinom{0\;-1}{0}\right)\\
&- \{m_{22}-m_{33}+\delta(M)\}\chi_{+}(M)(m_{13}-m_{12}+1)(m_{13}-m_{22}+1),\\
A^{\lambda}_{[32;1;00;2]}(M) =& \{m_{22}-m_{33}+\delta(M)\}\chi_{+}(M)C_2\left(M\tbinom{0\;-1}{0}[-1]\right),
\end{aligned}$$

and

$$\begin{aligned}
B^{\lambda}_{[32;1;11;0]}(M) =& (m_{22}-m_{33}-1)(m_{13}-m_{12})(m_{22}-m_{33}),\\
B^{\lambda}_{[32;1;11;1]}(M) =& -(m_{22}-m_{33})\bar{E}(M)\\
&+ \{m_{22}-m_{33}+\delta(M)-1\}\chi_{+}(M)(m_{13}-m_{12})(m_{22}-m_{33}),\\
B^{\lambda}_{[32;1;11;2]}(M) =& -\{m_{22}-m_{33}+\delta(M)\}\chi_{+}(M)\bar{E}(M),\\
B^{\lambda}_{[32;1;01;0]}(M) =& -(m_{22}-m_{33}-1)(m_{13}-m_{12})(m_{22}-m_{33}),\\
B^{\lambda}_{[32;1;01;1]}(M) =& (m_{22}-m_{33})\bar{F}(M)\\
&- \{m_{22}-m_{33}+\delta(M)-2\}\chi^{(1)}_{+}(M)(m_{13}-m_{12})(m_{22}-m_{33}),\\
B^{\lambda}_{[32;1;01;2]}(M) =& -(m_{22}-m_{33}+1)C_2(M)\chi_{+}(M)\\
&+ \{m_{22}-m_{33}+\delta(M)-1\}\chi^{(1)}_{+}(M)\bar{F}(M),\\
B^{\lambda}_{[32;1;01;3]}(M) =& -\{m_{22}-m_{33}+\delta(M)\}\chi^{(1)}_{+}(M)C_2(M)\chi_{+}(M),\\
B^{\lambda}_{[32;1;00;0]}(M) =& -(m_{22}-m_{33})(m_{13}-m_{12})(m_{13}-m_{22}+1)-(m_{13}-m_{12})(m_{22}-m_{33}),\\
B^{\lambda}_{[32;1;00;1]}(M) =& (m_{22}-m_{33}+1)C_2(M)\\
&- \{m_{22}-m_{33}+\delta(M)\}\chi_{+}(M)(m_{13}-m_{12})(m_{13}-m_{22}+1)+\bar{F}(M),\\
B^{\lambda}_{[32;1;00;2]}(M) =& \{m_{22}-m_{33}+\delta(M)+1\}\chi_{+}(M)C_2(M)-C_2(M)\chi_{+}(M).
\end{aligned}$$

We have

$$\begin{aligned}
&A^{\lambda}_{[32;1;11;1]}(M)-B^{\lambda}_{[32;1;11;1]}(M)\\
&=-(m_{22}-m_{33})(m_{13}-m_{33}-m_{12}+m_{22}+\delta(M))\chi_{+}(M)\\
&\quad+\{m_{22}-m_{33}+\delta(M)\}\chi_{+}(M)(m_{13}-m_{12}+1)(m_{22}-m_{33})\\
&\quad-\{m_{22}-m_{33}+\delta(M)-1\}\chi_{+}(M)(m_{13}-m_{12})(m_{22}-m_{33})\\
&=0.
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[32;1;11;1]}(M)=B^{\lambda}_{[32;1;11;1]}(M)$. Here we use the relation

$$\bar{E}\left(M\tbinom{0\;-1}{0}\right)=\bar{E}(M)+(m_{13}-m_{33}-m_{12}+m_{22}+\delta(M))\chi_{+}(M). \tag{4.32}$$

By the relation

$$\bar{E}\left(M\tbinom{0\;-1}{0}[-1]\right)\chi_{+}(M)=\bar{E}(M)\chi_{+}(M), \tag{4.33}$$

we obtain

$$A^{\lambda}_{[32;1;11;2]}(M) = B^{\lambda}_{[32;1;11;2]}(M).$$

We have

$$\begin{aligned}
&A^{\lambda}_{[32;1;01;1]}(M) - B^{\lambda}_{[32;1;01;1]}(M) \\
&= (m_{22}-m_{33})\{-(m_{13}-m_{12})(m_{22}-m_{33}-1)\chi_+^{(1)}(M) \\
&\quad + (m_{13}-m_{12}+1)(m_{22}-m_{33}+1)\chi_+(M) + (\delta(M)-1)\chi_+(M) \\
&\quad - (m_{22}-m_{33}+\delta(M))\chi_+(M)(m_{13}-m_{12}+1) \\
&\quad + (m_{22}-m_{33}+\delta(M)-2)\chi_+^{(1)}(M)(m_{13}-m_{12})\} \\
&= (m_{22}-m_{33})(m_{13}-m_{12})(\delta(M)-1)(\chi_+^{(1)}(M)-\chi_+(M)) \\
&= 0.
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[32;1;01;1]}(M) = B^{\lambda}_{[32;1;01;1]}(M)$. Here we use the relation

$$\begin{aligned}
\bar{F}\left(M\left(\begin{smallmatrix}0&-1\\&0\end{smallmatrix}\right)\right) =& \bar{F}(M) - (m_{13}-m_{12})(m_{22}-m_{33}-1)\chi_+^{(1)} \\
&+ \{(m_{13}-m_{12}+1)(m_{22}-m_{33}+1)+\delta(M)-1\}\chi_+(M).
\end{aligned} \tag{4.34}$$

We have

$$\begin{aligned}
&A^{\lambda}_{[32;1;01;2]}(M) - B^{\lambda}_{[32;1;01;2]}(M) \\
&= (m_{22}-m_{33}+\delta(M))(\bar{F}\left(M\left(\begin{smallmatrix}0&-1\\&0\end{smallmatrix}\right)[-1]\right)\chi_+(M) - \bar{F}(M)\chi_+^{(1)}(M)) \\
&\quad + \bar{F}(M)\chi_+^{(1)}(M) + (m_{22}-m_{33}+1)C_2(M)\chi_+(M) - (m_{22}-m_{33})C_2\left(M\left(\begin{smallmatrix}0&-1\\&0\end{smallmatrix}\right)\right)\chi_+^{(1)}(M) \\
&= (m_{22}-m_{33}+\delta(M))\{(m_{13}-m_{22}+1)\chi_+^{(1)}(M) + C_2(M)\chi_+^{(1)}(M)\} \\
&\quad - (\delta(M)-1)C_2(M)\chi_+(M) + \bar{F}(M)\chi_+^{(1)}(M) - (m_{22}-m_{33})C_2\left(M\left(\begin{smallmatrix}0&-1\\&0\end{smallmatrix}\right)\right)\chi_+^{(1)}(M) \\
&= (m_{22}-m_{33}+\delta(M))(m_{13}-m_{22}+1)\chi_+^{(1)}(M) \\
&\quad - (m_{22}-m_{33})(C_2\left(M\left(\begin{smallmatrix}0&-1\\&0\end{smallmatrix}\right)\right) - C_2(M))\chi_+^{(1)}(M) + C_2(M)\chi_+^{(1)}(M) + \bar{F}(M)\chi_+^{(1)}(M) \\
&= \chi_+^{(1)}(M)\{C_2(M) + (m_{13}-m_{12})(m_{22}-m_{33}) + (m_{13}-m_{33}+1)\delta(M) + \bar{F}(M)\chi_+^{(1)}(M)\} \\
&= 0.
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[32;1;01;2]}(M) = B^{\lambda}_{[32;1;01;2]}(M)$. Here we use the relations

$$\begin{aligned}
\bar{F}\left(M\left(\begin{smallmatrix}0&-1\\&0\end{smallmatrix}\right)[-1]\right)\chi_+(M) =& \bar{F}(M)\chi_+^{(1)}(M) + (m_{13}-m_{22}+1)\chi_+^{(1)}(M) \\
&+ C_2(M)(\chi_+^{(1)}(M)-\chi_+(M)), \\
C_2\left(M\left(\begin{smallmatrix}0&-1\\&0\end{smallmatrix}\right)\right)\chi_+^{(1)}(M) =& (C_2(M)+m_{12}-m_{22}+1-\delta(M))\chi_+^{(1)}(M).
\end{aligned} \tag{4.35}$$

We have

$$\begin{aligned}
A^{\lambda}_{[32;1;01;3]}(M) =& -\{m_{22}-m_{33}+\delta(M)\}\chi_+(M)(m_{11}-m_{22})(m_{23}-m_{22})\chi_+^{(1)}(M) \\
=& B^{\lambda}_{[32;1;01;3]}(M).
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[32;1;01;3]}(M) = B^{\lambda}_{[32;1;01;3]}(M)$.

We have

$$\begin{aligned}
&A^{\lambda}_{[32;1;00;1]}(M) - B^{\lambda}_{[32;1;00;1]}(M) \\
&= (m_{22}-m_{33})(m_{12}-m_{22}+1-\delta(M))\chi_+(M) - C_2(M) \\
&\quad - \{m_{22}-m_{33}+\delta(M)\}\chi_+(M)(m_{13}-m_{22}+1) - \bar{F}(M) \\
&= -C_2(M) - \chi_+(M)\{(m_{13}-m_{12})(m_{22}-m_{33}) + (m_{13}-m_{33}+1)\delta(M)\} - \bar{F}(M) \\
&= 0.
\end{aligned}$$

Hence we obtain $A^{\lambda}_{[32;1;00;1]}(M) = B^{\lambda}_{[32;1;00;1]}(M)$. Here we use the relation

$$C_2\left(M\left(\begin{smallmatrix}0\,-1\\0\end{smallmatrix}\right)\right) = C_2(M) + (m_{12} - m_{22} + 1 - \delta(M))\chi_+(M). \tag{4.36}$$

We have

$$A^{\lambda}_{[32;1;00;2]}(M) = B^{\lambda}_{[32;1;00;2]}(M).$$

Hence we obtain $A^{\lambda}_{[32;1;00;2]}(M) = B^{\lambda}_{[32;1;00;2]}(M)$.

It is trivial that the equations (4.17) hold for $(j,k,l) = (1,1,0),\ (0,1,0)$ and $(0,0,0)$.

□

4.2. **Irreducible decompositions of $V_\lambda \otimes_{\mathbf{C}} V_{2\mathbf{e}_1}$ and $V_\lambda \otimes_{\mathbf{C}} V_{-2\mathbf{e}_3}$.** For a vector space W, we denote by id_W the identity map of W. We denote $\lambda \pm (\mathbf{e}_i + \mathbf{e}_j)$ $(1 \le i \le j \le 3)$ by $\lambda[\pm ij]$ for the sake of simplicity. Generically the tensor product $V_\lambda \otimes_{\mathbf{C}} V_{2\mathbf{e}_1}$ has six irreducible components: $V_{\lambda[+ij]}$, $1 \le i \le j \le 3$. Here some components may vanish.

For $1 \le i \le j \le 3$, let $i^{\lambda}_{\mathbf{e}_i+\mathbf{e}_j}$ be a non-zero generator of $\mathrm{Hom}_K(V_{\lambda[+ij]}, V_\lambda \otimes_{\mathbf{C}} V_{2\mathbf{e}_1})$, which is unique up to scalar multiple if $V_\lambda \otimes_{\mathbf{C}} V_{2\mathbf{e}_1}$ has a non-zero $\tau_{\lambda[+ij]}$-isotypic component.

Lemma 4.3. *We define a linear map $P_{\mathbf{e}_1}\colon V_{\mathbf{e}_1} \otimes_{\mathbf{C}} V_{\mathbf{e}_1} \to V_{2\mathbf{e}_1}$ by*

$$P_{\mathbf{e}_1}\left(f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\1\end{smallmatrix}\right) \otimes f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\1\end{smallmatrix}\right)\right) = f\left(\begin{smallmatrix}2\,0\,0\\2\,0\\2\end{smallmatrix}\right), \qquad P_{\mathbf{e}_1}\left(f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\1\end{smallmatrix}\right) \otimes f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\0\end{smallmatrix}\right)\right) = f\left(\begin{smallmatrix}2\,0\,0\\2\,0\\1\end{smallmatrix}\right),$$

$$P_{\mathbf{e}_1}\left(f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\1\end{smallmatrix}\right) \otimes f\left(\begin{smallmatrix}1\,0\,0\\0\,0\\0\end{smallmatrix}\right)\right) = f\left(\begin{smallmatrix}2\,0\,0\\1\,0\\1\end{smallmatrix}\right), \qquad P_{\mathbf{e}_1}\left(f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\0\end{smallmatrix}\right) \otimes f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\1\end{smallmatrix}\right)\right) = f\left(\begin{smallmatrix}2\,0\,0\\2\,0\\1\end{smallmatrix}\right),$$

$$P_{\mathbf{e}_1}\left(f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\0\end{smallmatrix}\right) \otimes f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\0\end{smallmatrix}\right)\right) = f\left(\begin{smallmatrix}2\,0\,0\\2\,0\\0\end{smallmatrix}\right), \qquad P_{\mathbf{e}_1}\left(f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\0\end{smallmatrix}\right) \otimes f\left(\begin{smallmatrix}1\,0\,0\\0\,0\\0\end{smallmatrix}\right)\right) = f\left(\begin{smallmatrix}2\,0\,0\\1\,0\\0\end{smallmatrix}\right),$$

$$P_{\mathbf{e}_1}\left(f\left(\begin{smallmatrix}1\,0\,0\\0\,0\\0\end{smallmatrix}\right) \otimes f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\1\end{smallmatrix}\right)\right) = f\left(\begin{smallmatrix}2\,0\,0\\1\,0\\1\end{smallmatrix}\right), \qquad P_{\mathbf{e}_1}\left(f\left(\begin{smallmatrix}1\,0\,0\\0\,0\\0\end{smallmatrix}\right) \otimes f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\0\end{smallmatrix}\right)\right) = f\left(\begin{smallmatrix}2\,0\,0\\1\,0\\0\end{smallmatrix}\right),$$

$$P_{\mathbf{e}_1}\left(f\left(\begin{smallmatrix}1\,0\,0\\0\,0\\0\end{smallmatrix}\right) \otimes f\left(\begin{smallmatrix}1\,0\,0\\0\,0\\0\end{smallmatrix}\right)\right) = f\left(\begin{smallmatrix}2\,0\,0\\0\,0\\0\end{smallmatrix}\right).$$

Then $P_{\mathbf{e}_1}$ is a non-zero generator of $\mathrm{Hom}_K(V_{\mathbf{e}_1} \otimes_{\mathbf{C}} V_{\mathbf{e}_1}, V_{2\mathbf{e}_1})$.

Proof. Since $V_{\mathbf{e}_1} \otimes_{\mathbf{C}} V_{\mathbf{e}_1} \simeq V_{2\mathbf{e}_1} \oplus V_{\mathbf{e}_1+\mathbf{e}_2}$, in order to prove this lemma, it suffices to check $P_{\mathbf{e}_1} \circ i^{\mathbf{e}_1}_{\mathbf{e}_2} = 0$ and that $P_{\mathbf{e}_1} \circ i^{\mathbf{e}_1}_{\mathbf{e}_1}$ agree with the identity map up to scalar multiple. To consider the case $\lambda = \mathbf{e}_1$ in Proposition 4.2, we obtain the explicit expressions of $i^{\mathbf{e}_1}_{\mathbf{e}_1}$ and $i^{\mathbf{e}_1}_{\mathbf{e}_2}$ in terms of the monomial basis as follows:

$$i^{\mathbf{e}_1}_{\mathbf{e}_1}\left(f\left(\begin{smallmatrix}2\,0\,0\\2\,0\\2\end{smallmatrix}\right)\right) = -6f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\1\end{smallmatrix}\right) \otimes f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\1\end{smallmatrix}\right),$$

$$i^{\mathbf{e}_1}_{\mathbf{e}_1}\left(f\left(\begin{smallmatrix}2\,0\,0\\2\,0\\1\end{smallmatrix}\right)\right) = -3\left\{f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\1\end{smallmatrix}\right) \otimes f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\0\end{smallmatrix}\right) + f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\0\end{smallmatrix}\right) \otimes f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\1\end{smallmatrix}\right)\right\},$$

$$i^{\mathbf{e}_1}_{\mathbf{e}_1}\left(f\left(\begin{smallmatrix}2\,0\,0\\2\,0\\0\end{smallmatrix}\right)\right) = -6f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\0\end{smallmatrix}\right) \otimes f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\0\end{smallmatrix}\right),$$

$$i^{\mathbf{e}_1}_{\mathbf{e}_1}\left(f\left(\begin{smallmatrix}2\,0\,0\\1\,0\\1\end{smallmatrix}\right)\right) = -3\left\{f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\1\end{smallmatrix}\right) \otimes f\left(\begin{smallmatrix}1\,0\,0\\0\,0\\0\end{smallmatrix}\right) + f\left(\begin{smallmatrix}1\,0\,0\\0\,0\\0\end{smallmatrix}\right) \otimes f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\1\end{smallmatrix}\right)\right\},$$

$$i^{\mathbf{e}_1}_{\mathbf{e}_1}\left(f\left(\begin{smallmatrix}2\,0\,0\\1\,0\\0\end{smallmatrix}\right)\right) = -3\left\{f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\0\end{smallmatrix}\right) \otimes f\left(\begin{smallmatrix}1\,0\,0\\0\,0\\0\end{smallmatrix}\right) + f\left(\begin{smallmatrix}1\,0\,0\\0\,0\\0\end{smallmatrix}\right) \otimes f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\0\end{smallmatrix}\right)\right\},$$

$$i^{\mathbf{e}_1}_{\mathbf{e}_1}\left(f\left(\begin{smallmatrix}2\,0\,0\\0\,0\\0\end{smallmatrix}\right)\right) = -6f\left(\begin{smallmatrix}1\,0\,0\\0\,0\\0\end{smallmatrix}\right) \otimes f\left(\begin{smallmatrix}1\,0\,0\\0\,0\\0\end{smallmatrix}\right),$$

$$i^{\mathbf{e}_1}_{\mathbf{e}_2}\left(f\left(\begin{smallmatrix}1\,1\,0\\1\,1\\1\end{smallmatrix}\right)\right) = f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\0\end{smallmatrix}\right) \otimes f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\1\end{smallmatrix}\right) - f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\1\end{smallmatrix}\right) \otimes f\left(\begin{smallmatrix}1\,0\,0\\1\,0\\0\end{smallmatrix}\right),$$

$$
\begin{aligned}
i^{\mathbf{e}_1}_{\mathbf{e}_2}\left(f\left(\begin{smallmatrix}1\;1\;0\\1\;0\\1\end{smallmatrix}\right)\right) =& f\left(\begin{smallmatrix}1\;0\;0\\0\;0\\0\end{smallmatrix}\right)\otimes f\left(\begin{smallmatrix}1\;0\;0\\1\;0\\1\end{smallmatrix}\right)-f\left(\begin{smallmatrix}1\;0\;0\\1\;0\\1\end{smallmatrix}\right)\otimes f\left(\begin{smallmatrix}1\;0\;0\\0\;0\\0\end{smallmatrix}\right),\\
i^{\mathbf{e}_1}_{\mathbf{e}_2}\left(f\left(\begin{smallmatrix}1\;1\;0\\1\;0\\0\end{smallmatrix}\right)\right) =& f\left(\begin{smallmatrix}1\;0\;0\\0\;0\\0\end{smallmatrix}\right)\otimes f\left(\begin{smallmatrix}1\;0\;0\\1\;0\\0\end{smallmatrix}\right)-f\left(\begin{smallmatrix}1\;0\;0\\1\;0\\0\end{smallmatrix}\right)\otimes f\left(\begin{smallmatrix}1\;0\;0\\0\;0\\0\end{smallmatrix}\right).
\end{aligned}
$$

By direct computation, we can easily check $P_{\mathbf{e}_1}\circ i^{\mathbf{e}_1}_{\mathbf{e}_1}=-6\,\mathrm{id}_{V_{2\mathbf{e}_1}}$ and $P_{\mathbf{e}_1}\circ i^{\mathbf{e}_1}_{\mathbf{e}_2}=0$. □

By a composition of the projectors in Lemma 4.3 and the injectors in Proposition 4.2, we obtain following formulas.

Proposition 4.4. *For $1\le i\le j\le 3$ and a G-pattern M of type $\lambda[+ij]$, the image of the monomial basis $f(M)$ by the injector $i^{\lambda}_{\mathbf{e}_i+\mathbf{e}_j}\colon V_{\lambda[+ij]}\to V_\lambda\otimes_{\mathbf{C}}V_{2\mathbf{e}_1}$ is given by*

$$
i^{\lambda}_{\mathbf{e}_i+\mathbf{e}_j}(f(M))=\sum_{0\le k\le l\le 2}\left\{\sum_{m=0}^{r_{[ij;kl]}}c^{\lambda}_{[ij;kl;m]}(M)\,f\left(M\left(\begin{smallmatrix}-\mathbf{e}_i-\mathbf{e}_j\\0\;-l\\-k\end{smallmatrix}\right)[-m]\right)\right\}\otimes f\left(\begin{smallmatrix}2\;0\;0\\l\;0\\k\end{smallmatrix}\right).
$$

In the right hand side of the above formula, we put $f(M')=0$ if M' is a triangular array which does not satisfy the condition (3.1) of G-patterns.

The explicit expressions of the coefficients are given by the following formulas.

Formula 1: *The coefficients of the injector $i^{\lambda}_{2\mathbf{e}_1}\colon V_{\lambda[+11]}\to V_\lambda\otimes_{\mathbf{C}}V_{2\mathbf{e}_1}$ are given as follows: $(r_{[11;22]},r_{[11;12]},r_{[11;11]},r_{[11;02]},r_{[11;01]},r_{[11;00]})=(2,3,2,4,3,2)$ and*

$$
\begin{aligned}
c^{\lambda}_{[11;22;0]}(M)=&(m_{13}-m_{12})(m_{13}-m_{12}-1)(m_{22}-m_{33})(m_{22}-m_{33}-1),\\
c^{\lambda}_{[11;22;1]}(M)=&-2(m_{13}-m_{12})(m_{22}-m_{33})(\bar{E}(M)-C_1(M)),\\
c^{\lambda}_{[11;22;2]}(M)=&\bar{E}(M)(C_1(M)-1)(m_{13}-m_{33}-\bar{C}_1(M)),\\
c^{\lambda}_{[11;12;0]}(M)=&-2(m_{13}-m_{12})(m_{13}-m_{12}-1)(m_{22}-m_{33})(m_{22}-m_{33}-1),\\
c^{\lambda}_{[11;12;1]}(M)=&2(m_{13}-m_{12})(m_{22}-m_{33})\left\{\bar{F}\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;-1\\-1\end{smallmatrix}\right)\right)+\bar{E}(M)\right\},\\
c^{\lambda}_{[11;12;2]}(M)=&-2\left\{\bar{E}(M)\bar{F}\left(M\left(\begin{smallmatrix}-1\;0\;0\\-1\;0\\-1\end{smallmatrix}\right)\right)\right.\\
&\qquad\left.+(m_{13}-m_{12})(m_{22}-m_{33})C_1(M)(\bar{C}_1(M)+1)\chi_+(M)\right\},\\
c^{\lambda}_{[11;12;3]}(M)=&2(C_1(M)-1)\bar{C}_1(M)\bar{E}(M)\chi_+(M),\\
c^{\lambda}_{[11;11;0]}(M)=&-2(m_{13}-m_{12})(m_{13}-m_{12}-1)(m_{22}-m_{33})(m_{13}-m_{22}+1),\\
c^{\lambda}_{[11;11;1]}(M)=&2(m_{13}-m_{12})\left\{(m_{22}-m_{33})C_1(M)(\bar{C}_1(M)+1)+(m_{13}-m_{22})\bar{E}(M)\right\},\\
c^{\lambda}_{[11;11;2]}(M)=&-2\bar{E}(M)(C_1(M)-1)\bar{C}_1(M),\\
c^{\lambda}_{[11;02;0]}(M)=&(m_{13}-m_{12})(m_{13}-m_{12}-1)(m_{22}-m_{33})(m_{22}-m_{33}-1),\\
c^{\lambda}_{[11;02;1]}(M)=&-(m_{13}-m_{12})(m_{22}-m_{33})\left(\bar{F}(M)+\bar{F}\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;-1\\0\end{smallmatrix}\right)\right)\right),\\
c^{\lambda}_{[11;02;2]}(M)=&(m_{13}-m_{12}+1)(m_{22}-m_{33}+1)C_2(M)\chi_+(M)\\
&\qquad+(m_{13}-m_{12})(m_{22}-m_{33})(C_1(M)+1)(\bar{C}_1(M)+1)\chi^{(1)}_+(M)\\
&\qquad+\bar{F}(M)\bar{F}\left(M\left(\begin{smallmatrix}-1\;0\;0\\-1\;0\\0\end{smallmatrix}\right)\right),\\
c^{\lambda}_{[11;02;3]}(M)=&-C_2(M)\left\{\chi^{(1)}_+(M)\bar{F}(M)+\chi_+(M)\bar{F}\left(M\left(\begin{smallmatrix}-1\;0\;0\\-2\;1\\0\end{smallmatrix}\right)\right)\right\},\\
c^{\lambda}_{[11;02;4]}(M)=&C_2(M)(C_1(M)-1)(\bar{C}_1(M)-1)\chi^{(1)}_+(M),\\
c^{\lambda}_{[11;01;0]}(M)=&2(m_{13}-m_{12})(m_{13}-m_{12}-1)(m_{22}-m_{33})(m_{13}-m_{22}+1),
\end{aligned}
$$

$$
\begin{aligned}
c^{\lambda}_{[11;01;1]}(M) =& -2(m_{13}-m_{12})\left\{(m_{13}-m_{22})\bar{F}(M) + (m_{22}-m_{33})C_2\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & -1 \\ & & 0 \end{smallmatrix}\right)\right)\right\},\\
c^{\lambda}_{[11;01;2]}(M) =& 2\left\{C_2\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & -1 & 0 \\ & & 0 \end{smallmatrix}\right)\right)\bar{F}(M)\right.\\
&\left. +\chi_{+}(M)(m_{13}-m_{12}+1)(m_{13}-m_{22}-1)C_2(M)\right\},\\
c^{\lambda}_{[11;01;3]}(M) =& -2C_2(M)(C_1(M)-1)(\bar{C}_1(M)-1)\chi_{+}(M),\\
c^{\lambda}_{[11;00;0]}(M) =& (m_{13}-m_{12})(m_{13}-m_{12}-1)(m_{13}-m_{22}+1)(m_{13}-m_{22}),\\
c^{\lambda}_{[11;00;1]}(M) =& -2(m_{13}-m_{12})(m_{13}-m_{22})C_2(M),\\
c^{\lambda}_{[11;00;2]}(M) =& C_2(M)(C_1(M)-1)(\bar{C}_1(M)-1).
\end{aligned}
$$

Formula 2: *The coefficients of the injector* $i^{\lambda}_{2\mathbf{e}_2}\colon V_{\lambda[+22]} \to V_{\lambda} \otimes_{\mathbf{C}} V_{2\mathbf{e}_1}$ *are given as follows:* $(r_{[22;22]}, r_{[22;12]}, r_{[22;11]}, r_{[22;02]}, r_{[22;01]}, r_{[22;00]}) = (2,2,2,2,2,2)$ *and*

$$
\begin{aligned}
c^{\lambda}_{[22;22;0]}(M) =& (m_{22}-m_{33})(m_{22}-m_{33}-1),\\
c^{\lambda}_{[22;22;1]}(M) =& -(m_{22}-m_{33})\{\bar{D}(M)\chi_{-}(M) + (\bar{D}(M)+2)\chi^{(1)}_{-}(M)\},\\
c^{\lambda}_{[22;22;2]}(M) =& \bar{D}(M)(\bar{D}(M)+1)\chi^{(1)}_{-}(M),\\
c^{\lambda}_{[22;12;0]}(M) =& -2(m_{22}-m_{33})(m_{22}-m_{33}-1),\\
c^{\lambda}_{[22;12;1]}(M) =& 2(m_{22}-m_{33})\{\bar{C}_1(M) + (\bar{D}(M)+1)\chi_{-}(M)\},\\
c^{\lambda}_{[22;12;2]}(M) =& -2\bar{C}_1(M)\bar{D}(M)\chi_{-}(M),\\
c^{\lambda}_{[22;11;0]}(M) =& -2(m_{22}-m_{33})(m_{23}-m_{22}),\\
c^{\lambda}_{[22;11;1]}(M) =& 2\{\bar{D}(M)(m_{23}-m_{22}-1)\chi_{-}(M) - (m_{22}-m_{33})(\bar{C}_1(M)+1)\chi^{(1)}_{-}(M)\},\\
c^{\lambda}_{[22;11;2]}(M) =& 2\bar{C}_1(M)\bar{D}(M)\chi^{(1)}_{-}(M),\\
c^{\lambda}_{[22;02;0]}(M) =& (m_{22}-m_{33})(m_{22}-m_{33}-1),\\
c^{\lambda}_{[22;02;1]}(M) =& -2(m_{22}-m_{33})\bar{C}_1(M),\\
c^{\lambda}_{[22;02;2]}(M) =& \bar{C}_1(M)(\bar{C}_1(M)-1),\\
c^{\lambda}_{[22;01;0]}(M) =& 2(m_{22}-m_{33})(m_{23}-m_{22}),\\
c^{\lambda}_{[22;01;1]}(M) =& 2\bar{C}_1(M)\{(m_{22}-m_{33})\chi_{-}(M) - (m_{23}-m_{22}-1)\},\\
c^{\lambda}_{[22;01;2]}(M) =& -2\bar{C}_1(M)(\bar{C}_1(M)-1)\chi_{-}(M),\\
c^{\lambda}_{[22;00;0]}(M) =& (m_{23}-m_{22})(m_{23}-m_{22}-1),\\
c^{\lambda}_{[22;00;1]}(M) =& \bar{C}_1(M)\{(m_{23}-m_{22}-2)\chi_{-}(M) + (m_{23}-m_{22})\chi^{(1)}_{-}(M)\},\\
c^{\lambda}_{[22;00;2]}(M) =& \bar{C}_1(M)(\bar{C}_1(M)-1)\chi^{(1)}_{-}(M).
\end{aligned}
$$

Formula 3: *The coefficients of the injector* $i^{\lambda}_{2\mathbf{e}_3}\colon V_{\lambda[+33]} \to V_{\lambda} \otimes_{\mathbf{C}} V_{2\mathbf{e}_1}$ *are given as follows:* $(r_{[33;22]}, r_{[33;12]}, r_{[33;11]}, r_{[33;02]}, r_{[33;01]}, r_{[33;00]}) = (0,1,0,2,1,0)$ *and*

$$
\begin{aligned}
c^{\lambda}_{[33;22;0]}(M) =& 1, & c^{\lambda}_{[33;12;0]}(M) =& -2,\\
c^{\lambda}_{[33;12;1]}(M) =& -2\chi_{+}(M), & c^{\lambda}_{[33;11;0]}(M) =& 2,\\
c^{\lambda}_{[33;02;0]}(M) =& 1, & c^{\lambda}_{[33;02;1]}(M) =& \chi^{(1)}_{+}(M) + \chi_{+}(M),\\
c^{\lambda}_{[33;02;2]}(M) =& \chi^{(1)}_{+}(M), & c^{\lambda}_{[33;01;0]}(M) =& -2,\\
c^{\lambda}_{[33;01;1]}(M) =& -2\chi_{+}(M), & c^{\lambda}_{[33;00;0]}(M) =& 1.
\end{aligned}
$$

Formula 4: *The coefficients of the injector* $i^{\lambda}_{\mathbf{e}_1+\mathbf{e}_2}\colon V_{\lambda[+12]} \to V_\lambda \otimes_{\mathbf{C}} V_{2\mathbf{e}_1}$ *are given as follows* $(r_{[12;22]}, r_{[12;12]}, r_{[12;11]}, r_{[12;02]}, r_{[12;01]}, r_{[12;00]}) = (2,2,2,3,2,2)$ *and*

$$
\begin{aligned}
c^{\lambda}_{[12;22;0]}(M) =&(m_{13}-m_{12})(m_{22}-m_{33})(m_{22}-m_{33}-1),\\
c^{\lambda}_{[12;22;1]}(M) =&-(m_{22}-m_{33})\left\{\bar{E}(M)+\chi_-(M)(m_{13}-m_{12})(\bar{D}(M)+1)\right\},\\
c^{\lambda}_{[12;22;2]}(M) =&\bar{D}(M)\bar{E}(M)\chi_-(M),\\
c^{\lambda}_{[12;12;0]}(M) =&-2(m_{13}-m_{12})(m_{22}-m_{33})(m_{22}-m_{33}-1),\\
c^{\lambda}_{[12;12;1]}(M) =&(m_{22}-m_{33})\left\{\bar{E}(M)+\bar{F}(M)\right.\\
&\left.+(m_{13}-m_{12})\left\{\bar{C}_1(M)+1+\bar{D}(M)(1-\chi_+(M))\right\}\right\},\\
c^{\lambda}_{[12;12;2]}(M) =&-\bar{C}_1(M)\bar{E}(M)-C_2(M)\left(1-\bar{D}(M)+\delta(M)\chi_+(M)\right),\\
c^{\lambda}_{[12;11;0]}(M) =&(m_{13}-m_{12})(m_{22}-m_{33})(2m_{22}-m_{13}-m_{23}-2),\\
c^{\lambda}_{[12;11;1]}(M) =&\bar{E}(M)(m_{23}-m_{22})+C_2(M)(m_{22}-m_{33}+1)\\
&+(m_{13}-m_{12})\chi_-(M)\left\{\bar{D}(M)(m_{13}-m_{22}+1)-(m_{22}-m_{33})(\bar{C}_1(M)+1)\right\},\\
c^{\lambda}_{[12;11;2]}(M) =&C_2(M)\chi_-(M)(m_{13}-m_{33}+2-\bar{C}_1(M)-\bar{D}(M)),\\
c^{\lambda}_{[12;02;0]}(M) =&(m_{13}-m_{12})(m_{22}-m_{33})(m_{22}-m_{33}-1),\\
c^{\lambda}_{[12;02;1]}(M) =&-(m_{22}-m_{33})\left\{\bar{F}(M)+(m_{13}-m_{12})(\bar{C}_1(M)+\chi_+(M))\right\},\\
c^{\lambda}_{[12;02;2]}(M) =&(\bar{C}_1(M)+\chi_+(M)-1)\bar{F}(M)+(m_{22}-m_{33}+1)C_2(M)\chi_+(M),\\
c^{\lambda}_{[12;02;3]}(M) =&-C_2(M)(\bar{C}_1(M)-1)\chi_+(M),\\
c^{\lambda}_{[12;01;0]}(M) =&-(m_{13}-m_{12})(m_{22}-m_{33})(2m_{22}-m_{13}-m_{23}-2),\\
c^{\lambda}_{[12;01;1]}(M) =&(m_{13}-m_{12})\bar{C}_1(M)\left\{(m_{22}-m_{33})(1-\chi_+(M))-(m_{13}-m_{22}+1)\right\}\\
&-(m_{23}-m_{22})\bar{F}(M)-(m_{22}-m_{33}+1)C_2(M),\\
c^{\lambda}_{[12;01;2]}(M) =&2C_2(M)(\bar{C}_1(M)-1),\\
c^{\lambda}_{[12;00;0]}(M) =&(m_{13}-m_{12})(m_{13}-m_{22}+1)(m_{23}-m_{22}),\\
c^{\lambda}_{[12;00;1]}(M) =&(m_{13}-m_{12})(m_{13}-m_{22}+1)\bar{C}_1(M)\chi_-(M)-(m_{23}-m_{22}-1)C_2(M),\\
c^{\lambda}_{[12;00;2]}(M) =&-C_2(M)(\bar{C}_1(M)-1)\chi_-(M).
\end{aligned}
$$

Formula 5: *The coefficients of the injector* $i^{\lambda}_{\mathbf{e}_1+\mathbf{e}_3}\colon V_{\lambda[+13]} \to V_\lambda \otimes_{\mathbf{C}} V_{2\mathbf{e}_1}$ *are given as follows:* $(r_{[13;22]}, r_{[13;12]}, r_{[13;11]}, r_{[13;02]}, r_{[13;01]}, r_{[13;00]}) = (1,2,1,3,2,1)$ *and*

$$
\begin{aligned}
c^{\lambda}_{[13;22;0]}(M) =&(m_{13}-m_{12})(m_{22}-m_{33}),\\
c^{\lambda}_{[13;22;1]}(M) =&-\bar{E}(M),\\
c^{\lambda}_{[13;12;0]}(M) =&-2(m_{13}-m_{12})(m_{22}-m_{33}),\\
c^{\lambda}_{[13;12;1]}(M) =&\bar{E}(M)+\bar{F}(M)-(m_{13}-m_{12})(m_{22}-m_{33})\chi_+(M),\\
c^{\lambda}_{[13;12;2]}(M) =&(\bar{E}(M)-C_2(M))\chi_+(M),\\
c^{\lambda}_{[13;11;0]}(M) =&(m_{13}-m_{12})(2m_{22}-m_{13}-m_{33}-1),\\
c^{\lambda}_{[13;11;1]}(M) =&C_2(M)-\bar{E}(M),\\
c^{\lambda}_{[13;02;0]}(M) =&(m_{13}-m_{12})(m_{22}-m_{33}),\\
c^{\lambda}_{[13;02;1]}(M) =&(m_{13}-m_{12})(m_{22}-m_{33})\chi^{(1)}_+(M)-\bar{F}(M),
\end{aligned}
$$

$$
\begin{aligned}
c^{\lambda}_{[13;02;2]}(M) =& C_2(M)\chi_+(M) - \bar{F}(M)\chi^{(1)}_+(M),\\
c^{\lambda}_{[13;02;3]}(M) =& C_2(M)\chi^{(1)}_+(M),\\
c^{\lambda}_{[13;01;0]}(M) =& -(m_{13} - m_{12})(2m_{22} - m_{13} - m_{23} - 1),\\
c^{\lambda}_{[13;01;1]}(M) =& \bar{F}(M) - C_2(M) + (m_{13} - m_{12})(m_{13} - m_{22} + 1)\chi_+(M),\\
c^{\lambda}_{[13;01;2]}(M) =& -2C_2(M)\chi_+(M),\\
c^{\lambda}_{[13;00;0]}(M) =& -(m_{13} - m_{12})(m_{13} - m_{22} + 1),\\
c^{\lambda}_{[13;00;1]}(M) =& C_2(M).
\end{aligned}
$$

Formula 6: *The coefficients of the injector* $i^{\lambda}_{\mathbf{e}_2+\mathbf{e}_3} \colon V_{\lambda[+23]} \to V_\lambda \otimes_{\mathbf{C}} V_{2\mathbf{e}_1}$ *are given as follows:* $(r_{[23;22]}, r_{[23;12]}, r_{[23;11]}, r_{[23;02]}, r_{[23;01]}, r_{[23;00]}) = (1, 1, 1, 2, 1, 1)$ *and*

$$
\begin{aligned}
c^{\lambda}_{[23;22;0]}(M) =& m_{22} - m_{33}, & c^{\lambda}_{[23;22;1]}(M) =& -\bar{D}(M)\chi_-(M),\\
c^{\lambda}_{[23;12;0]}(M) =& -2(m_{22} - m_{33}), & c^{\lambda}_{[23;12;1]}(M) =& \bar{C}_1(M) - (m_{22} - m_{33}) + \delta(M)\chi_-(M),\\
c^{\lambda}_{[23;11;0]}(M) =& 2m_{22} - m_{23} - m_{33}, & c^{\lambda}_{[23;11;1]}(M) =& -(\bar{C}_1(M) + \bar{D}(M))\chi_-(M),\\
c^{\lambda}_{[23;02;0]}(M) =& m_{22} - m_{33}, & c^{\lambda}_{[23;02;1]}(M) =& -\{\bar{C}_1(M) - (m_{22} - m_{33})\chi_+(M)\},\\
c^{\lambda}_{[23;02;2]}(M) =& -\bar{C}_1(M)\chi_+(M), & c^{\lambda}_{[23;01;0]}(M) =& -(2m_{22} - m_{23} - m_{33}),\\
c^{\lambda}_{[23;01;1]}(M) =& 2\bar{C}_1(M), & c^{\lambda}_{[23;00;0]}(M) =& -(m_{23} - m_{22}),\\
c^{\lambda}_{[23;00;1]}(M) =& -\bar{C}_1(M)\chi_-(M).
\end{aligned}
$$

Proof. For $1 \leq i \leq j \leq 3$, let $i^{\lambda}_{\mathbf{e}_i+\mathbf{e}_j} \colon V_{\lambda[+ij]} \to V_\lambda \otimes_{\mathbf{C}} V_{2\mathbf{e}_1}$ be the composite of three K-homomorphisms

$$
\begin{aligned}
i^{\lambda+\mathbf{e}_j}_{\mathbf{e}_i} &\colon V_{\lambda[+ij]} \to V_{\lambda+\mathbf{e}_j} \otimes_{\mathbf{C}} V_{\mathbf{e}_1}\\
i^{\lambda}_{\mathbf{e}_j} \otimes_{\mathbf{C}} \mathrm{id}_{V_{\mathbf{e}_1}} &\colon V_{\lambda+\mathbf{e}_j} \otimes_{\mathbf{C}} V_{\mathbf{e}_1} \to V_\lambda \otimes_{\mathbf{C}} V_{\mathbf{e}_1} \otimes_{\mathbf{C}} V_{\mathbf{e}_1}\\
\mathrm{id}_{V_\lambda} \otimes_{\mathbf{C}} P_{\mathbf{e}_1} &\colon V_\lambda \otimes_{\mathbf{C}} V_{\mathbf{e}_1} \otimes_{\mathbf{C}} V_{\mathbf{e}_1} \to V_\lambda \otimes_{\mathbf{C}} V_{2\mathbf{e}_1}.
\end{aligned}
$$

Then $i^{\lambda}_{\mathbf{e}_i+\mathbf{e}_j}$ is an element of $\mathrm{Hom}_K(V_{\lambda[+ij]}, V_\lambda \otimes_{\mathbf{C}} V_{2\mathbf{e}_1})$. By direct computation, we confirm that $i^{\lambda}_{\mathbf{e}_i+\mathbf{e}_j}$ is non-zero and obtain the explicit expression of this K-homomorphism.

$$
\begin{aligned}
&i^{\lambda}_{\mathbf{e}_i+\mathbf{e}_j}(f(M)) = (\mathrm{id}_{V_\lambda} \otimes_{\mathbf{C}} P_{\mathbf{e}_1}) \circ (i^{\lambda}_{\mathbf{e}_j} \otimes_{\mathbf{C}} \mathrm{id}_{V_{\mathbf{e}_1}}) \circ i^{\lambda+\mathbf{e}_j}_{\mathbf{e}_i}(f(M))\\
&= \sum_{0\leq k\leq l\leq 1} \sum_{m=0}^{r_{[i;kl]}} c^{\lambda}_{[i;kl;m]}(M)\, (\mathrm{id}_{V_\lambda} \otimes_{\mathbf{C}} P_{\mathbf{e}_1}) \circ (i^{\lambda}_{\mathbf{e}_j} \otimes_{\mathbf{C}} \mathrm{id}_{V_{\mathbf{e}_1}}) \left(f\left(M \left(\begin{smallmatrix} -\mathbf{e}_i \\ 0 \; -l \\ -k \end{smallmatrix} \right) [-m]\right) \otimes f\left(\begin{smallmatrix} 1\,0\,0 \\ l\;0 \\ k \end{smallmatrix} \right)\right)\\
&= \sum_{0\leq k\leq l\leq 1} \sum_{m=0}^{r_{[i;kl]}} c^{\lambda}_{[i;kl;m]}(M) \left\{ \sum_{0\leq p\leq q\leq 1} \sum_{r=0}^{r_{[j;pq]}} c^{\lambda}_{[j;pq;r]}\left(M \left(\begin{smallmatrix} -\mathbf{e}_i \\ 0 \; -l \\ -k \end{smallmatrix} \right) [-m]\right)\right.\\
&\qquad \left. \times f\left(M \left(\begin{smallmatrix} -\mathbf{e}_i-\mathbf{e}_j \\ 0 \; -l-q \\ -k-p \end{smallmatrix} \right) [-m-r]\right) \right\} \otimes P_{\mathbf{e}_1}\left(f\left(\begin{smallmatrix} 1\,0\,0 \\ q\;0 \\ p \end{smallmatrix} \right) \otimes f\left(\begin{smallmatrix} 1\,0\,0 \\ l\;0 \\ k \end{smallmatrix} \right)\right)\\
&= \sum_{0\leq s\leq t\leq 2} \left\{ \sum_{u=0}^{R_{[ij;st]}} c^{\lambda}_{[ij;st;u]}(M)\, f\left(M \left(\begin{smallmatrix} -\mathbf{e}_i-\mathbf{e}_j \\ 0 \; -t \\ -s \end{smallmatrix} \right) [-u]\right) \right\} \otimes f\left(\begin{smallmatrix} 2\,0\,0 \\ t\;0 \\ s \end{smallmatrix} \right),
\end{aligned}
$$

where

$$c^{\lambda}_{[ij;st;u]}(M)=\sum_{\substack{0\le p\le q\le 1,\\ 0\le k\le l\le 1,\\ p+k=s,\ q+l=t}}\ \sum_{\substack{0\le m\le r_{[i;kl]},\\ 0\le r\le r_{[j;pq]}.\\ m+r=u}} c^{\lambda}_{[i;kl;m]}(M)\, c^{\lambda}_{[j;pq;r]}\left(M\left(\begin{smallmatrix}-\mathbf{e}_i\\ 0\ -l\\ -k\end{smallmatrix}\right)[-m]\right),$$

and

$$R_{[ij;st]}=\max\{r_{[i;kl]}+r_{[j;pq]}\mid 0\le k\le l\le 1,\ 0\le p\le q\le 1,\ p+k=s,\ q+l=t\}.$$

Now we simplify each coefficients by direct computation.

First, we compute the case of formula 1.

- the case $(s,t)=(2,2)$.

We have

$$\begin{aligned}
c^{\lambda}_{[11;22;0]}(M)&=c^{\lambda}_{[1;11;0]}(M)\,c^{\lambda}_{[1;11;0]}\left(M\left(\begin{smallmatrix}-1\,0\,0\\ 0\,-1\\ -1\end{smallmatrix}\right)\right)\\
&=(m_{13}-m_{12})(m_{22}-m_{33})(m_{13}-m_{12}-1)(m_{22}-m_{33}-1),\\
c^{\lambda}_{[11;22;1]}(M)&=c^{\lambda}_{[1;11;0]}(M)\,c^{\lambda}_{[1;11;1]}\left(M\left(\begin{smallmatrix}-1\,0\,0\\ 0\,-1\\ -1\end{smallmatrix}\right)\right)+c^{\lambda}_{[1;11;1]}(M)\,c^{\lambda}_{[1;11;0]}\left(M\left(\begin{smallmatrix}-1\,0\,0\\ 0\,-1\\ -1\end{smallmatrix}\right)[-1]\right)\\
&=-(m_{13}-m_{12})(m_{22}-m_{33})\bar{E}\left(M\left(\begin{smallmatrix}-1\,0\,0\\ 0\,-1\\ -1\end{smallmatrix}\right)\right)-\bar{E}(M)(m_{13}-m_{12})(m_{22}-m_{33})\\
&=-2(m_{13}-m_{12})(m_{22}-m_{33})(\bar{E}(M)-C_1(M)),\\
c^{\lambda}_{[11;22;2]}(M)&=c^{\lambda}_{[1;11;1]}(M)\,c^{\lambda}_{[1;11;1]}\left(M\left(\begin{smallmatrix}-1\,0\,0\\ 0\,-1\\ -1\end{smallmatrix}\right)[-1]\right)\\
&=\bar{E}(M)\bar{E}\left(M\left(\begin{smallmatrix}-1\,0\,0\\ 0\,-1\\ -1\end{smallmatrix}\right)[-1]\right)\\
&=\bar{E}(M)(C_1(M)-1)(m_{13}-m_{33}-\bar{C}_1(M)).
\end{aligned}$$

Here we use the relations:

$$\bar{E}\left(M\left(\begin{smallmatrix}-1\,0\,0\\ 0\,-1\\ -1\end{smallmatrix}\right)\right)=\bar{E}(M)-2C_1(M),\tag{4.37}$$

$$C_1\left(M\left(\begin{smallmatrix}-1\,0\,0\\ 0\,-1\\ -1\end{smallmatrix}\right)\right)=C_1(M),\quad \bar{C}_1\left(M\left(\begin{smallmatrix}-1\,0\,0\\ 0\,-1\\ -1\end{smallmatrix}\right)\right)=\bar{C}_1(M)+1,\tag{4.38}$$

$$C_1\left(M[-1]\right)=C_1(M)-1,\quad \bar{C}_1\left(M[-1]\right)=\bar{C}_1(M)-1.\tag{4.39}$$

- the case $(s,t)=(1,2)$.

We have

$$\begin{aligned}
c^{\lambda}_{[11;12;0]}(M)&=c^{\lambda}_{[1;11;0]}(M)\,c^{\lambda}_{[1;01;0]}\left(M\left(\begin{smallmatrix}-1\,0\,0\\ 0\,-1\\ -1\end{smallmatrix}\right)\right)+c^{\lambda}_{[1;01;0]}(M)\,c^{\lambda}_{[1;11;0]}\left(M\left(\begin{smallmatrix}-1\,0\,0\\ 0\,-1\\ 0\end{smallmatrix}\right)\right)\\
&=-2(m_{13}-m_{12})(m_{22}-m_{33})(m_{13}-m_{12}-1)(m_{22}-m_{33}-1),\\
c^{\lambda}_{[11;12;1]}(M)&=c^{\lambda}_{[1;11;0]}(M)\,c^{\lambda}_{[1;01;1]}\left(M\left(\begin{smallmatrix}-1\,0\,0\\ 0\,-1\\ -1\end{smallmatrix}\right)\right)+c^{\lambda}_{[1;01;1]}(M)\,c^{\lambda}_{[1;11;0]}\left(M\left(\begin{smallmatrix}-1\,0\,0\\ 0\,-1\\ 0\end{smallmatrix}\right)[-1]\right)\\
&\quad+c^{\lambda}_{[1;01;0]}(M)\,c^{\lambda}_{[1;11;1]}\left(M\left(\begin{smallmatrix}-1\,0\,0\\ 0\,-1\\ 0\end{smallmatrix}\right)\right)+c^{\lambda}_{[1;11;1]}(M)\,c^{\lambda}_{[1;01;0]}\left(M\left(\begin{smallmatrix}-1\,0\,0\\ 0\,-1\\ -1\end{smallmatrix}\right)[-1]\right)\\
&=(m_{13}-m_{12})(m_{22}-m_{33})\bar{F}\left(M\left(\begin{smallmatrix}-1\,0\,0\\ 0\,-1\\ -1\end{smallmatrix}\right)\right)+\bar{F}(M)(m_{13}-m_{12})(m_{22}-m_{33})\\
&\quad+(m_{13}-m_{12})(m_{22}-m_{33})\bar{E}\left(M\left(\begin{smallmatrix}-1\,0\,0\\ 0\,-1\\ 0\end{smallmatrix}\right)\right)+\bar{E}(M)(m_{13}-m_{12})(m_{22}-m_{33})\\
&=2(m_{13}-m_{12})(m_{22}-m_{33})\left(\bar{F}\left(M\left(\begin{smallmatrix}-1\,0\,0\\ 0\,-1\\ -1\end{smallmatrix}\right)\right)+\bar{E}(M)\right),\\
c^{\lambda}_{[11;12;2]}(M)&=c^{\lambda}_{[1;11;1]}(M)\,c^{\lambda}_{[1;01;1]}\left(M\left(\begin{smallmatrix}-1\,0\,0\\ 0\,-1\\ -1\end{smallmatrix}\right)[-1]\right)+c^{\lambda}_{[1;01;1]}(M)\,c^{\lambda}_{[1;11;1]}\left(M\left(\begin{smallmatrix}-1\,0\,0\\ 0\,-1\\ 0\end{smallmatrix}\right)[-1]\right)
\end{aligned}$$

$$
\begin{aligned}
&+ c^{\lambda}_{[1;11;0]}(M)\, c^{\lambda}_{[1;01;2]}\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ -1\\ -1\end{smallmatrix}\right)\right) + c^{\lambda}_{[1;01;2]}(M)\, c^{\lambda}_{[1;11;0]}\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ -1\\ 0\end{smallmatrix}\right)[-2]\right)\\
&= -\bar{E}(M)\bar{F}\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ -1\\ -1\end{smallmatrix}\right)[-1]\right) - \bar{F}(M)\bar{E}\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ -1\\ 0\end{smallmatrix}\right)[-1]\right)\\
&\quad - (m_{13}-m_{12})(m_{22}-m_{33})C_2\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ -1\\ -1\end{smallmatrix}\right)\right)\chi_+\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ -1\\ -1\end{smallmatrix}\right)\right)\\
&\quad - C_2(M)\chi_+(M)(m_{13}-m_{12}+1)(m_{22}-m_{33}+1)\\
&= -2\bar{E}(M)\bar{F}\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ -1\\ -1\end{smallmatrix}\right)[-1]\right) - (m_{13}-m_{12})(m_{22}-m_{33})C_1(M)\chi_+(M)\\
&\quad + C_2(M)\chi_+(M)(m_{13}-m_{33}+1-m_{12}+m_{22})\\
&\quad - (m_{13}-m_{12})(m_{22}-m_{33})C_1(M)(\bar{C}_1(M)+1)\chi_+(M)\\
&\quad - C_2(M)\chi_+(M)(m_{13}-m_{12}+1)(m_{22}-m_{33}+1)\\
&= -2\Bigg\{\bar{E}(M)\bar{F}\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ -1\\ -1\end{smallmatrix}\right)[-1]\right)\\
&\quad + (m_{13}-m_{12})(m_{22}-m_{33})C_1(M)(\bar{C}_1(M)+1)\chi_+(M)\Bigg\},
\end{aligned}
$$

$$
\begin{aligned}
c^{\lambda}_{[11;12;3]}(M) &= c^{\lambda}_{[1;11;1]}(M)\, c^{\lambda}_{[1;01;2]}\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ -1\\ 0\end{smallmatrix}\right)[-1]\right) + c^{\lambda}_{[1;01;2]}(M)\, c^{\lambda}_{[1;11;1]}\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ -1\\ 0\end{smallmatrix}\right)[-2]\right)\\
&= \bar{E}(M)C_2\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ -1\\ 0\end{smallmatrix}\right)[-1]\right)\chi_+\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ -1\\ 0\end{smallmatrix}\right)[-1]\right)\\
&\quad + C_2(M)\chi_+(M)\bar{E}\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ -1\\ 0\end{smallmatrix}\right)[-2]\right)\\
&= 2(C_1(M)-1)\bar{C}_1(M)\bar{E}(M)\chi_+(M).
\end{aligned}
$$

Here we use the relations:

$$
\bar{F}\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ -1\\ -1\end{smallmatrix}\right)\right) - \bar{F}(M) = \bar{E}\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ -1\\ 0\end{smallmatrix}\right)\right) - \bar{E}(M), \tag{4.40}
$$

$$
\begin{aligned}
&\bar{F}(M)\bar{E}\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ -1\\ 0\end{smallmatrix}\right)[-1]\right)\\
&= \bar{E}(M)\bar{F}\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ -1\\ -1\end{smallmatrix}\right)[-1]\right) + (m_{13}-m_{12})(m_{22}-m_{33})C_1(M)\chi_+(M)\\
&\quad - C_2(M)\chi_+(M)(m_{13}-m_{33}+1-m_{12}+m_{22}),
\end{aligned} \tag{4.41}
$$

$$
C_2\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ -1\\ -1\end{smallmatrix}\right)\right)\chi_+\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ -1\\ -1\end{smallmatrix}\right)\right) = C_1(M)(\bar{C}_1(M)+1)\chi_+(M), \tag{4.42}
$$

$$
C_2\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ -1\\ 0\end{smallmatrix}\right)[-1]\right)\chi_+\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ -1\\ 0\end{smallmatrix}\right)[-1]\right) = (C_1(M)-1)\bar{C}_1(M)\chi_+(M), \tag{4.43}
$$

$$
C_2(M)\chi_+(M)\bar{E}\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ -1\\ 0\end{smallmatrix}\right)[-2]\right) = (C_1(M)-1)\bar{C}_1(M)\bar{E}(M)\chi_+(M). \tag{4.44}
$$

- the case $(s, t) = (1, 1)$.
  We have

$$
\begin{aligned}
c^{\lambda}_{[11;11;0]}(M) &= c^{\lambda}_{[1;11;0]}(M)\, c^{\lambda}_{[1;00;0]}\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ -1\\ -1\end{smallmatrix}\right)\right) + c^{\lambda}_{[1;00;0]}(M)\, c^{\lambda}_{[1;11;0]}\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ 0\\ 0\end{smallmatrix}\right)\right)\\
&= -(m_{13}-m_{12})(m_{22}-m_{33})(m_{13}-m_{12}-1)(m_{13}-m_{22}+1)\\
&\quad - (m_{13}-m_{12})(m_{13}-m_{22}+1)(m_{13}-m_{12}-1)(m_{22}-m_{33})\\
&= -2(m_{13}-m_{12})(m_{13}-m_{12}-1)(m_{22}-m_{33})(m_{13}-m_{22}+1),\\
c^{\lambda}_{[11;11;1]}(M) &= c^{\lambda}_{[1;11;0]}(M)\, c^{\lambda}_{[1;00;1]}\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ -1\\ -1\end{smallmatrix}\right)\right) + c^{\lambda}_{[1;00;1]}(M)\, c^{\lambda}_{[1;11;0]}\left(M\left(\begin{smallmatrix}-1\ 0\ 0\\ 0\ 0\\ 0\end{smallmatrix}\right)[-1]\right)
\end{aligned}
$$

$$
\begin{aligned}
&\quad + c^{\lambda}_{[1;00;0]}(M)\, c^{\lambda}_{[1;11;1]}\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;0\\0\end{smallmatrix}\right)\right) + c^{\lambda}_{[1;11;1]}(M)\, c^{\lambda}_{[1;00;0]}\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;-1\\-1\end{smallmatrix}\right)[-1]\right)\\
&= (m_{13}-m_{12})(m_{22}-m_{33})C_2\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;-1\\-1\end{smallmatrix}\right)\right)\\
&\quad + C_2(M)(m_{13}-m_{12})(m_{22}-m_{33}+1)\\
&\quad + (m_{13}-m_{12})(m_{13}-m_{22}+1)\bar{E}\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;0\\0\end{smallmatrix}\right)\right)\\
&\quad + \bar{E}(M)(m_{13}-m_{12})(m_{13}-m_{22})\\
&= (m_{13}-m_{12})\big\{(m_{22}-m_{33})C_1(M)(\bar{C}_1(M)+1) + C_2(M)(m_{22}-m_{33}+1)\\
&\quad + (m_{13}-m_{22}+1)(\bar{E}(M)-C_1(M)) + \bar{E}(M)(m_{13}-m_{22})\big\}\\
&= (m_{13}-m_{12})\big\{2(m_{22}-m_{33})C_1(M)(\bar{C}_1(M)+1)\\
&\quad + 2(m_{13}-m_{22})\bar{E}(M) + \bar{E}(M) - C_1(M)(m_{13}-m_{33}+1-\bar{C}_1(M))\big\}\\
&= 2(m_{13}-m_{12})\big\{(m_{22}-m_{33})C_1(M)(\bar{C}_1(M)+1) + (m_{13}-m_{22})\bar{E}(M)\big\},
\end{aligned}
$$

$$
\begin{aligned}
c^{\lambda}_{[11;11;2]}(M) &= c^{\lambda}_{[1;11;1]}(M)\, c^{\lambda}_{[1;00;1]}\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;-1\\-1\end{smallmatrix}\right)[-1]\right) + c^{\lambda}_{[1;00;1]}(M)\, c^{\lambda}_{[1;11;1]}\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;0\\0\end{smallmatrix}\right)[-1]\right)\\
&= -\bar{E}(M)C_2\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;-1\\-1\end{smallmatrix}\right)[-1]\right) - C_2(M)\bar{E}\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;0\\0\end{smallmatrix}\right)[-1]\right)\\
&= -2\bar{E}(M)(C_1(M)-1)\bar{C}_1(M).
\end{aligned}
$$

Here we use the relations:

$$
\bar{E}\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;0\\0\end{smallmatrix}\right)\right) = \bar{E}(M) - C_1(M), \tag{4.45}
$$

$$
C_2(M)\bar{E}\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;0\\0\end{smallmatrix}\right)[-1]\right) = (C_1(M)-1)\bar{C}_1(M)\bar{E}(M), \tag{4.46}
$$

(4.38) and (4.39).

- the case $(s,t)=(0,2)$.

We have

$$
\begin{aligned}
c^{\lambda}_{[11;02;0]}(M) &= c^{\lambda}_{[1;01;0]}(M)\, c^{\lambda}_{[1;01;0]}\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;-1\\0\end{smallmatrix}\right)\right)\\
&= (m_{13}-m_{12})(m_{22}-m_{33})(m_{13}-m_{12}-1)(m_{22}-m_{33}-1),\\
c^{\lambda}_{[11;02;1]}(M) &= c^{\lambda}_{[1;01;0]}(M)\, c^{\lambda}_{[1;01;1]}\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;-1\\0\end{smallmatrix}\right)\right) + c^{\lambda}_{[1;01;1]}(M)\, c^{\lambda}_{[1;01;0]}\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;-1\\0\end{smallmatrix}\right)[-1]\right)\\
&= -(m_{13}-m_{12})(m_{22}-m_{33})\left(\bar{F}\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;-1\\0\end{smallmatrix}\right)\right) + \bar{F}(M)\right),\\
c^{\lambda}_{[11;02;2]}(M) &= c^{\lambda}_{[1;01;0]}(M)\, c^{\lambda}_{[1;01;2]}\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;-1\\0\end{smallmatrix}\right)\right) + c^{\lambda}_{[1;01;1]}(M)\, c^{\lambda}_{[1;01;1]}\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;-1\\0\end{smallmatrix}\right)[-1]\right)\\
&\quad + c^{\lambda}_{[1;01;2]}(M)\, c^{\lambda}_{[1;01;0]}\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;-1\\0\end{smallmatrix}\right)[-2]\right)\\
&= (m_{13}-m_{12})(m_{22}-m_{33})C_2\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;-1\\0\end{smallmatrix}\right)\right)\chi_{+}\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;-1\\0\end{smallmatrix}\right)\right)\\
&\quad + \bar{F}(M)\bar{F}\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;-1\\0\end{smallmatrix}\right)[-1]\right) + C_2(M)\chi_{+}(M)(m_{13}-m_{12}+1)(m_{22}-m_{33}+1)\\
&= (m_{13}-m_{12}+1)(m_{22}-m_{33}+1)C_2(M)\chi_{+}(M)\\
&\quad + (m_{13}-m_{12})(m_{22}-m_{33})(C_1(M)+1)(\bar{C}_1(M)+1)\chi_{+}^{(1)}(M)\\
&\quad + \bar{F}(M)\bar{F}\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;-1\\0\end{smallmatrix}\right)[-1]\right),
\end{aligned}
$$

$$\begin{aligned}
c^{\lambda}_{[11;02;3]}(M) &= c^{\lambda}_{[1;01;1]}(M)\,c^{\lambda}_{[1;01;2]}\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1&\\&0&\end{smallmatrix}\right)[-1]\right)+c^{\lambda}_{[1;01;2]}(M)\,c^{\lambda}_{[1;01;1]}\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1&\\&0&\end{smallmatrix}\right)[-2]\right)\\
&= -\bar{F}(M)C_2\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1&\\&0&\end{smallmatrix}\right)[-1]\right)\chi_+\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1&\\&0&\end{smallmatrix}\right)[-1]\right)\\
&\quad - C_2(M)\chi_+(M)\bar{F}\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1&\\&0&\end{smallmatrix}\right)[-2]\right)\\
&= -C_2(M)\left(\chi_+^{(1)}(M)\bar{F}(M)+\chi_+(M)\bar{F}\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1&\\&0&\end{smallmatrix}\right)[-2]\right)\right),\\
c^{\lambda}_{[11;02;4]}(M) &= c^{\lambda}_{[1;01;2]}(M)\,c^{\lambda}_{[1;01;2]}\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1&\\&0&\end{smallmatrix}\right)[-2]\right)\\
&= C_2(M)\chi_+(M)C_2\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1&\\&0&\end{smallmatrix}\right)[-2]\right)\chi_+\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1&\\&0&\end{smallmatrix}\right)[-2]\right)\\
&= C_2(M)(C_1(M)-1)(\bar{C}_1(M)-1)\chi_+^{(1)}(M).
\end{aligned}$$

Here we use the equations:

$$C_2\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1&\\&0&\end{smallmatrix}\right)\right)\chi_+\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1&\\&0&\end{smallmatrix}\right)\right)=(C_1(M)+1)(\bar{C}_1(M)+1)\chi_+^{(1)}(M), \tag{4.47}$$

(4.5), (4.11) and (4.39).

- the case $(s,t)=(0,1)$.

We have

$$\begin{aligned}
c^{\lambda}_{[11;01;0]}(M) &= c^{\lambda}_{[1;01;0]}(M)\,c^{\lambda}_{[1;00;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1&\\&0&\end{smallmatrix}\right)\right)+c^{\lambda}_{[1;00;0]}(M)\,c^{\lambda}_{[1;01;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\0&0&\\&0&\end{smallmatrix}\right)\right)\\
&= (m_{13}-m_{12})(m_{22}-m_{33})(m_{13}-m_{12}-1)(m_{13}-m_{22}+1)\\
&\quad +(m_{13}-m_{12})(m_{13}-m_{22}+1)(m_{13}-m_{12}-1)(m_{22}-m_{33})\\
&= 2(m_{13}-m_{12})(m_{13}-m_{12}-1)(m_{22}-m_{33})(m_{13}-m_{22}+1),\\
c^{\lambda}_{[11;01;1]}(M) &= c^{\lambda}_{[1;01;0]}(M)\,c^{\lambda}_{[1;00;1]}\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1&\\&0&\end{smallmatrix}\right)\right)+c^{\lambda}_{[1;00;1]}(M)\,c^{\lambda}_{[1;01;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\0&0&\\&0&\end{smallmatrix}\right)[-1]\right)\\
&\quad + c^{\lambda}_{[1;00;0]}(M)\,c^{\lambda}_{[1;01;1]}\left(M\left(\begin{smallmatrix}-1&0&0\\0&0&\\&0&\end{smallmatrix}\right)\right)+c^{\lambda}_{[1;01;1]}(M)\,c^{\lambda}_{[1;00;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1&\\&0&\end{smallmatrix}\right)[-1]\right)\\
&= -(m_{13}-m_{12})(m_{22}-m_{33})C_2\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1&\\&0&\end{smallmatrix}\right)\right)\\
&\quad - C_2(M)(m_{13}-m_{12})(m_{22}-m_{33}+1)\\
&\quad - (m_{13}-m_{12})(m_{13}-m_{22}+1)\bar{F}\left(M\left(\begin{smallmatrix}-1&0&0\\0&0&\\&0&\end{smallmatrix}\right)\right)\\
&\quad - \bar{F}(M)(m_{13}-m_{12})(m_{13}-m_{22})\\
&= -(m_{13}-m_{12})\Big\{(m_{22}-m_{33})C_2\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1&\\&0&\end{smallmatrix}\right)\right)+C_2(M)(m_{22}-m_{33}+1)\\
&\quad + (m_{13}-m_{22}+1)\{\bar{F}(M)+(m_{22}-m_{33}+\delta(M))\chi_+(M)\}\\
&\quad + \bar{F}(M)(m_{13}-m_{22})\Big\}\\
&= -(m_{13}-m_{12})\Big\{(m_{22}-m_{33})\Big(C_2\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1&\\&0&\end{smallmatrix}\right)\right)\\
&\quad + C_2(M)+(m_{12}-m_{22}+1-\delta(M))\chi_+(M)\Big)+2(m_{13}-m_{22})\bar{F}(M)+\bar{F}(M)\\
&\quad + C_2(M)+\chi_+(M)\{(m_{13}-m_{12})(m_{22}-m_{33})+(m_{13}-m_{33}+1)\delta(M)\}\Big\}\\
&= -2(m_{13}-m_{12})\Big\{(m_{22}-m_{33})C_2\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1&\\&0&\end{smallmatrix}\right)\right)+(m_{13}-m_{22})\bar{F}(M)\Big\},
\end{aligned}$$

$$\begin{aligned}
c^{\lambda}_{[11;01;2]}(M) &= c^{\lambda}_{[1;01;1]}(M)\, c^{\lambda}_{[1;00;1]}\left(M\left(\begin{smallmatrix}-1\,0\,0\\0\,-1\\0\end{smallmatrix}\right)[-1]\right) + c^{\lambda}_{[1;00;1]}(M)\, c^{\lambda}_{[1;01;1]}\left(M\left(\begin{smallmatrix}-1\,0\,0\\0\,0\\0\end{smallmatrix}\right)[-1]\right)\\
&\quad + c^{\lambda}_{[1;01;2]}(M)\, c^{\lambda}_{[1;00;0]}\left(M\left(\begin{smallmatrix}-1\,0\,0\\0\,-1\\0\end{smallmatrix}\right)[-2]\right) + c^{\lambda}_{[1;00;0]}(M)\, c^{\lambda}_{[1;01;2]}\left(M\left(\begin{smallmatrix}-1\,0\,0\\0\,0\\0\end{smallmatrix}\right)\right)\\
&= \bar{F}(M)C_2\left(M\left(\begin{smallmatrix}-1\,0\,0\\0\,-1\\0\end{smallmatrix}\right)[-1]\right) + C_2(M)\bar{F}\left(M\left(\begin{smallmatrix}-1\,0\,0\\0\,0\\0\end{smallmatrix}\right)[-1]\right)\\
&\quad + C_2(M)\chi_+(M)(m_{13}-m_{12}+1)(m_{13}-m_{22}-1)\\
&\quad + (m_{13}-m_{12})(m_{13}-m_{22}+1)C_2\left(M\left(\begin{smallmatrix}-1\,0\,0\\0\,0\\0\end{smallmatrix}\right)\right)\chi_+\left(M\left(\begin{smallmatrix}-1\,0\,0\\0\,0\\0\end{smallmatrix}\right)\right)\\
&= \bar{F}(M)C_2\left(M\left(\begin{smallmatrix}-1\,0\,0\\0\,-1\\0\end{smallmatrix}\right)[-1]\right)\\
&\quad + C_2(M)\bigl\{\bar{F}(M)+m_{12}-m_{22}-1+\delta(M)-\chi_+(M)(m_{13}-m_{12}+\delta(M))\bigr\}\\
&\quad + C_2(M)\chi_+(M)\{(m_{13}-m_{12}+1)(m_{13}-m_{22}-1)\\
&\quad + (m_{13}-m_{12})(m_{13}-m_{22}+1)\}\\
&= 2\bar{F}(M)C_2\left(M\left(\begin{smallmatrix}-1\,0\,0\\0\,-1\\0\end{smallmatrix}\right)[-1]\right) - (m_{12}-m_{22}-1+\delta(M))C_2(M)(1-\chi_+(M))\\
&\quad + C_2(M)\bigl\{m_{12}-m_{22}-1+\delta(M)-\chi_+(M)(m_{13}-m_{12}+\delta(M))\bigr\}\\
&\quad + C_2(M)\chi_+(M)\{(m_{13}-m_{12}+1)(m_{13}-m_{22}-1)\\
&\quad + (m_{13}-m_{12})(m_{13}-m_{22}+1)\}\\
&= 2\{\bar{F}(M)C_2\left(M\left(\begin{smallmatrix}-1\,0\,0\\0\,-1\\0\end{smallmatrix}\right)[-1]\right)\\
&\quad + C_2(M)\chi_+(M)(m_{13}-m_{12}+1)(m_{13}-m_{22}-1)\},\\
c^{\lambda}_{[11;01;3]}(M) &= c^{\lambda}_{[1;00;1]}(M)\, c^{\lambda}_{[1;01;2]}\left(M\left(\begin{smallmatrix}-1\,0\,0\\0\,0\\0\end{smallmatrix}\right)[-1]\right) + c^{\lambda}_{[1;01;2]}(M)\, c^{\lambda}_{[1;00;1]}\left(M\left(\begin{smallmatrix}-1\,0\,0\\0\,-1\\0\end{smallmatrix}\right)[-2]\right)\\
&= -C_2(M)C_2\left(M\left(\begin{smallmatrix}-1\,0\,0\\0\,0\\0\end{smallmatrix}\right)[-1]\right)\chi_+\left(M\left(\begin{smallmatrix}-1\,0\,0\\0\,0\\0\end{smallmatrix}\right)[-1]\right)\\
&\quad - C_2(M)\chi_+(M)C_2\left(M\left(\begin{smallmatrix}-1\,0\,0\\0\,-1\\0\end{smallmatrix}\right)[-2]\right)\\
&= -2C_2(M)(C_1(M)-1)(\bar{C}_1(M)-1)\chi_+(M).
\end{aligned}$$

Here we use the relations:

$$\bar{F}\left(M\left(\begin{smallmatrix}-1\,0\,0\\0\,0\\0\end{smallmatrix}\right)\right) = \bar{F}(M) + (m_{22}-m_{33}+\delta(M))\chi_+(M), \tag{4.48}$$

$$\begin{aligned} C_2\left(M\left(\begin{smallmatrix}-1\,0\,0\\0\,-1\\0\end{smallmatrix}\right)\right) &= C_2(M) + (m_{12}-m_{22}+1-\delta(M))\chi_+(M),\\ &= (C_1(M)+\chi_+(M))(\bar{C}_1(M)+\chi_+(M)), \end{aligned}\tag{4.49}$$

$$C_2\left(M\left(\begin{smallmatrix}-1\,0\,0\\0\,0\\0\end{smallmatrix}\right)\right) = C_2(M), \tag{4.50}$$

$$\begin{aligned} \bar{F}\left(M\left(\begin{smallmatrix}-1\,0\,0\\0\,0\\0\end{smallmatrix}\right)\right) &= \bar{F}(M) + m_{12}-m_{22}-1+\delta(M)\\ &\quad - \chi_+(M)(m_{13}-m_{12}+\delta(M)), \end{aligned}\tag{4.51}$$

$$C_2\left(M\left(\begin{smallmatrix}-1\,0\,0\\0\,-1\\0\end{smallmatrix}\right)[-1]\right) = C_2(M) - (m_{12}-m_{22}-1+\delta(M))(1-\chi_+(M)), \tag{4.52}$$

$$\bar{F}(M)(1-\chi_+(M)) = -C_2(M)(1-\chi_+(M)), \tag{4.53}$$

(4.5) and (4.39).

- the case $(s,t)=(0,0)$.

We have

$$\begin{aligned}
c^{\lambda}_{[11;00;0]}(M) &= c^{\lambda}_{[1;00;0]}(M)\, c^{\lambda}_{[1;00;0]}\left(M\left(\begin{smallmatrix}-1&&0&&0\\&0&&0&\\&&0&&\end{smallmatrix}\right)\right)\\
&= (m_{13}-m_{12})(m_{13}-m_{22}+1)(m_{13}-m_{12}-1)(m_{13}-m_{22}),\\
c^{\lambda}_{[11;00;1]}(M) &= c^{\lambda}_{[1;00;0]}(M)\, c^{\lambda}_{[1;00;1]}\left(M\left(\begin{smallmatrix}-1&&0&&0\\&0&&0&\\&&0&&\end{smallmatrix}\right)\right) + c^{\lambda}_{[1;00;1]}(M)\, c^{\lambda}_{[1;00;0]}\left(M\left(\begin{smallmatrix}-1&&0&&0\\&0&&0&\\&&0&&\end{smallmatrix}\right)[-1]\right)\\
&= -(m_{13}-m_{12})(m_{13}-m_{22}+1)C_2\left(M\left(\begin{smallmatrix}-1&&0&&0\\&0&&0&\\&&0&&\end{smallmatrix}\right)\right)\\
&\quad - C_2(M)(m_{13}-m_{12})(m_{13}-m_{22}-1)\\
&= -2(m_{13}-m_{12})(m_{13}-m_{22})C_2(M),\\
c^{\lambda}_{[11;00;2]}(M) &= +c^{\lambda}_{[1;00;1]}(M)\, c^{\lambda}_{[1;00;1]}\left(M\left(\begin{smallmatrix}-1&&0&&0\\&0&&0&\\&&0&&\end{smallmatrix}\right)[-1]\right)\\
&= C_2(M)C_2\left(M\left(\begin{smallmatrix}-1&&0&&0\\&0&&0&\\&&0&&\end{smallmatrix}\right)[-1]\right)\\
&= C_2(M)(C_1(M)-1)(\bar{C}_1(M)-1).
\end{aligned}$$

Here we use the relations (4.39) and (4.50).

Next, we compute the case of formula 2.

- the case $(s,t)=(2,2)$.

We have

$$\begin{aligned}
c^{\lambda}_{[22;22;0]}(M) =& c^{\lambda}_{[2;11;0]}(M)\, c^{\lambda}_{[2;11;0]}\left(M\left(\begin{smallmatrix}0&&-1&&0\\&0&&-1&\\&&-1&&\end{smallmatrix}\right)\right) = (m_{22}-m_{33})(m_{22}-m_{33}-1),\\
c^{\lambda}_{[22;22;1]}(M) =& c^{\lambda}_{[2;11;0]}(M)\, c^{\lambda}_{[2;11;1]}\left(M\left(\begin{smallmatrix}0&&-1&&0\\&0&&-1&\\&&-1&&\end{smallmatrix}\right)\right) + c^{\lambda}_{[2;11;1]}(M)\, c^{\lambda}_{[2;11;0]}\left(M\left(\begin{smallmatrix}0&&-1&&0\\&0&&-1&\\&&-1&&\end{smallmatrix}\right)[-1]\right)\\
=& -(m_{22}-m_{33})\bar{D}\left(M\left(\begin{smallmatrix}0&&-1&&0\\&0&&-1&\\&&-1&&\end{smallmatrix}\right)\right)\chi_{-}\left(M\left(\begin{smallmatrix}0&&-1&&0\\&0&&-1&\\&&-1&&\end{smallmatrix}\right)\right) - \bar{D}(M)\chi_{-}(M)(m_{22}-m_{33})\\
=& -(m_{22}-m_{33})\{\bar{D}(M)\chi_{-}(M) + (\bar{D}(M)+2)\chi_{-}^{(1)}(M)\},\\
c^{\lambda}_{[22;22;2]}(M) =& c^{\lambda}_{[2;11;1]}(M)\, c^{\lambda}_{[2;11;1]}\left(M\left(\begin{smallmatrix}0&&-1&&0\\&0&&-1&\\&&-1&&\end{smallmatrix}\right)[-1]\right)\\
=& \bar{D}(M)\chi_{-}(M)\bar{D}\left(M\left(\begin{smallmatrix}0&&-1&&0\\&0&&-1&\\&&-1&&\end{smallmatrix}\right)[-1]\right)\chi_{-}\left(M\left(\begin{smallmatrix}0&&-1&&0\\&0&&-1&\\&&-1&&\end{smallmatrix}\right)[-1]\right)\\
=& \bar{D}(M)(\bar{D}(M)+1)\chi_{-}^{(1)}(M).
\end{aligned}$$

Here we use the relations

$$\bar{D}\left(M\left(\begin{smallmatrix}0&&-1&&0\\&0&&-1&\\&&-1&&\end{smallmatrix}\right)\right) = \bar{D}(M)+2, \tag{4.54}$$

(4.6), (4.12) and (4.62).

- the case $(s,t)=(1,2)$.

We have

$$\begin{aligned}
c^{\lambda}_{[22;12;0]}(M) =& c^{\lambda}_{[2;11;0]}(M)\, c^{\lambda}_{[2;01;0]}\left(M\left(\begin{smallmatrix}0&&-1&&0\\&0&&-1&\\&&-1&&\end{smallmatrix}\right)\right) + c^{\lambda}_{[2;01;0]}(M)\, c^{\lambda}_{[2;11;0]}\left(M\left(\begin{smallmatrix}0&&-1&&0\\&0&&-1&\\&&0&&\end{smallmatrix}\right)\right)\\
=& -(m_{22}-m_{33})(m_{22}-m_{33}-1) - (m_{22}-m_{33}))(m_{22}-m_{33}-1)\\
=& -2(m_{22}-m_{33})(m_{22}-m_{33}-1),\\
c^{\lambda}_{[22;12;1]}(M) =& c^{\lambda}_{[2;11;0]}(M)\, c^{\lambda}_{[2;01;1]}\left(M\left(\begin{smallmatrix}0&&-1&&0\\&0&&-1&\\&&-1&&\end{smallmatrix}\right)\right) + c^{\lambda}_{[2;01;1]}(M)\, c^{\lambda}_{[2;11;0]}\left(M\left(\begin{smallmatrix}0&&-1&&0\\&0&&-1&\\&&0&&\end{smallmatrix}\right)[-1]\right)\\
&+ c^{\lambda}_{[2;01;0]}(M)\, c^{\lambda}_{[2;11;1]}\left(M\left(\begin{smallmatrix}0&&-1&&0\\&0&&-1&\\&&0&&\end{smallmatrix}\right)\right) + c^{\lambda}_{[2;11;1]}(M)\, c^{\lambda}_{[2;01;0]}\left(M\left(\begin{smallmatrix}0&&-1&&0\\&0&&-1&\\&&-1&&\end{smallmatrix}\right)[-1]\right)
\end{aligned}$$

$$\begin{aligned}
&=(m_{22}-m_{33})\bar{C}_1\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&-1\end{smallmatrix}\right)\right)+\bar{C}_1(M)(m_{22}-m_{33})\\
&\quad+(m_{22}-m_{33})\bar{D}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&0\end{smallmatrix}\right)\right)\chi_-\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&0\end{smallmatrix}\right)\right)+\bar{D}(M)\chi_-(M)(m_{22}-m_{33})\\
&=2(m_{22}-m_{33})\{\bar{C}_1(M)+(\bar{D}(M)+1)\chi_-(M)\},\\
c^{\lambda}_{[22;12;2]}(M)&=c^{\lambda}_{[2;11;1]}(M)\,c^{\lambda}_{[2;01;1]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&-1\end{smallmatrix}\right)[-1]\right)+c^{\lambda}_{[2;01;1]}(M)\,c^{\lambda}_{[2;11;1]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&0\end{smallmatrix}\right)[-1]\right)\\
&=-\bar{D}(M)\chi_-(M)\bar{C}_1\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&-1\end{smallmatrix}\right)[-1]\right)\\
&\quad-\bar{C}_1(M)\bar{D}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&0\end{smallmatrix}\right)[-1]\right)\chi_-\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&0\end{smallmatrix}\right)[-1]\right)\\
&=-2\bar{C}_1(M)\bar{D}(M)\chi_-(M).
\end{aligned}$$

Here we use the relations

$$\bar{D}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&0\end{smallmatrix}\right)\right)=\bar{D}(M)+1, \tag{4.55}$$

$$\bar{C_1}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&-1\end{smallmatrix}\right)\right)=\bar{C_1}(M)+chi_-(M), \tag{4.56}$$

$$\bar{C_1}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&-1\end{smallmatrix}\right)[-1]\right)\chi_-(M)=\bar{C_1}(M)\chi_-(M), \tag{4.57}$$

(4.6) and (4.62).

- the case $(s,t)=(1,1)$.

  We have

$$\begin{aligned}
c^{\lambda}_{[22;11;0]}(M)&=c^{\lambda}_{[2;11;0]}(M)\,c^{\lambda}_{[2;00;0]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&-1\end{smallmatrix}\right)\right)+c^{\lambda}_{[2;00;0]}(M)\,c^{\lambda}_{[2;11;0]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)\right)\\
&=-(m_{22}-m_{33})(m_{23}-m_{22})-(m_{23}-m_{22})(m_{22}-m_{33})\\
&=-2(m_{22}-m_{33})(m_{23}-m_{22}),\\
c^{\lambda}_{[22;11;1]}(M)&=c^{\lambda}_{[2;11;0]}(M)\,c^{\lambda}_{[2;00;1]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&-1\end{smallmatrix}\right)\right)+c^{\lambda}_{[2;00;1]}(M)\,c^{\lambda}_{[2;11;0]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)[-1]\right)\\
&\quad+c^{\lambda}_{[2;11;1]}(M)\,c^{\lambda}_{[2;00;0]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&-1\end{smallmatrix}\right)[-1]\right)+c^{\lambda}_{[2;00;0]}(M)\,c^{\lambda}_{[2;11;1]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)\right)\\
&=-(m_{22}-m_{33})\bar{C}_1\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&-1\end{smallmatrix}\right)\right)\chi_-\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&-1\end{smallmatrix}\right)\right)\\
&\quad-\bar{C}_1(M)\chi_-(M)(m_{22}-m_{33}+1)+\bar{D}(M)\chi_-(M)(m_{23}-m_{22}-1)\\
&\quad+(m_{23}-m_{22})\bar{D}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)\right)\chi_-\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)\right)\\
&=-(m_{22}-m_{33})(m_{12}-m_{11}+1)\chi_-^{(1)}(M)-(m_{12}-m_{11})\chi_-(M)(m_{22}-m_{33}+1)\\
&\quad+(-m_{22}+m_{33}+\delta(M))\chi_-(M)(m_{23}-m_{22}-1)\\
&\quad+(m_{23}-m_{22})(-m_{22}+m_{33}+\delta(M)+1)\chi_-^{(1)}(M)\\
&=-2(m_{22}-m_{33})(m_{12}-m_{11}+1)\chi_-^{(1)}(M)\\
&\quad+2(-m_{22}+m_{33}+\delta(M))\chi_-(M)(m_{23}-m_{22}-1)\\
&\quad+(m_{23}-m_{33})(\delta(M)+1)(\chi_-^{(1)}(M)-\chi_-(M))\\
&=2\{\bar{D}(M)(m_{23}-m_{22}-1)\chi_-(M)-(m_{22}-m_{33})(\bar{C}_1(M)+1)\chi_-^{(1)}(M)\},\\
c^{\lambda}_{[22;11;2]}(M)&=c^{\lambda}_{[2;11;1]}(M)\,c^{\lambda}_{[2;00;1]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&-1\end{smallmatrix}\right)[-1]\right)+c^{\lambda}_{[2;00;1]}(M)\,c^{\lambda}_{[2;11;1]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)[-1]\right)\\
&=\bar{D}(M)\chi_-(M)\bar{C}_1\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&-1\end{smallmatrix}\right)[-1]\right)\chi_-\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&-1\end{smallmatrix}\right)[-1]\right)
\end{aligned}$$

$$\begin{aligned}
&+ \bar{C}_1(M)\chi_-(M)\bar{D}\left(M\left(\begin{smallmatrix} 0 & -1 & 0 \\ & 0 & 0 \\ & & 0\end{smallmatrix}\right)[-1]\right)\chi_-\left(M\left(\begin{smallmatrix} 0 & -1 & 0 \\ & 0 & 0 \\ & & 0\end{smallmatrix}\right)[-1]\right)\\
=&\bar{D}(M)(m_{12}-m_{11})\chi_-^{(1)}(M) + \bar{C}_1(M)\bar{D}(M)\chi_-^{(1)}(M)\\
=&2\bar{C}_1(M)\bar{D}(M)\chi_-^{(1)}(M).
\end{aligned}$$

Here we use the relations

$$\bar{D}\left(M\left(\begin{smallmatrix} 0 & -1 & 0 \\ & 0 & 0 \\ & & 0\end{smallmatrix}\right)\right) = \bar{D}(M)+1, \tag{4.58}$$

(4.4), (4.6), (4.12), (4.16), (4.56) and (4.62).

- the case $(s,t)=(0,2)$.
  We have

$$\begin{aligned}
c^{\lambda}_{[22;02;0]}(M) =&c^{\lambda}_{[2;01;0]}(M)\,c^{\lambda}_{[2;01;0]}\left(M\left(\begin{smallmatrix} 0 & -1 & 0 \\ & 0 & -1 \\ & & 0\end{smallmatrix}\right)\right)\\
=&(m_{22}-m_{33})(m_{22}-m_{33}-1)\\
c^{\lambda}_{[22;02;1]}(M) =&c^{\lambda}_{[2;01;0]}(M)\,c^{\lambda}_{[2;01;1]}\left(M\left(\begin{smallmatrix} 0 & -1 & 0 \\ & 0 & -1 \\ & & 0\end{smallmatrix}\right)\right) + c^{\lambda}_{[2;01;1]}(M)\,c^{\lambda}_{[2;01;0]}\left(M\left(\begin{smallmatrix} 0 & -1 & 0 \\ & 0 & -1 \\ & & 0\end{smallmatrix}\right)[-1]\right)\\
=&-(m_{22}-m_{33})\bar{C}_1\left(M\left(\begin{smallmatrix} 0 & -1 & 0 \\ & 0 & -1 \\ & & 0\end{smallmatrix}\right)\right) - \bar{C}_1(M)(m_{22}-m_{33})\\
=&-2(m_{22}-m_{33})\bar{C}_1(M),\\
c^{\lambda}_{[22;02;2]}(M) =&c^{\lambda}_{[2;01;1]}(M)\,c^{\lambda}_{[2;01;1]}\left(M\left(\begin{smallmatrix} 0 & -1 & 0 \\ & 0 & -1 \\ & & 0\end{smallmatrix}\right)[-1]\right)\\
=&\bar{C}_1(M)\bar{C}_1\left(M\left(\begin{smallmatrix} 0 & -1 & 0 \\ & 0 & -1 \\ & & 0\end{smallmatrix}\right)[-1]\right)\\
=&\bar{C}_1(M)(\bar{C}_1(M)-1).
\end{aligned}$$

Here we use the relations

$$\bar{C}_1\left(M\left(\begin{smallmatrix} 0 & -1 & 0 \\ & 0 & -1 \\ & & 0\end{smallmatrix}\right)\right) = \bar{C}_1(M), \tag{4.59}$$

and (4.39).

- the case $(s,t)=(0,1)$.
  We have

$$\begin{aligned}
c^{\lambda}_{[22;01;0]}(M) =&c^{\lambda}_{[2;01;0]}(M)\,c^{\lambda}_{[2;00;0]}\left(M\left(\begin{smallmatrix} 0 & -1 & 0 \\ & 0 & -1 \\ & & 0\end{smallmatrix}\right)\right) + c^{\lambda}_{[2;00;0]}(M)\,c^{\lambda}_{[2;01;0]}\left(M\left(\begin{smallmatrix} 0 & -1 & 0 \\ & 0 & 0 \\ & & 0\end{smallmatrix}\right)\right)\\
=&(m_{22}-m_{33})(m_{23}-m_{22}) + (m_{23}-m_{22})(m_{22}-m_{33})\\
=&2(m_{22}-m_{33})(m_{23}-m_{22}),\\
c^{\lambda}_{[22;01;1]}(M) =&c^{\lambda}_{[2;01;0]}(M)\,c^{\lambda}_{[2;00;1]}\left(M\left(\begin{smallmatrix} 0 & -1 & 0 \\ & 0 & -1 \\ & & 0\end{smallmatrix}\right)\right) + c^{\lambda}_{[2;00;1]}(M)\,c^{\lambda}_{[2;01;0]}\left(M\left(\begin{smallmatrix} 0 & -1 & 0 \\ & 0 & 0 \\ & & 0\end{smallmatrix}\right)[-1]\right)\\
&+ c^{\lambda}_{[2;01;1]}(M)\,c^{\lambda}_{[2;00;0]}\left(M\left(\begin{smallmatrix} 0 & -1 & 0 \\ & 0 & -1 \\ & & 0\end{smallmatrix}\right)[-1]\right) + c^{\lambda}_{[2;00;0]}(M)\,c^{\lambda}_{[2;01;1]}\left(M\left(\begin{smallmatrix} 0 & -1 & 0 \\ & 0 & 0 \\ & & 0\end{smallmatrix}\right)\right)\\
=&(m_{22}-m_{33})\bar{C}_1\left(M\left(\begin{smallmatrix} 0 & -1 & 0 \\ & 0 & -1 \\ & & 0\end{smallmatrix}\right)\right)\chi_-\left(M\left(\begin{smallmatrix} 0 & -1 & 0 \\ & 0 & -1 \\ & & 0\end{smallmatrix}\right)\right) + \bar{C}_1(M)\chi_-(M)(m_{22}-m_{33}+1)\\
&- \bar{C}_1(M)(m_{23}-m_{22}-1) - (m_{23}-m_{22})\bar{C}_1\left(M\left(\begin{smallmatrix} 0 & -1 & 0 \\ & 0 & 0 \\ & & 0\end{smallmatrix}\right)\right)\\
=&(m_{22}-m_{33})\bar{C}_1(M)\chi_-(M) + \bar{C}_1(M)\chi_-(M)(m_{22}-m_{33}+1)\\
&- \bar{C}_1(M)(m_{23}-m_{22}-1) - (m_{23}-m_{22})(\bar{C}_1(M)-\chi_+^{(-1)}(M))\\
=&2(m_{22}-m_{33})\bar{C}_1(M)\chi_-(M)\\
&- 2(m_{23}-m_{22})\bar{C}_1(M) + \bar{C}_1(M)(1+\chi_-(M)+\chi_+^{(-1)}(M))
\end{aligned}$$

$$
\begin{aligned}
&=2\bar{C}_1(M)\{(m_{22}-m_{33})\chi_-(M)-(m_{23}-m_{22}-1)\},\\
c^{\lambda}_{[22;01;2]}(M) &=c^{\lambda}_{[2;00;1]}(M)\,c^{\lambda}_{[2;01;1]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)[-1]\right)+c^{\lambda}_{[2;01;1]}(M)\,c^{\lambda}_{[2;00;1]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&0\end{smallmatrix}\right)[-1]\right)\\
&=-\bar{C}_1(M)\chi_-(M)\bar{C}_1\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)[-1]\right)\\
&\quad-\bar{C}_1(M)\bar{C}_1\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&0\end{smallmatrix}\right)[-1]\right)\chi_-\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&0\end{smallmatrix}\right)[-1]\right)\\
&=-\bar{C}_1(M)\chi_-(M)(\bar{C}_1(M)-\chi_+^{(-1)}(M)-1)-\bar{C}_1(M)(\bar{C}_1(M)-1)\chi_-(M)\\
&=-2\bar{C}_1(M)(\bar{C}_1(M)-1)\chi_-(M).
\end{aligned}
$$

Here we use the equations

$$
\bar{C}_1\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)\right)=\bar{C}_1(M)-\chi_+^{(-1)}(M), \tag{4.60}
$$

(4.6), (4.10), (4.9), (4.15), (4.59) and (4.39).

- the case $(s,t)=(0,0)$.
  We have

$$
\begin{aligned}
c^{\lambda}_{[22;00;0]}(M) &=c^{\lambda}_{[2;00;0]}(M)\,c^{\lambda}_{[2;00;0]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)\right)\\
&=(m_{23}-m_{22}))(m_{23}-m_{22}-1)\\
c^{\lambda}_{[22;00;1]}(M) &=c^{\lambda}_{[2;00;0]}(M)\,c^{\lambda}_{[2;00;1]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)\right)+c^{\lambda}_{[2;00;1]}(M)\,c^{\lambda}_{[2;00;0]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)[-1]\right)\\
&=(m_{23}-m_{22})\bar{C}_1\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)\right)\chi_-\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)\right)+\bar{C}_1(M)\chi_-(M)(m_{23}-m_{22}-2)\\
&=\bar{C}_1(M)\{(m_{23}-m_{22}-2)\chi_-(M)+(m_{23}-m_{22})\chi_-^{(1)}(M)\},\\
c^{\lambda}_{[22;00;2]}(M) &=c^{\lambda}_{[2;00;1]}(M)\,c^{\lambda}_{[2;00;1]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)[-1]\right)\\
&=\bar{C}_1(M)\chi_-(M)\bar{C}_1\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)[-1]\right)\chi_-\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)[-1]\right)\\
&=\bar{C}_1(M)(\bar{C}_1(M)-1)\chi_-^{(1)}(M).
\end{aligned}
$$

Here we use the relations (4.6), (4.10), (4.12), (4.39) and (4.60).

Next, we compute the case of formula 3.

- the case $(s,t)=(2,2)$.
  We have

$$
c^{\lambda}_{[33;22;0]}(M)=c^{\lambda}_{[3;11;0]}(M)\,c^{\lambda}_{[3;11;0]}\left(M\left(\begin{smallmatrix}0&0&-1\\&0&-1\\&&-1\end{smallmatrix}\right)\right)=1.
$$

- the case $(s,t)=(1,2)$.
  We have

$$
\begin{aligned}
c^{\lambda}_{[33;12;0]}(M) &=c^{\lambda}_{[3;11;0]}(M)\,c^{\lambda}_{[3;01;0]}\left(M\left(\begin{smallmatrix}0&0&-1\\&0&-1\\&&-1\end{smallmatrix}\right)\right)+c^{\lambda}_{[3;01;0]}(M)\,c^{\lambda}_{[3;11;0]}\left(M\left(\begin{smallmatrix}0&0&-1\\&0&-1\\&&0\end{smallmatrix}\right)\right)\\
&=-1-1=-2,\\
c^{\lambda}_{[33;12;1]}(M) &=c^{\lambda}_{[3;11;0]}(M)\,c^{\lambda}_{[3;01;1]}\left(M\left(\begin{smallmatrix}0&0&-1\\&0&-1\\&&-1\end{smallmatrix}\right)\right)+c^{\lambda}_{[3;01;1]}(M)\,c^{\lambda}_{[3;11;0]}\left(M\left(\begin{smallmatrix}0&0&-1\\&0&-1\\&&0\end{smallmatrix}\right)[-1]\right)\\
&=-\chi_+\left(M\left(\begin{smallmatrix}0&0&-1\\&0&-1\\&&-1\end{smallmatrix}\right)\right)-\chi_+(M)=-2\chi_+(M).
\end{aligned}
$$

Here we use the relation (4.5).

- the case $(s,t) = (1,1)$.
  We have

$$\begin{aligned} c^{\lambda}_{[33;11;0]}(M) =& c^{\lambda}_{[3;11;0]}(M)\, c^{\lambda}_{[3;00;0]}\left(M\left(\begin{smallmatrix}0&0&-1\\&0&-1\\&&-1\end{smallmatrix}\right)\right) + c^{\lambda}_{[3;00;0]}(M)\, c^{\lambda}_{[3;11;0]}\left(M\left(\begin{smallmatrix}0&0&-1\\&0&0\\&&0\end{smallmatrix}\right)\right) \\ =& 1+1=2. \end{aligned}$$

- the case $(s,t) = (0,2)$.
  We have

$$\begin{aligned} c^{\lambda}_{[33;02;0]}(M) =& c^{\lambda}_{[3;01;0]}(M)\, c^{\lambda}_{[3;01;0]}\left(M\left(\begin{smallmatrix}0&0&-1\\&0&-1\\&&0\end{smallmatrix}\right)\right) = 1. \\ c^{\lambda}_{[33;02;1]}(M) =& c^{\lambda}_{[3;01;0]}(M)\, c^{\lambda}_{[3;01;1]}\left(M\left(\begin{smallmatrix}0&0&-1\\&0&-1\\&&0\end{smallmatrix}\right)\right) + c^{\lambda}_{[3;01;1]}(M)\, c^{\lambda}_{[3;01;0]}\left(M\left(\begin{smallmatrix}0&0&-1\\&0&-1\\&&0\end{smallmatrix}\right)[-1]\right) \\ =& \chi_{+}\left(M\left(\begin{smallmatrix}0&0&-1\\&0&-1\\&&0\end{smallmatrix}\right)\right) + \chi_{+}(M) = \chi_{+}^{(1)}(M) + \chi_{+}(M) \\ c^{\lambda}_{[33;02;2]}(M) =& c^{\lambda}_{[3;01;1]}(M)\, c^{\lambda}_{[3;01;1]}\left(M\left(\begin{smallmatrix}0&0&-1\\&0&-1\\&&0\end{smallmatrix}\right)[-1]\right) = \chi_{+}(M)\chi_{+}\left(M\left(\begin{smallmatrix}0&0&-1\\&0&-1\\&&0\end{smallmatrix}\right)[-1]\right) \\ =& \chi_{+}^{(1)}(M). \end{aligned}$$

Here we use the relation (4.5).

- the case $(s,t) = (0,1)$.
  We have

$$\begin{aligned} c^{\lambda}_{[33;01;0]}(M) =& c^{\lambda}_{[3;01;0]}(M)\, c^{\lambda}_{[3;00;0]}\left(M\left(\begin{smallmatrix}0&0&-1\\&0&-1\\&&0\end{smallmatrix}\right)\right) + c^{\lambda}_{[3;00;0]}(M)\, c^{\lambda}_{[3;01;0]}\left(M\left(\begin{smallmatrix}0&0&-1\\&0&0\\&&0\end{smallmatrix}\right)\right) \\ =& -1-1=-2, \\ c^{\lambda}_{[33;01;1]}(M) =& c^{\lambda}_{[3;01;1]}(M)\, c^{\lambda}_{[3;00;0]}\left(M\left(\begin{smallmatrix}0&0&-1\\&0&-1\\&&0\end{smallmatrix}\right)[-1]\right) + c^{\lambda}_{[3;00;0]}(M)\, c^{\lambda}_{[3;01;1]}\left(M\left(\begin{smallmatrix}0&0&-1\\&0&0\\&&0\end{smallmatrix}\right)\right) \\ =& -\chi_{+}(M) - \chi_{+}\left(M\left(\begin{smallmatrix}0&0&-1\\&0&0\\&&0\end{smallmatrix}\right)\right) = -2\chi_{+}(M). \end{aligned}$$

Here we use the relation (4.5).

- the case $(s,t) = (0,0)$.
  We have

$$c^{\lambda}_{[33;00;0]}(M) = c^{\lambda}_{[3;00;0]}(M)\, c^{\lambda}_{[3;00;0]}\left(M\left(\begin{smallmatrix}0&0&-1\\&0&0\\&&0\end{smallmatrix}\right)\right) = 1.$$

Next, we compute the case of formula 4.

- the case $(s,t) = (2,2)$.
  We have

$$\begin{aligned} c^{\lambda}_{[12;22;0]}(M) =& c^{\lambda}_{[1;11;0]}(M)\, c^{\lambda}_{[2;11;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&-1\end{smallmatrix}\right)\right) \\ =& (m_{13}-m_{12})(m_{22}-m_{33})(m_{22}-m_{33}-1), \\ c^{\lambda}_{[12;22;1]}(M) =& c^{\lambda}_{[1;11;0]}(M)\, c^{\lambda}_{[2;11;1]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&-1\end{smallmatrix}\right)\right) + c^{\lambda}_{[1;11;1]}(M)\, c^{\lambda}_{[2;11;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&-1\end{smallmatrix}\right)[-1]\right) \\ =& -(m_{13}-m_{12})(m_{22}-m_{33})\bar{D}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&-1\end{smallmatrix}\right)\right)\chi_{-}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&-1\end{smallmatrix}\right)\right) \\ & -\bar{E}(M)(m_{22}-m_{33}) \end{aligned}$$

$$
\begin{aligned}
&=-(m_{22}-m_{33})\{\bar{E}(M)+(m_{13}-m_{12})(\bar{D}(M)+1)\chi_-(M)\},\\
c^{\lambda}_{[12;22;2]}(M)=&c^{\lambda}_{[1;11;1]}(M)\,c^{\lambda}_{[2;11;1]}\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1\\-1\end{smallmatrix}\right)[-1]\right)\\
=&\bar{E}(M)\bar{D}\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1\\-1\end{smallmatrix}\right)[-1]\right)\chi_-\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1\\-1\end{smallmatrix}\right)[-1]\right)\\
=&\bar{E}(M)\bar{D}(M)\chi_-(M).
\end{aligned}
$$

Here we use the relations:

$$
\bar{D}\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1\\-1\end{smallmatrix}\right)\right)=\bar{D}(M)+1, \tag{4.61}
$$

$$
\bar{D}\left(M[-1]\right)=\bar{D}(M)-1, \tag{4.62}
$$

and (4.6).

- the case $(s,t)=(1,2)$.

We have

$$
\begin{aligned}
c^{\lambda}_{[12;12;0]}(M)=&c^{\lambda}_{[1;11;0]}(M)\,c^{\lambda}_{[2;01;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1\\-1\end{smallmatrix}\right)\right)+c^{\lambda}_{[1;01;0]}(M)\,c^{\lambda}_{[2;11;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1\\0\end{smallmatrix}\right)\right)\\
=&-(m_{13}-m_{12})(m_{22}-m_{33})(m_{22}-m_{33}-1)\\
&-(m_{13}-m_{12})(m_{22}-m_{33})(m_{22}-m_{33}-1)\\
=&-2(m_{13}-m_{12})(m_{22}-m_{33})(m_{22}-m_{33}-1),\\
c^{\lambda}_{[12;12;1]}(M)=&c^{\lambda}_{[1;11;0]}(M)\,c^{\lambda}_{[2;01;1]}\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1\\-1\end{smallmatrix}\right)\right)+c^{\lambda}_{[1;01;1]}(M)\,c^{\lambda}_{[2;11;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1\\0\end{smallmatrix}\right)[-1]\right)\\
&+c^{\lambda}_{[1;01;0]}(M)\,c^{\lambda}_{[2;11;1]}\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1\\0\end{smallmatrix}\right)\right)+c^{\lambda}_{[1;11;1]}(M)\,c^{\lambda}_{[2;01;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1\\-1\end{smallmatrix}\right)[-1]\right)\\
=&(m_{13}-m_{12})(m_{22}-m_{33})\bar{C}_1\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1\\-1\end{smallmatrix}\right)\right)\\
&+\bar{F}(M)(m_{22}-m_{33})\\
&+(m_{13}-m_{12})(m_{22}-m_{33})\bar{D}\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1\\0\end{smallmatrix}\right)\right)\chi_-\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1\\0\end{smallmatrix}\right)\right)\\
&+\bar{E}(M)(m_{22}-m_{33})\\
=&(m_{22}-m_{33})\big\{\bar{E}(M)+\bar{F}(M)\\
&+(m_{13}-m_{12})\{\bar{C}_1(M)+1+\bar{D}(M)(1-\chi_+(M))\}\big\},\\
c^{\lambda}_{[12;12;2]}(M)=&c^{\lambda}_{[1;11;1]}(M)\,c^{\lambda}_{[2;01;1]}\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1\\-1\end{smallmatrix}\right)[-1]\right)+c^{\lambda}_{[1;01;1]}(M)\,c^{\lambda}_{[2;11;1]}\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1\\0\end{smallmatrix}\right)[-1]\right)\\
&+c^{\lambda}_{[1;01;2]}(M)\,c^{\lambda}_{[2;11;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1\\0\end{smallmatrix}\right)[-2]\right)\\
=&-\bar{E}(M)\bar{C}_1\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1\\-1\end{smallmatrix}\right)[-1]\right)\\
&-\bar{F}(M)\bar{D}\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1\\0\end{smallmatrix}\right)[-1]\right)\chi_-\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1\\0\end{smallmatrix}\right)[-1]\right)\\
&-C_2(M)\chi_+(M)(m_{22}-m_{33}+1)\\
=&-\bar{E}(M)\bar{C}_1(M)-C_2(M)(1-\bar{D}(M)+\delta(M)\chi_+(M)),\\
c^{\lambda}_{[12;12;3]}(M)=&c^{\lambda}_{[1;01;2]}(M)\,c^{\lambda}_{[2;11;1]}\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1\\0\end{smallmatrix}\right)[-2]\right)\\
=&C_2(M)\chi_+(M)\bar{D}\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1\\0\end{smallmatrix}\right)[-2]\right)\chi_-\left(M\left(\begin{smallmatrix}-1&0&0\\0&-1\\0\end{smallmatrix}\right)[-2]\right)\\
=&0.
\end{aligned}
$$

Here we use the relations:

$$\bar{D}\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & -1 \\ & & 0 \end{smallmatrix}\right)\right) = \bar{D}(M), \tag{4.63}$$

$$\bar{C}_1\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & -1 \\ & & -1 \end{smallmatrix}\right)\right) = \bar{C}_1(M) + 1, \tag{4.64}$$

(4.6), (4.9), (4.10), (4.62), (4.26) and (4.39).

- the case $(s, t) = (1, 1)$.
  We have

$$\begin{aligned}
c^{\lambda}_{[12;11;0]}(M) =& c^{\lambda}_{[1;11;0]}(M)\, c^{\lambda}_{[2;00;0]}\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & -1 \\ & & -1 \end{smallmatrix}\right)\right) + c^{\lambda}_{[1;00;0]}(M)\, c^{\lambda}_{[2;11;0]}\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & 0 \\ & & 0 \end{smallmatrix}\right)\right) \\
=& -(m_{13} - m_{12})(m_{22} - m_{33})(m_{23} - m_{22} + 1) \\
& -(m_{13} - m_{12})(m_{13} - m_{22} + 1)(m_{22} - m_{33}) \\
=& (m_{13} - m_{12})(m_{22} - m_{33})(2m_{22} - m_{13} - m_{23} - 2), \\
c^{\lambda}_{[12;11;1]}(M) =& c^{\lambda}_{[1;11;0]}(M)\, c^{\lambda}_{[2;00;1]}\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & -1 \\ & & -1 \end{smallmatrix}\right)\right) + c^{\lambda}_{[1;00;1]}(M)\, c^{\lambda}_{[2;11;0]}\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & 0 \\ & & 0 \end{smallmatrix}\right)[-1]\right) \\
& + c^{\lambda}_{[1;11;1]}(M)\, c^{\lambda}_{[2;00;0]}\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & -1 \\ & & -1 \end{smallmatrix}\right)[-1]\right) + c^{\lambda}_{[1;00;0]}(M)\, c^{\lambda}_{[2;11;1]}\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & 0 \\ & & 0 \end{smallmatrix}\right)\right) \\
=& -(m_{13} - m_{12})(m_{22} - m_{33})\bar{C}_1\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & -1 \\ & & -1 \end{smallmatrix}\right)\right)\chi_-\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & -1 \\ & & -1 \end{smallmatrix}\right)\right) \\
& + C_2(M)(m_{22} - m_{33} + 1) + \bar{E}(M)(m_{23} - m_{22}) \\
& + (m_{13} - m_{12})(m_{13} - m_{22} + 1)\bar{D}\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & 0 \\ & & 0 \end{smallmatrix}\right)\right)\chi_-\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & 0 \\ & & 0 \end{smallmatrix}\right)\right) \\
=& \bar{E}(M)(m_{23} - m_{22}) + C_2(M)(m_{22} - m_{33} + 1) \\
& + (m_{13} - m_{12})\chi_-(M)\{(m_{13} - m_{22} + 1)\bar{D}(M) - (m_{22} - m_{33})(\bar{C}_1(M) + 1)\}, \\
c^{\lambda}_{[12;11;2]}(M) =& c^{\lambda}_{[1;11;1]}(M)\, c^{\lambda}_{[2;00;1]}\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & -1 \\ & & -1 \end{smallmatrix}\right)[-1]\right) + c^{\lambda}_{[1;00;1]}(M)\, c^{\lambda}_{[2;11;1]}\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & 0 \\ & & 0 \end{smallmatrix}\right)[-1]\right) \\
=& \bar{E}(M)\bar{C}_1\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & -1 \\ & & -1 \end{smallmatrix}\right)[-1]\right)\chi_-\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & -1 \\ & & -1 \end{smallmatrix}\right)[-1]\right) \\
& - C_2(M)\bar{D}\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & 0 \\ & & 0 \end{smallmatrix}\right)[-1]\right)\chi_-\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & 0 \\ & & 0 \end{smallmatrix}\right)[-1]\right) \\
=& C_2(M)(m_{13} - m_{33} + 2 - \bar{C}_1(M) - \bar{D}(M)).
\end{aligned}$$

Here we use the relations:

$$\bar{D}\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & 0 \\ & & 0 \end{smallmatrix}\right)\right) = \bar{D}(M), \tag{4.65}$$

(4.6), (4.62), (4.64) and (4.39).

- the case $(s, t) = (0, 2)$.
  We have

$$\begin{aligned}
c^{\lambda}_{[12;02;0]}(M) =& c^{\lambda}_{[1;01;0]}(M)\, c^{\lambda}_{[2;01;0]}\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & -1 \\ & & 0 \end{smallmatrix}\right)\right) \\
=& (m_{13} - m_{12})(m_{22} - m_{33})(m_{22} - m_{33} - 1), \\
c^{\lambda}_{[12;02;1]}(M) =& c^{\lambda}_{[1;01;0]}(M)\, c^{\lambda}_{[2;01;1]}\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & -1 \\ & & 0 \end{smallmatrix}\right)\right) + c^{\lambda}_{[1;01;1]}(M)\, c^{\lambda}_{[2;01;0]}\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & -1 \\ & & 0 \end{smallmatrix}\right)[-1]\right) \\
=& -(m_{13} - m_{12})(m_{22} - m_{33})\bar{C}_1\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & -1 \\ & & 0 \end{smallmatrix}\right)\right) - \bar{F}(M)(m_{22} - m_{33}) \\
=& -(m_{22} - m_{33})\{(m_{13} - m_{12})(\bar{C}_1(M) + \chi_+(M)) + \bar{F}(M)\},
\end{aligned}$$

$$\begin{aligned}
c^{\lambda}_{[12;02;2]}(M) =& c^{\lambda}_{[1;01;2]}(M)\, c^{\lambda}_{[2;01;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&0&\end{smallmatrix}\right)[-2]\right) + c^{\lambda}_{[1;01;1]}(M)\, c^{\lambda}_{[2;01;1]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&0&\end{smallmatrix}\right)[-1]\right)\\
=& C_2(M)\chi_+(M)(m_{22}-m_{33}+1) + \bar{F}(M)\bar{C}_1\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&0&\end{smallmatrix}\right)[-1]\right)\\
=& (\bar{C}_1(M)+\chi_+(M)-1)\bar{F}(M) + (m_{22}-m_{33}+1)C_2(M)\chi_+(M),\\
c^{\lambda}_{[12;02;3]}(M) =& c^{\lambda}_{[1;01;2]}(M)\, c^{\lambda}_{[2;01;1]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&0&\end{smallmatrix}\right)[-2]\right)\\
=& -C_2(M)\chi_+(M)\bar{C}_1\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&0&\end{smallmatrix}\right)[-2]\right)\\
=& -C_2(M)(\bar{C}_1(M)-1)\chi_+(M).
\end{aligned}$$

Here we use the relations:

$$\bar{C}_1\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&0&\end{smallmatrix}\right)\right) = \bar{C}_1(M)+\chi_+(M), \tag{4.66}$$

and (4.39).

- the case $(s,t)=(0,1)$.
  We have

$$\begin{aligned}
c^{\lambda}_{[12;01;0]}(M) =& c^{\lambda}_{[1;01;0]}(M)\, c^{\lambda}_{[2;00;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&0&\end{smallmatrix}\right)\right) + c^{\lambda}_{[1;00;0]}(M)\, c^{\lambda}_{[2;01;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&0\\&0&\end{smallmatrix}\right)\right)\\
=& (m_{13}-m_{12})(m_{22}-m_{33})(m_{23}-m_{22}+1)\\
&+ (m_{13}-m_{12})(m_{13}-m_{22}+1)(m_{22}-m_{33})\\
=& -(m_{13}-m_{12})(m_{22}-m_{33})(2m_{22}-m_{13}-m_{23}-2),\\
c^{\lambda}_{[12;01;1]}(M) =& c^{\lambda}_{[1;01;0]}(M)\, c^{\lambda}_{[2;00;1]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&0&\end{smallmatrix}\right)\right) + c^{\lambda}_{[1;00;1]}(M)\, c^{\lambda}_{[2;01;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&0\\&0&\end{smallmatrix}\right)[-1]\right)\\
&+ c^{\lambda}_{[1;01;1]}(M)\, c^{\lambda}_{[2;00;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&0&\end{smallmatrix}\right)[-1]\right) + c^{\lambda}_{[1;00;0]}(M)\, c^{\lambda}_{[2;01;1]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&0\\&0&\end{smallmatrix}\right)\right)\\
=& (m_{13}-m_{12})(m_{22}-m_{33})\bar{C}_1\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&0&\end{smallmatrix}\right)\right)\chi_-\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&0&\end{smallmatrix}\right)\right)\\
&- C_2(M)(m_{22}-m_{33}+1) - \bar{F}(M)(m_{23}-m_{22})\\
&- (m_{13}-m_{12})(m_{13}-m_{22}+1)\bar{C}_1\left(M\left(\begin{smallmatrix}-1&0&0\\&0&0\\&0&\end{smallmatrix}\right)\right)\\
=& (m_{13}-m_{12})\bar{C}_1(M)\{(m_{22}-m_{33})(1-\chi_+(M)) - (m_{13}-m_{22}+1)\}\\
&- C_2(M)(m_{22}-m_{33}+1) - \bar{F}(M)(m_{23}-m_{22}),\\
c^{\lambda}_{[12;01;2]}(M) =& c^{\lambda}_{[1;00;1]}(M)\, c^{\lambda}_{[2;01;1]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&0\\&0&\end{smallmatrix}\right)[-1]\right) + c^{\lambda}_{[1;01;1]}(M)\, c^{\lambda}_{[2;00;1]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&0&\end{smallmatrix}\right)[-1]\right)\\
&+ c^{\lambda}_{[1;01;2]}(M)\, c^{\lambda}_{[2;00;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&0&\end{smallmatrix}\right)[-2]\right)\\
=& C_2(M)\bar{C}_1\left(M\left(\begin{smallmatrix}-1&0&0\\&0&0\\&0&\end{smallmatrix}\right)[-1]\right)\\
&- \bar{F}(M)\bar{C}_1\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&0&\end{smallmatrix}\right)[-1]\right)\chi_-\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&0&\end{smallmatrix}\right)[-1]\right)\\
&+ C_2(M)\chi_+(M)(m_{23}-m_{22}-1)\\
=& C_2(M)(\bar{C}_1(M)-1)(1+\chi_-^{(-1)}(M)+\chi_+(M))\\
=& 2C_2(M)(\bar{C}_1(M)-1),\\
c^{\lambda}_{[12;01;3]}(M) =& c^{\lambda}_{[1;01;2]}(M)\, c^{\lambda}_{[2;00;1]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&0&\end{smallmatrix}\right)[-2]\right)
\end{aligned}$$

$$\begin{aligned}
&=C_2(M)\chi_+(M)\bar{C}_1\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&0\end{smallmatrix}\right)[-2]\right)\chi_-\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&0\end{smallmatrix}\right)[-2]\right)\\
&=0.
\end{aligned}$$

Here we use the relations:

$$\bar{C}_1\left(M\left(\begin{smallmatrix}-1&0&0\\&0&0\\&&0\end{smallmatrix}\right)\right)=\bar{C}_1(M), \tag{4.67}$$

(4.6), (4.9), (4.10), (4.15), (4.26), (4.39) and (4.66).

- the case $(s,t)=(0,0)$.
  We have

$$\begin{aligned}
c^{\lambda}_{[12;00;0]}(M)=&c^{\lambda}_{[1;00;0]}(M)\,c^{\lambda}_{[2;00;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&0\\&&0\end{smallmatrix}\right)\right)\\
=&(m_{13}-m_{12})(m_{13}-m_{22}+1)(m_{23}-m_{22}),\\
c^{\lambda}_{[12;00;1]}(M)=&c^{\lambda}_{[1;00;0]}(M)\,c^{\lambda}_{[2;00;1]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&0\\&&0\end{smallmatrix}\right)\right)+c^{\lambda}_{[1;00;1]}(M)\,c^{\lambda}_{[2;00;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&0\\&&0\end{smallmatrix}\right)[-1]\right)\\
=&(m_{13}-m_{12})(m_{13}-m_{22}+1)\bar{C}_1\left(M\left(\begin{smallmatrix}-1&0&0\\&0&0\\&&0\end{smallmatrix}\right)\right)\chi_-\left(M\left(\begin{smallmatrix}-1&0&0\\&0&0\\&&0\end{smallmatrix}\right)\right)\\
&-C_2(M)(m_{23}-m_{22}-1)\\
=&(m_{13}-m_{12})(m_{13}-m_{22}+1)\bar{C}_1(M)\chi_-(M)-C_2(M)(m_{23}-m_{22}-1),\\
c^{\lambda}_{[12;00;2]}(M)=&c^{\lambda}_{[1;00;1]}(M)\,c^{\lambda}_{[2;00;1]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&0\\&&0\end{smallmatrix}\right)[-1]\right)\\
=&-C_2(M)\bar{C}_1\left(M\left(\begin{smallmatrix}-1&0&0\\&0&0\\&&0\end{smallmatrix}\right)[-1]\right)\chi_-\left(M\left(\begin{smallmatrix}-1&0&0\\&0&0\\&&0\end{smallmatrix}\right)[-1]\right)\\
=&-C_2(M)(\bar{C}_1(M)-1)\chi_-(M).
\end{aligned}$$

Here we use the relations (4.6), (4.39) and (4.67).

Next, we compute the case of formula 5.

- the case $(s,t)=(2,2)$.
  We have

$$\begin{aligned}
c^{\lambda}_{[13;22;0]}(M)=&c^{\lambda}_{[1;11;0]}(M)\,c^{\lambda}_{[3;11;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&-1\end{smallmatrix}\right)\right)=(m_{13}-m_{12})(m_{22}-m_{33}),\\
c^{\lambda}_{[13;22;1]}(M)=&c^{\lambda}_{[1;11;1]}(M)\,c^{\lambda}_{[3;11;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&-1\end{smallmatrix}\right)[-1]\right)=-\bar{E}(M).
\end{aligned}$$

- the case $(s,t)=(1,2)$.
  We have

$$\begin{aligned}
c^{\lambda}_{[13;12;0]}(M)=&c^{\lambda}_{[1;11;0]}(M)\,c^{\lambda}_{[3;01;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&-1\end{smallmatrix}\right)\right)+c^{\lambda}_{[1;01;0]}(M)\,c^{\lambda}_{[3;11;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&0\end{smallmatrix}\right)\right)\\
=&-(m_{13}-m_{12})(m_{22}-m_{33})-(m_{13}-m_{12})(m_{22}-m_{33})\\
=&-2(m_{13}-m_{12})(m_{22}-m_{33}),\\
c^{\lambda}_{[13;12;1]}(M)=&c^{\lambda}_{[1;11;0]}(M)\,c^{\lambda}_{[3;01;1]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&-1\end{smallmatrix}\right)\right)+c^{\lambda}_{[1;01;1]}(M)\,c^{\lambda}_{[3;11;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&0\end{smallmatrix}\right)[-1]\right)\\
&+c^{\lambda}_{[1;11;1]}(M)\,c^{\lambda}_{[3;01;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&-1\end{smallmatrix}\right)[-1]\right)\\
=&-(m_{13}-m_{12})(m_{22}-m_{33})\chi_+(M)+\bar{F}(M)+\bar{E}(M),\\
c^{\lambda}_{[13;12;2]}(M)=&c^{\lambda}_{[1;11;1]}(M)\,c^{\lambda}_{[3;01;1]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&-1\end{smallmatrix}\right)[-1]\right)+c^{\lambda}_{[1;01;2]}(M)\,c^{\lambda}_{[3;11;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&0\end{smallmatrix}\right)[-2]\right)
\end{aligned}$$

$$\begin{aligned}
&=\bar{E}(M)\chi_{+}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&-1\end{smallmatrix}\right)[-1]\right)-C_2(M)\chi_{+}(M)\\
&=(\bar{E}(M)-C_2(M))\chi_{+}(M).
\end{aligned}$$

Here we use the relation (4.5).

- the case $(s,t)=(1,1)$.
  We have

$$\begin{aligned}
c^{\lambda}_{[13;11;0]}(M)=&c^{\lambda}_{[1;11;0]}(M)\,c^{\lambda}_{[3;00;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&-1\end{smallmatrix}\right)\right)+c^{\lambda}_{[1;00;0]}(M)\,c^{\lambda}_{[3;11;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&0\\&&0\end{smallmatrix}\right)\right)\\
=&(m_{13}-m_{12})(m_{22}-m_{33})-(m_{13}-m_{12})(m_{13}-m_{22}+1)\\
=&(m_{13}-m_{12})(2m_{22}-m_{13}-m_{33}-1),\\
c^{\lambda}_{[13;11;1]}(M)=&c^{\lambda}_{[1;00;1]}(M)\,c^{\lambda}_{[3;11;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&0\\&&0\end{smallmatrix}\right)[-1]\right)+c^{\lambda}_{[1;11;1]}(M)\,c^{\lambda}_{[3;00;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&-1\end{smallmatrix}\right)[-1]\right)\\
=&C_2(M)-\bar{E}(M).
\end{aligned}$$

- the case $(s,t)=(0,2)$.
  We have

$$\begin{aligned}
c^{\lambda}_{[13;02;0]}(M)=&c^{\lambda}_{[1;01;0]}(M)\,c^{\lambda}_{[3;01;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&0\end{smallmatrix}\right)\right)=(m_{13}-m_{12})(m_{22}-m_{33}),\\
c^{\lambda}_{[13;02;1]}(M)=&c^{\lambda}_{[1;01;0]}(M)\,c^{\lambda}_{[3;01;1]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&0\end{smallmatrix}\right)\right)+c^{\lambda}_{[1;01;1]}(M)\,c^{\lambda}_{[3;01;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&0\end{smallmatrix}\right)[-1]\right)\\
=&(m_{13}-m_{12})(m_{22}-m_{33})\chi_{+}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&0\end{smallmatrix}\right)\right)-\bar{F}(M)\\
=&(m_{13}-m_{12})(m_{22}-m_{33})\chi_{+}^{(1)}(M)-\bar{F}(M),\\
c^{\lambda}_{[13;02;2]}(M)=&c^{\lambda}_{[1;01;2]}(M)\,c^{\lambda}_{[3;01;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&0\end{smallmatrix}\right)[-2]\right)+c^{\lambda}_{[1;01;1]}(M)\,c^{\lambda}_{[3;01;1]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&0\end{smallmatrix}\right)[-1]\right)\\
=&C_2(M)\chi_{+}(M)-\bar{F}(M)\chi_{+}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&0\end{smallmatrix}\right)[-1]\right)\\
=&C_2(M)\chi_{+}(M)-\bar{F}(M)\chi_{+}^{(1)}(M),\\
c^{\lambda}_{[13;02;3]}(M)=&c^{\lambda}_{[1;01;2]}(M)\,c^{\lambda}_{[3;01;1]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&0\end{smallmatrix}\right)[-2]\right)\\
=&C_2(M)\chi_{+}(M)\chi_{+}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&0\end{smallmatrix}\right)[-2]\right)\\
=&C_2(M)\chi_{+}^{(1)}(M).
\end{aligned}$$

Here we use the relations (4.5) and (4.11).

- the case $(s,t)=(0,1)$.
  We have

$$\begin{aligned}
c^{\lambda}_{[13;01;0]}(M)=&c^{\lambda}_{[1;01;0]}(M)\,c^{\lambda}_{[3;00;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&0\end{smallmatrix}\right)\right)+c^{\lambda}_{[1;00;0]}(M)\,c^{\lambda}_{[3;01;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&0\\&&0\end{smallmatrix}\right)\right)\\
=&-(m_{13}-m_{12})(m_{22}-m_{33})+(m_{13}-m_{12})(m_{13}-m_{22}+1)\\
=&-(m_{13}-m_{12})(2m_{22}-m_{13}-m_{33}-1),\\
c^{\lambda}_{[13;01;1]}(M)=&c^{\lambda}_{[1;00;1]}(M)\,c^{\lambda}_{[3;01;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&0\\&&0\end{smallmatrix}\right)[-1]\right)\\
&+c^{\lambda}_{[1;01;1]}(M)\,c^{\lambda}_{[3;00;0]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&0\end{smallmatrix}\right)[-1]\right)+c^{\lambda}_{[1;00;0]}(M)\,c^{\lambda}_{[3;01;1]}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&0\\&&0\end{smallmatrix}\right)\right)
\end{aligned}$$

$$\begin{aligned}
&=-\,C_2(M)+\bar{F}(M)+(m_{13}-m_{12})(m_{13}-m_{22}+1)\chi_+\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;0\\0\end{smallmatrix}\right)\right)\\
&=\bar{F}(M)-C_2(M)+(m_{13}-m_{12})(m_{13}-m_{22}+1)\chi_+(M),\\
c^{\lambda}_{[13;01;2]}(M)&=c^{\lambda}_{[1;00;1]}(M)\,c^{\lambda}_{[3;01;1]}\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;0\\0\end{smallmatrix}\right)[-1]\right)+c^{\lambda}_{[1;01;2]}(M)\,c^{\lambda}_{[3;00;0]}\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;-1\\0\end{smallmatrix}\right)[-2]\right)\\
&=-\,C_2(M)\chi_+\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;0\\0\end{smallmatrix}\right)[-1]\right)-C_2(M)\chi_+(M)\\
&=-\,2C_2(M)\chi_+(M).
\end{aligned}$$

Here we use the relation (4.5).

- the case $(s,t)=(0,0)$.
  We have

$$\begin{aligned}
c^{\lambda}_{[13;00;0]}(M)&=c^{\lambda}_{[1;00;0]}(M)\,c^{\lambda}_{[3;00;0]}\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;0\\0\end{smallmatrix}\right)\right)=-(m_{13}-m_{12})(m_{13}-m_{22}+1),\\
c^{\lambda}_{[13;00;1]}(M)&=c^{\lambda}_{[1;00;1]}(M)\,c^{\lambda}_{[3;00;0]}\left(M\left(\begin{smallmatrix}-1\;0\;0\\0\;0\\0\end{smallmatrix}\right)[-1]\right)=C_2(M).
\end{aligned}$$

At last, we compute the case of formula 6.

- the case $(s,t)=(2,2)$.
  We have

$$\begin{aligned}
c^{\lambda}_{[23;22;0]}(M)&=c^{\lambda}_{[2;11;0]}(M)\,c^{\lambda}_{[3;11;0]}\left(M\left(\begin{smallmatrix}0\;-1\;0\\0\;-1\\-1\end{smallmatrix}\right)\right)=m_{22}-m_{33},\\
c^{\lambda}_{[23;22;1]}(M)&=c^{\lambda}_{[2;11;1]}(M)\,c^{\lambda}_{[3;11;0]}\left(M\left(\begin{smallmatrix}0\;-1\;0\\0\;-1\\-1\end{smallmatrix}\right)[-1]\right)=-\bar{D}(M)\chi_-(M).
\end{aligned}$$

- the case $(s,t)=(1,2)$.
  We have

$$\begin{aligned}
c^{\lambda}_{[23;12;0]}(M)&=c^{\lambda}_{[2;11;0]}(M)\,c^{\lambda}_{[3;01;0]}\left(M\left(\begin{smallmatrix}0\;-1\;0\\0\;-1\\-1\end{smallmatrix}\right)\right)+c^{\lambda}_{[2;01;0]}(M)\,c^{\lambda}_{[3;11;0]}\left(M\left(\begin{smallmatrix}0\;-1\;0\\0\;-1\\0\end{smallmatrix}\right)\right)\\
&=-\,(m_{22}-m_{33})-(m_{22}-m_{33})=-2(m_{22}-m_{33}),\\
c^{\lambda}_{[23;12;1]}(M)&=c^{\lambda}_{[2;11;0]}(M)\,c^{\lambda}_{[3;01;1]}\left(M\left(\begin{smallmatrix}0\;-1\;0\\0\;-1\\-1\end{smallmatrix}\right)\right)+c^{\lambda}_{[2;01;1]}(M)\,c^{\lambda}_{[3;11;0]}\left(M\left(\begin{smallmatrix}0\;-1\;0\\0\;-1\\0\end{smallmatrix}\right)[-1]\right)\\
&\qquad+c^{\lambda}_{[2;11;1]}(M)\,c^{\lambda}_{[3;01;0]}\left(M\left(\begin{smallmatrix}0\;-1\;0\\0\;-1\\-1\end{smallmatrix}\right)[-1]\right)\\
&=-\,(m_{22}-m_{33})\chi_+\left(M\left(\begin{smallmatrix}0\;-1\;0\\0\;-1\\-1\end{smallmatrix}\right)\right)+\bar{C}_1(M)+\bar{D}(M)\chi_-(M)\\
&=\bar{C}_1(M)-(m_{22}-m_{33})(\chi_-(M)+\chi_+^{(-1)}(M))+\delta(M)\chi_-(M)\\
&=\bar{C}_1(M)-(m_{22}-m_{33})+\delta(M)\chi_-(M),\\
c^{\lambda}_{[23;12;2]}(M)&=c^{\lambda}_{[2;11;1]}(M)\,c^{\lambda}_{[3;01;1]}\left(M\left(\begin{smallmatrix}0\;-1\;0\\0\;-1\\-1\end{smallmatrix}\right)[-1]\right)\\
&=\bar{D}(M)\chi_-(M)\chi_+\left(M\left(\begin{smallmatrix}0\;-1\;0\\0\;-1\\-1\end{smallmatrix}\right)[-1]\right)=0.
\end{aligned}$$

Here we use the relations (4.5), (4.9) and (4.10).

- the case $(s,t)=(1,1)$.
  We have

$$c^{\lambda}_{[23;11;0]}(M)=c^{\lambda}_{[2;11;0]}(M)\,c^{\lambda}_{[3;00;0]}\left(M\left(\begin{smallmatrix}0\;-1\;0\\0\;-1\\-1\end{smallmatrix}\right)\right)+c^{\lambda}_{[2;00;0]}(M)\,c^{\lambda}_{[3;11;0]}\left(M\left(\begin{smallmatrix}0\;-1\;0\\0\;0\\0\end{smallmatrix}\right)\right)$$

$$\begin{aligned}
&=(m_{22}-m_{33})-(m_{23}-m_{22})=2m_{22}-m_{23}-m_{33},\\
c^{\lambda}_{[23;11;1]}(M) &=c^{\lambda}_{[2;00;1]}(M)\, c^{\lambda}_{[3;11;0]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)[-1]\right)+c^{\lambda}_{[2;11;1]}(M)\, c^{\lambda}_{[3;00;0]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&-1\end{smallmatrix}\right)[-1]\right)\\
&=-\bar{C}_1(M)\chi_-(M)-\bar{D}(M)\chi_-(M)=-(\bar{C}_1(M)+\bar{D}(M))\chi_-(M).
\end{aligned}$$

- the case $(s,t)=(0,2)$.

  We have

$$\begin{aligned}
c^{\lambda}_{[23;02;0]}(M) &=c^{\lambda}_{[2;01;0]}(M)\, c^{\lambda}_{[3;01;0]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&0\end{smallmatrix}\right)\right)=m_{22}-m_{33},\\
c^{\lambda}_{[23;02;1]}(M) &=c^{\lambda}_{[2;01;0]}(M)\, c^{\lambda}_{[3;01;1]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&0\end{smallmatrix}\right)\right)+c^{\lambda}_{[2;01;1]}(M)\, c^{\lambda}_{[3;01;0]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&0\end{smallmatrix}\right)[-1]\right)\\
&=(m_{22}-m_{33})\chi_+\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&0\end{smallmatrix}\right)\right)-\bar{C}_1(M)=-\{\bar{C}_1(M)-(m_{22}-m_{33})\chi_+(M)\},\\
c^{\lambda}_{[23;02;2]}(M) &=c^{\lambda}_{[2;01;1]}(M)\, c^{\lambda}_{[3;01;1]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&0\end{smallmatrix}\right)[-1]\right)=-\bar{C}_1(M)\chi_+\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&0\end{smallmatrix}\right)[-1]\right)\\
&=-\bar{C}_1(M)\chi_+(M).
\end{aligned}$$

Here we use the relation (4.5).

- the case $(s,t)=(0,1)$.

  We have

$$\begin{aligned}
c^{\lambda}_{[23;01;0]}(M) &=c^{\lambda}_{[2;01;0]}(M)\, c^{\lambda}_{[3;00;0]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&0\end{smallmatrix}\right)\right)+c^{\lambda}_{[2;00;0]}(M)\, c^{\lambda}_{[3;01;0]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)\right)\\
&=-(m_{22}-m_{33})+(m_{23}-m_{22})=-(2m_{22}-m_{23}-m_{33}),\\
c^{\lambda}_{[23;01;1]}(M) &=c^{\lambda}_{[2;00;1]}(M)\, c^{\lambda}_{[3;01;0]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)[-1]\right)\\
&\quad+c^{\lambda}_{[2;01;1]}(M)\, c^{\lambda}_{[3;00;0]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&-1\\&&0\end{smallmatrix}\right)[-1]\right)+c^{\lambda}_{[2;00;0]}(M)\, c^{\lambda}_{[3;01;1]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)\right)\\
&=\bar{C}_1(M)\chi_-(M)+\bar{C}_1(M)+(m_{23}-m_{22})\chi_+\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)\right)\\
&=\bar{C}_1(M)(1+\chi_-(M)+\chi_+^{(-1)}(M))=2\bar{C}_1(M),\\
c^{\lambda}_{[23;01;2]}(M) &=c^{\lambda}_{[2;00;1]}(M)\, c^{\lambda}_{[3;01;1]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)[-1]\right)\\
&=\bar{C}_1(M)\chi_-(M)\chi_+\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)[-1]\right)=0.
\end{aligned}$$

(4.5), (4.9), (4.10) and (4.15).

- the case $(s,t)=(0,0)$.

  We have

$$\begin{aligned}
c^{\lambda}_{[23;00;0]}(M) &=c^{\lambda}_{[2;00;0]}(M)\, c^{\lambda}_{[3;00;0]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)\right)=-(m_{23}-m_{22}),\\
c^{\lambda}_{[23;00;1]}(M) &=c^{\lambda}_{[2;00;1]}(M)\, c^{\lambda}_{[3;00;0]}\left(M\left(\begin{smallmatrix}0&-1&0\\&0&0\\&&0\end{smallmatrix}\right)[-1]\right)=-\bar{C}_1(M)\chi_-(M).
\end{aligned}$$

$\square$

For $1\le i\le j\le 3$, we define a linear map $i^{\lambda}_{-\mathbf{e}_i-\mathbf{e}_j}\colon V_{\lambda[-ij]}\to V_{\lambda}\otimes_{\mathbf{C}} V_{-2\mathbf{e}_3}$ by

$$i^{\lambda}_{-\mathbf{e}_i-\mathbf{e}_j}=(T_{\hat{\lambda}}\otimes_{\mathbf{C}} T_{2\mathbf{e}_1})\circ i^{\widehat{\lambda}}_{\mathbf{e}_{4-j}+\mathbf{e}_{4-i}}\circ T_{\lambda-\mathbf{e}_i-\mathbf{e}_j}.$$

By Proposition 3.2, we see that

$$X \circ i^{\lambda}_{-\mathbf{e}_i-\mathbf{e}_j} = i^{\lambda}_{-\mathbf{e}_i-\mathbf{e}_j} \circ \omega^2(X), \quad X \in \mathfrak{g}.$$

Since $\omega^2 = \mathrm{id}_{\mathfrak{g}}$, $i^{\lambda}_{-\mathbf{e}_i-\mathbf{e}_j}$ is a non-zero generator of $\mathrm{Hom}_K(V_{\lambda[-ij]}, V_\lambda \otimes_{\mathbf{C}} V_{-2\mathbf{e}_3})$, which is unique up to scalar multiple. So, we obtain the following proposition from Proposition 4.4 easily.

Proposition 4.5. *For $1 \leq i \leq j \leq 3$ and G-pattern M of type $\lambda[-ij]$, the image of the monomial basis $f(M)$ by the injector $i^{\lambda}_{-\mathbf{e}_i-\mathbf{e}_j} \colon V_{\lambda[-ij]} \to V_\lambda \otimes_{\mathbf{C}} V_{-2\mathbf{e}_3}$ is given by*

$$i^{\lambda}_{-\mathbf{e}_i-\mathbf{e}_j}(f(M)) = \sum_{0 \leq k \leq l \leq 2} \left\{ \sum_{m=0}^{r_{[4-j\,4-i;kl]}} c^{\widehat{\lambda}}_{[4-j\,4-i;kl;m]}(\hat{M}) f\left(M \begin{pmatrix} \begin{smallmatrix} \mathbf{e}_i+\mathbf{e}_j \\ l\ 0 \\ k \end{smallmatrix} \end{pmatrix} [-m] \right) \right\} \otimes f\begin{pmatrix} \begin{smallmatrix} 0\ 0\ -2 \\ 0\ -l \\ -k \end{smallmatrix} \end{pmatrix}$$

In the right hand side of the above formula, we put $f(M') = 0$ if M' is a triangular array which does not satisfy the condition (3.1) of G-patterns.

5. $(\mathfrak{g}, K)$-MODULE STRUCTURE OF PRINCIPAL SERIES REPRESENTATION OF $Sp(3,\mathbf{R})$

5.1. **Irreducible decomposition of** $(\pi_{(\sigma,\nu)}|_K, H_{(\sigma,\nu)})$**.** For an irreducible finite dimensional representation (τ, V) of K, we denote by (τ^*, V^*) its contragradient representation. For a basis $\{v_i\}_{i\in I}$ of V, we denote by $\{v_i^*\}_{i \in I}$ its dual basis of V^*. Let L^+ be the set of dominant weights of length three.

We set

$$L^2_{(M_{\min},\sigma)}(K) = \{ f \in L^2(K) \mid f(mx) = \sigma(m) f(x) \text{ for a.e. } m \in M_{\min},\ x \in K \}$$

and give K-module structure by the right regular action of K. Then the restriction map $r_K \colon H_{(\sigma,\nu)} \ni f \mapsto f|_K \in L^2_{(M_{\min},\sigma)}(K)$ is an isomorphism of K-modules.

$L^2(K)$ has a $K \times K$-bimodule structure by the two sided regular action:

$$((k_1,k_2)f)(x) = f(k_1^{-1} x k_2), \quad x \in K,\ f \in L^2(K),\ (k_1,k_2) \in K \times K,$$

and a homomorphism $\Phi_\lambda \colon V_\lambda^* \otimes_{\mathbf{C}} V_\lambda \to L^2(K)$ of $K \times K$-bimodules by

$$w \otimes v \mapsto (x \mapsto \langle w, \tau_\lambda(x) v \rangle)$$

where $\langle , \rangle$ is a canonical pairing on $V_\lambda^* \otimes_{\mathbf{C}} V_\lambda$. Then the Peter-Weyl theorem tells that

$$\widehat{\bigoplus_{\lambda \in L^+}} \Phi_\lambda \colon \widehat{\bigoplus_{\lambda \in L^+}} V_\lambda^* \otimes_{\mathbf{C}} V_\lambda \to L^2(K)$$

is an isomorphism as $K \times K$-bimodules. Here $\widehat{\bigoplus}$ means a Hilbert space direct sum.

Since $L^2_{(M_{\min},\sigma)}(K) \subset L^2(K)$, we have an irreducible decomposition of $L^2_{(M_{\min},\sigma)}(K)$:

$$L^2_{(M_{\min},\sigma)}(K) \simeq \widehat{\bigoplus_{\lambda \in L^+}} (V_\lambda^*[\sigma]) \otimes_{\mathbf{C}} V_\lambda.$$

Here $V_\lambda^*[\sigma]$ is the σ-isotypic component in $(\tau_\lambda^*|_{M_{\min}}, V^*)$, i.e.,

$$V_\lambda^*[\sigma] = \bigoplus_{\gamma \equiv (\sigma_1,\sigma_2,\sigma_3) \bmod 2} V_\lambda^*(\gamma) = \bigoplus_{M \in G_\sigma(\lambda)} \mathbf{C} f(M)^*$$

where $G_\sigma(\lambda) = \{ M \in G(\lambda) \mid \gamma^M \equiv (\sigma_1,\sigma_2,\sigma_3) \bmod 2 \}$.

From above arguments, We obtain an isomorphism

$$r_K^{-1} \circ \widehat{\bigoplus_{\lambda \in L^+}} \Phi_\lambda \colon \widehat{\bigoplus_{\lambda \in L^+}} (V_\lambda^*[\sigma]) \otimes_{\mathbf{C}} V_\lambda \to H_{(\sigma,\nu)}.$$

Now we define elementary function $s(M,N) \in H_{(\sigma,\nu)}$ by $s(M,N) = r_K^{-1} \circ \Phi_\lambda(f(M)^* \otimes f(N))$ for $M \in G_\sigma(\lambda)$ and $N \in G(\lambda)$. Let $l(M)$ be the order of the set $\{ N \in G(\lambda) \mid \gamma^N > \gamma^M \ \text{ or } \ N =$

$M[k]$ for some $k \geq 0\}$ for a G-pattern M of type λ. For a G-pattern M of type λ, let $S_\lambda(M)$ be a column vector of degree $d_\lambda = \dim V_\lambda$ with its $l(N)$-th component $s(M,N)$ ($N \in G(\lambda)$), i.e.,

$$S_\lambda(M) = {}^t\!\left(s\left(M, \left(\begin{smallmatrix}\lambda_1 & & \lambda_2 & & \lambda_3\\ & \lambda_1 & & \lambda_2 & \\ & & \lambda_1 & & \end{smallmatrix}\right)\right), \cdots, \overset{\overset{l(N)\text{-th}}{\downarrow}}{s(M,N)}, \cdots, s\left(M, \left(\begin{smallmatrix}\lambda_1 & & \lambda_2 & & \lambda_3\\ & \lambda_2 & & \lambda_3 & \\ & & \lambda_3 & & \end{smallmatrix}\right)\right) \right).$$

Moreover we denote by $\langle S_\lambda(M)\rangle$ the subspace of $H_{(\sigma,\nu)}$ generated by the functions in the entries of the vector $S_\lambda(M)$, i.e., $\langle S_\lambda(M)\rangle = \bigoplus_{N\in G(\lambda)} \mathbf{C}s(M,N)$ ($\simeq V_\lambda$). We take the marking $\{s(M,N)\}_{N\in G(\lambda)}$ for the simple K-module $\langle S_\lambda(M)\rangle$.

Proposition 5.1. *As a unitary representation of K, it has an irreducible decomposition:*

$$H_{(\sigma,\nu)} \simeq \widehat{\bigoplus_{\lambda\in L^+}} (V_\lambda^*[\sigma]) \otimes_{\mathbf{C}} V_\lambda.$$

The direct sum

$$\bigoplus_{M\in G_\sigma(\lambda)} \langle S_\lambda(M)\rangle$$

is the τ_λ-isotypic component $(V_\lambda^*[\sigma]) \otimes_{\mathbf{C}} V_\lambda$ in $H_{(\sigma,\nu)}$.

5.2. **General setting.** The K-finite part $H_{(\sigma,\nu),K}$ of $H_{(\sigma,\nu)}$ is also a $\mathfrak{k}$-module. Because of the Cartan decomposition $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$, in order to describe the action of $\mathfrak{g}$ or $\mathfrak{g}_{\mathbf{C}}$ it suffices to investigate the action of $\mathfrak{p}$ or $\mathfrak{p}_{\mathbf{C}} = \mathfrak{p}_+ \oplus \mathfrak{p}_-$.

For a K-type $(\tau_\lambda, V_\lambda)$ of $(\pi_{(\sigma,\nu)}, H_{(\sigma,\nu),K})$ and a nonzero K-homomorphism $\eta\colon V_\lambda \to H_{(\sigma,\nu),K}$, we define linear map $\tilde{\eta}\colon \mathfrak{p}_{\mathbf{C}} \otimes_{\mathbf{C}} V_\lambda \to H_{(\sigma,\nu),K}$ by $X \otimes v \mapsto X \cdot \eta(v)$. Then $\tilde{\eta}$ is K-homomorphism with $\mathfrak{p}_{\mathbf{C}}$ endowed with the adjoint action Ad of K.

Since

$$V_\lambda \otimes_{\mathbf{C}} \mathfrak{p}_+ \simeq V_\lambda \otimes_{\mathbf{C}} V_{2\mathbf{e}_1} \simeq \bigoplus_{1\leq i\leq j\leq 3} V_{\lambda[+ij]}$$

$$\left(\text{resp. } V_\lambda \otimes_{\mathbf{C}} \mathfrak{p}_- \simeq V_\lambda \otimes_{\mathbf{C}} V_{-2\mathbf{e}_3} \simeq \bigoplus_{1\leq i\leq j\leq 3} V_{\lambda[-ij]}\right),$$

there are six injective K-homomorphisms

$$I^\lambda_{+ij} = (\mathrm{id}_{V_\lambda} \otimes_{\mathbf{C}} i_{\mathfrak{p}_+}) \circ i^\lambda_{\mathbf{e}_i+\mathbf{e}_j} \colon V_{\lambda[+ij]} \to V_\lambda \otimes_{\mathbf{C}} \mathfrak{p}_+, \quad 1 \leq i \leq j \leq 3$$

$$\left(\text{ resp. } I^\lambda_{-ij} = (\mathrm{id}_{V_\lambda} \otimes_{\mathbf{C}} i_{\mathfrak{p}_-}) \circ i^\lambda_{-\mathbf{e}_i-\mathbf{e}_j} \colon V_{\lambda[-ij]} \to V_\lambda \otimes_{\mathbf{C}} \mathfrak{p}_-, \quad 1 \leq i \leq j \leq 3\right)$$

for general λ. Then we define $\mathbf{C}$-linear maps

$$\Gamma^\lambda_{\pm ij}\colon \mathrm{Hom}_K(V_\lambda, H_{(\sigma,\nu),K}) \to \mathrm{Hom}_K(V_{\lambda[\pm ij]}, H_{(\sigma,\nu),K}) \quad, \ 1 \leq i \leq j \leq 3$$

by $\eta \mapsto \tilde{\eta} \circ I^\lambda_{\pm ij}$.

Now we settle two purposes of this paper:

- **(i):** Describe the injective K-homomorphism $I^\lambda_{\pm ij}$ in terms of the monomial basis.
- **(ii):** Determine the matrix representations of the linear homomorphisms $\Gamma^\lambda_{\pm ij}$ with respect to the induced basis defined in the next subsection.

We have already accomplished the first purpose in Lemma 3.4, Proposition 4.4 and 4.5. We accomplish the second purpose in the subsection 5.5. As a result, we obtain infinite number of 'contiguous relations', a kind of system of differential-difference relations among vectors in $H_{(\sigma,\nu)}[\tau_\lambda]$ and $H_{(\sigma,\nu)}[\tau_{\lambda[\pm ij]}]$. Here $H_{(\sigma,\nu)}[\tau]$ is τ-isotypic component of $H_{(\sigma,\nu)}$.

5.3. **The canonical blocks of elementary functions.** Let $\eta \colon V_\lambda \to H_{(\sigma,\nu),K}$ be a non-zero K-homomorphism, where V_λ is a simple $K-module$ with the marking $\{f(M)\}_{M\in G(\lambda)}$. Then we identify η with the column vector of degree $d_\lambda = \dim V_\lambda$ whose $l(N)$-th component is $\eta\,(f(N))$ for $N \in G(\lambda)$, i.e.,

$$ {}^{t}\!\left(\eta\left(f\left(\begin{smallmatrix}\lambda_1 & & \lambda_2 & & \lambda_3\\ & \lambda_1 & & \lambda_2 & \\ & & \lambda_1 & & \end{smallmatrix}\right)\right), \cdots, \overset{\substack{l(N)\text{-th}\\ \downarrow}}{\eta\,(f(N))}, \cdots, \eta\left(f\left(\begin{smallmatrix}\lambda_1 & & \lambda_2 & & \lambda_3\\ & \lambda_2 & & \lambda_3 & \\ & & \lambda_3 & & \end{smallmatrix}\right)\right)\right). $$

By this identification, we identify $S_\lambda\,(M)$ with the injective K-homomorphism

$$ V_\lambda \ni f(N) \mapsto s(M,N) \in H_{(\sigma,\nu),K},\ N \in G(\lambda) $$

for $M \in G_\sigma(\lambda)$. We note that $\{S_\lambda\,(M)\}_{M\in G_\sigma(\lambda)}$ is a basis of $\mathrm{Hom}_K(V_\lambda, H_{(\sigma,\nu),K})$ and we call it *the induced basis from the monomial basis.*

We define a certain matrix of elementary functions corresponding to the induced basis $\{S_\lambda\,(M)\}_{M\in G_\sigma(\lambda)}$ of $\mathrm{Hom}_K(V_\lambda, H_{(\sigma,\nu),K})$ for each K-type τ_λ of our principal series representation $\pi_{(\sigma,\nu)}$.

Definition 5.2. *For $M \in G_\sigma(\lambda)$, let d^σ_λ and $l^\sigma(M)$ be the orders of the set $G_\sigma(\lambda)$ and $\{N \in G_\sigma(\lambda) \mid l(M) \le l(N)\}$, respectively. The $d_\lambda \times d^\sigma_\lambda$ matrix $\mathbf{S}_\sigma(\lambda)$ whose $l^\sigma(M)$-th column is $S_\lambda\,(M)$ for $M \in G_\sigma(\lambda)$ is called the canonical block of elementary functions for τ_λ-isotypic component.*

5.4. **The $\mathfrak{p}_\pm$-matrix corresponding to $I^\lambda_{\pm ij}$.** In this subsection, we define $\mathfrak{p}_\pm$-matrix $\mathcal{C}^\lambda_{\pm ij}$ of size $d_{\lambda[\pm ij]} \times d_\lambda$ corresponding to $I^\lambda_{\pm ij}$ with respect to the monomial basis.

Definition 5.3. *We define a $\mathfrak{p}_\pm$-matrix $\mathcal{C}^\lambda_{\pm ij}$ of size $d_{\lambda[\pm ij]} \times d_\lambda$ as follows.*
(i) For $1 \le i \le j \le 3$, we define a $\mathfrak{p}_+$-matrix $\mathcal{C}^\lambda_{+ij} \in M(d_{\lambda[+ij]}, d_\lambda, \mathbf{C}) \otimes \mathfrak{p}_+$ by

$$ \begin{aligned} \mathcal{C}^\lambda_{+ij} = &\left\{\sum_{m=0}^{r_{[ij;22]}} L^\lambda_{+ij}\left(\left(\begin{smallmatrix}0 & & -2\\ & -2 & \end{smallmatrix}\right)[-m]\right)\right\} \otimes X_{+11} + \left\{\sum_{m=0}^{r_{[ij;12]}} L^\lambda_{+ij}\left(\left(\begin{smallmatrix}0 & & -2\\ & -1 & \end{smallmatrix}\right)[-m]\right)\right\} \otimes X_{+12} \\ &+ \left\{\sum_{m=0}^{r_{[ij;11]}} L^\lambda_{+ij}\left(\left(\begin{smallmatrix}0 & & -1\\ & -1 & \end{smallmatrix}\right)[-m]\right)\right\} \otimes X_{+13} + \left\{\sum_{m=0}^{r_{[ij;02]}} L^\lambda_{+ij}\left(\left(\begin{smallmatrix}0 & & -2\\ & 0 & \end{smallmatrix}\right)[-m]\right)\right\} \otimes X_{+22} \\ &+ \left\{\sum_{m=0}^{r_{[ij;01]}} L^\lambda_{+ij}\left(\left(\begin{smallmatrix}0 & & -1\\ & 0 & \end{smallmatrix}\right)[-m]\right)\right\} \otimes X_{+23} + \left\{\sum_{m=0}^{r_{[ij;00]}} L^\lambda_{+ij}\left(\left(\begin{smallmatrix}0 & & 0\\ & 0 & \end{smallmatrix}\right)[-m]\right)\right\} \otimes X_{+33}. \end{aligned} $$

Here $L^\lambda_{+ij}\left(\left(\begin{smallmatrix}0 & & -l\\ & -k & \end{smallmatrix}\right)[-m]\right)$ is a matrix of size $d_{\lambda[+ij]} \times d_\lambda$ whose $l(M)$-th row is given by

$$ \begin{cases} (\overbrace{0, \cdots, 0}^{l-1}, c^\lambda_{[ij;kl;m]}\,(M), \overbrace{0, \cdots, 0}^{d_\lambda - l}) & \left(l = l\left(M\left(\begin{smallmatrix} & -\mathbf{e}_i-\mathbf{e}_j & \\ 0 & & -l\\ & -k & \end{smallmatrix}\right)[-m]\right)\right) \\ & \textit{if } M\left(\begin{smallmatrix} & -\mathbf{e}_i-\mathbf{e}_j & \\ 0 & & -l\\ & -k & \end{smallmatrix}\right)[-m] \in G(\lambda) \\ \mathbf{0} & \textit{otherwise} \end{cases} $$

for $M \in G(\lambda[+ij])$.
(ii) For $1 \le i \le j \le 3$, we define a $\mathfrak{p}_-$-matrix $\mathcal{C}^\lambda_{-ij} \in M(d_{\lambda[-ij]}, d_\lambda, \mathbf{C}) \otimes \mathfrak{p}_-$ by

$$ \begin{aligned} \mathcal{C}^\lambda_{-ij} = &\left\{\sum_{m=0}^{r_{[4-j\,4-i;22]}} L^\lambda_{-ij}\left(\left(\begin{smallmatrix}2 & & 0\\ & 2 & \end{smallmatrix}\right)[-m]\right)\right\} \otimes X_{-11} - \left\{\sum_{m=0}^{r_{[4-j\,4-i;12]}} L^\lambda_{-ij}\left(\left(\begin{smallmatrix}2 & & 0\\ & 1 & \end{smallmatrix}\right)[-m]\right)\right\} \otimes X_{-12} \\ &+ \left\{\sum_{m=0}^{r_{[4-j\,4-i;11]}} L^\lambda_{-ij}\left(\left(\begin{smallmatrix}1 & & 0\\ & 1 & \end{smallmatrix}\right)[-m]\right)\right\} \otimes X_{-13} + \left\{\sum_{m=0}^{r_{[4-j\,4-i;02]}} L^\lambda_{-ij}\left(\left(\begin{smallmatrix}2 & & 0\\ & 0 & \end{smallmatrix}\right)[-m]\right)\right\} \otimes X_{-22} \end{aligned} $$

$$-\left\{\sum_{m=0}^{r_{[4-j\,4-i;01]}} L^{\lambda}_{-ij}\left(\left(\begin{smallmatrix}1\,0\\0\end{smallmatrix}\right)[-m]\right)\right\}\otimes X_{-23}+\left\{\sum_{m=0}^{r_{[4-j\,4-i;00]}} L^{\lambda}_{-ij}\left(\left(\begin{smallmatrix}0\,0\\0\end{smallmatrix}\right)[-m]\right)\right\}\otimes X_{-33}.$$

Here $L^{\lambda}_{-ij}\left(\left(\begin{smallmatrix}l\,0\\k\end{smallmatrix}\right)[-m]\right)$ *is a matrix of size* $d_{\lambda[-ij]}\times d_{\lambda}$ *whose* $l(M)$*-th row is given by*

$$\begin{cases}(\overbrace{0,\cdots,0}^{l-1},c^{\widehat{\lambda}}_{[4-j\,4-i;kl;m]}(\hat{M}),\overbrace{0,\cdots,0}^{d_{\lambda}-l}) & \left(\, l=l\left(M\left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\l\,0\\k\end{smallmatrix}\right)[-m]\right)\,\right)\\ & \textit{if}\ M\left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\l\,0\\k\end{smallmatrix}\right)[-m]\in G(\lambda)\\ \mathbf{0} & \textit{otherwise}\end{cases}$$

for $M\in G(\lambda[-ij])$.

Now we define $\mathcal{C}^{\lambda}_{\pm ij}S_{\lambda}(M)\in\left(H_{(\sigma,\nu),K}\right)^{\oplus d_{\lambda[\pm ij]}}\simeq\mathbf{C}^{d_{\lambda[\pm ij]}}\otimes_{\mathbf{C}}H_{(\sigma,\nu),K}$ by the action

$$(L\otimes X)(v\otimes f)=L(v)\otimes Xf,$$
$$\left(\ L\otimes X\in\mathrm{Hom}_{\mathbf{C}}(\mathbf{C}^{d_{\lambda}},\mathbf{C}^{d_{\lambda[\pm ij]}})\otimes_{\mathbf{C}}\mathfrak{p}_{\pm},\quad v\otimes f\in\mathbf{C}^{d_{\lambda}}\otimes_{\mathbf{C}}H_{(\sigma,\nu),K}\ \right)$$

for

$$\begin{aligned}&S_{\lambda}(M)\in\left(H_{(\sigma,\nu),K}\right)^{\oplus d_{\lambda}}\simeq\mathbf{C}^{d_{\lambda}}\otimes_{\mathbf{C}}H_{(\sigma,\nu),K},\\&\mathcal{C}^{\lambda}_{\pm ij}\in M(d_{\lambda[\pm ij]},d_{\lambda},\mathbf{C})\otimes_{\mathbf{C}}\mathfrak{p}_{\pm}\simeq\mathrm{Hom}_{\mathbf{C}}(\mathbf{C}^{d_{\lambda}},\mathbf{C}^{d_{\lambda[\pm ij]}})\otimes_{\mathbf{C}}\mathfrak{p}_{\pm}.\end{aligned}$$

By the definition of $\mathcal{C}^{\lambda}_{\pm ij}$, we note that the vector $\mathcal{C}^{\lambda}_{\pm ij}S_{\lambda}(M)$ is identified with the image of $S_{\lambda}(M)$ by $\Gamma^{\lambda}_{\pm ij}$.

5.5. **The contiguous relations.** To compute the matrix representation of $\Gamma^{\lambda}_{\pm ij}$ with respect to the induced basis, we prepare the following lemmas.

Lemma 5.4. *The root vectors* $X_{\pm ij}$ $(0\le i\le j\le 3)$ *in* $\mathfrak{p}_{\pm}$ *have the following expressions according to the Iwasawa decomposition of* $\mathfrak{g}$.

$$X_{+ij}=\begin{cases}2\sqrt{-1}E_{2e_i}+H_i+\kappa(E_{ii}), & i=j\\ (E_{e_i-e_j}+\sqrt{-1}E_{e_i+e_j})+\kappa(E_{ji}), & i<j\end{cases},$$
$$X_{-ij}=\begin{cases}-2\sqrt{-1}E_{2e_i}+H_i-\kappa(E_{ii}), & i=j\\ (E_{e_i-e_j}-\sqrt{-1}E_{e_i+e_j})-\kappa(E_{ij}), & i<j\end{cases}.$$

Proof. These are obtained by direct computation. □

Lemma 5.5. *The coefficients* $c^{\lambda}_{[ij;kl;m]}(M)$ *in Proposition 4.4 have the following relations:*

$$\begin{aligned}k_{ij}(M)\,c^{\lambda}_{[ij;22;m]}\left(M\left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\0\,2\\2\end{smallmatrix}\right)[m]\right)=&(m_{12}-m_{23}+1)\chi_{-}^{(-1)}(M)c^{\lambda}_{[ij;12;m-1]}\left(M\left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\0\,2\\2\end{smallmatrix}\right)[m]\right)\\&+(m_{11}-m_{22}+1)c^{\lambda}_{[ij;12;m]}\left(M\left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\0\,2\\2\end{smallmatrix}\right)[m]\right)\\&+(C_1(M)+1)c^{\lambda}_{[ij;11;m-1]}\left(M\left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\0\,2\\2\end{smallmatrix}\right)[m]\right)\\&+(m_{33}-m_{22}-1)c^{\lambda}_{[ij;11;m]}\left(M\left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\0\,2\\2\end{smallmatrix}\right)[m]\right)\end{aligned}$$

for $1\le i\le j\le 3$, $m\in\mathbf{Z}$ *and* $M\in G(\lambda)$. *Here*

$$c^{\lambda}_{[ij;kl;m]}\left(M\left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\0\,2\\2\end{smallmatrix}\right)[m]\right)=0\qquad\qquad \textit{if}\ r_{[ij;kl]}<m\quad\textit{or}\quad m<0.$$

and

$$\begin{aligned}&k_{11}(M)=-2m_{11}+2m_{13}, && k_{12}(M)=-2m_{11}+m_{13}+m_{23}-2,\\&k_{22}(M)=-2m_{11}+2m_{23}-2, && k_{13}(M)=-2m_{11}+m_{13}+m_{33}-3,\end{aligned}$$

$$k_{33}(M) = -2m_{11} + 2m_{33} - 4, \qquad k_{23}(m) = -2m_{11} + m_{23} + m_{33} - 4.$$

Proof. In order to prove the assertion, it suffices to confirm the equations

$$\begin{aligned}
(5.1)\quad & k_{ij}\left(M\left(\begin{smallmatrix}-\mathbf{e}_i\,-\mathbf{e}_j\\ 0\,-2\\ -2\end{smallmatrix}\right)[-m]\right) c^{\lambda}_{[ij;22;m]}(M)\\
&= (m_{12} - m_{23} - m + 1 + \delta_{2i} + \delta_{2j})\chi_-^{(-1)}\left(M\left(\begin{smallmatrix}-\mathbf{e}_i\,-\mathbf{e}_j\\ 0\,-2\\ -2\end{smallmatrix}\right)[-m]\right) c^{\lambda}_{[ij;12;m-1]}(M)\\
&+ (m_{11} - m_{22} - m + 1)c^{\lambda}_{[ij;12;m]}(M) + \left(C_1\left(M\left(\begin{smallmatrix}-\mathbf{e}_i\,-\mathbf{e}_j\\ 0\,-2\\ -2\end{smallmatrix}\right)[-m]\right) + 1\right) c^{\lambda}_{[ij;11;m-1]}(M)\\
&+ (m_{33} - m_{22} - m + 1 - \delta_{3i} - \delta_{3j})c^{\lambda}_{[ij;11;m]}(M)
\end{aligned}$$

for $1 \le i \le j \le 3$, $m \in \mathbf{Z}$ and $M \in G(\lambda[+ij])$. Since

$$\begin{aligned}
k_{ij}\left(M\left(\begin{smallmatrix}-\mathbf{e}_i\,-\mathbf{e}_j\\ 0\,-2\\ -2\end{smallmatrix}\right)[-m]\right) &= k_{ij}(M) + 2(1 - \delta_{ij}),\\
\chi_-^{(-1)}\left(M\left(\begin{smallmatrix}-\mathbf{e}_i\,-\mathbf{e}_j\\ 0\,-2\\ -2\end{smallmatrix}\right)[-m]\right) &= \chi_-^{(-1+\delta_{2i}+\delta_{2j})}(M),\\
C_1\left(M\left(\begin{smallmatrix}-\mathbf{e}_i\,-\mathbf{e}_j\\ 0\,-2\\ -2\end{smallmatrix}\right)[-m]\right) + 1 &= C_1(M) - m + 1 + \delta_{2i} \cdot \chi_-^{(\delta_{2j})}(M) + \delta_{2j} \cdot \chi_-(M),
\end{aligned}$$

(4.1), (4.4) and (4.6), the equations (5.1) are equivalent to the equations

$$\begin{aligned}
(5.2)\qquad & \{k_{ij}(M) + 2(1 - \delta_{ij})\}c^{\lambda}_{[ij;22;m]}(M)\\
&= (m_{12} - m_{23} - m + 1 + \delta_{2i} + \delta_{2j})\chi_-^{(-1+\delta_{2i}+\delta_{2j})}(M)c^{\lambda}_{[ij;12;m-1]}(M)\\
&\quad + (m_{11} - m_{22} - m + 1)c^{\lambda}_{[ij;12;m]}(M)\\
&\quad + \left(C_1(M) - m + 1 + \delta_{2i} \cdot \chi_-^{(\delta_{2j})}(M) + \delta_{2j} \cdot \chi_-(M)\right) c^{\lambda}_{[ij;11;m-1]}(M)\\
&\quad + (m_{33} - m_{22} - m + 1 - \delta_{3i} - \delta_{3j})c^{\lambda}_{[ij;11;m]}(M)
\end{aligned}$$

for $1 \le i \le j \le 3$, $m \in \mathbf{Z}$ and $M \in G(\lambda[+ij])$.

We prove the equations (5.2) by direct computation.

- the proof of the case of the $(i, j) = (1, 1)$.

Since $(r_{[11;22]}, r_{[11;12]}, r_{[11;11]}) = (2, 3, 2)$, we have to confirm the following equations:

$$\begin{aligned}
(5.3)\qquad & (m_{11} - m_{22} + 1)c^{\lambda}_{[11;12;0]}(M) + (m_{33} - m_{22} + 1)c^{\lambda}_{[11;11;0]}(M)\\
&= (-2m_{11} + 2m_{13})c^{\lambda}_{[11;22;0]}(M),
\end{aligned}$$

$$\begin{aligned}
(5.4)\qquad & (m_{12} - m_{23})\chi_-^{(-1)}(M)c^{\lambda}_{[11;12;0]}(M) + (m_{11} - m_{22})c^{\lambda}_{[11;12;1]}(M)\\
&+ C_1(M)c^{\lambda}_{[11;11;0]}(M) + (m_{33} - m_{22})c^{\lambda}_{[11;11;1]}(M)\\
&= (-2m_{11} + 2m_{13})c^{\lambda}_{[11;22;1]}(M),
\end{aligned}$$

$$\begin{aligned}
(5.5)\qquad & (m_{12} - m_{23} - 1)\chi_-^{(-1)}(M)c^{\lambda}_{[11;12;1]}(M) + (m_{11} - m_{22} - 1)c^{\lambda}_{[11;12;2]}(M)\\
&+ (C_1(M) - 1)\, c^{\lambda}_{[11;11;1]}(M) + (m_{33} - m_{22} - 1)c^{\lambda}_{[11;11;2]}(M)\\
&= (-2m_{11} + 2m_{13})c^{\lambda}_{[11;22;2]}(M),
\end{aligned}$$

$$\begin{aligned}
(5.6)\qquad & (m_{12} - m_{23} - 2)\chi_-^{(-1)}(M)c^{\lambda}_{[11;12;2]}(M) + (m_{11} - m_{22} - 2)c^{\lambda}_{[11;12;3]}(M)\\
&+ (C_1(M) - 2)\, c^{\lambda}_{[11;11;2]}(M) = 0,
\end{aligned}$$

$$(5.7)\qquad (m_{12} - m_{23} - 3)\chi_-^{(-1)}(M)c^{\lambda}_{[11;12;3]}(M) = 0.$$

We have

$$\begin{aligned}
&(m_{11}-m_{22}+1)c^{\lambda}_{[11;12;0]}(M)+(m_{33}-m_{22}+1)c^{\lambda}_{[11;11;0]}(M)\\
&=-2(m_{11}-m_{22}+1)(m_{13}-m_{12})(m_{13}-m_{12}-1)(m_{22}-m_{33})(m_{22}-m_{33}-1)\\
&\quad-2(m_{33}-m_{22}+1)(m_{13}-m_{12})(m_{13}-m_{12}-1)(m_{22}-m_{33})(m_{13}-m_{22}+1)\\
&=-2(m_{11}-m_{13})(m_{13}-m_{12})(m_{13}-m_{12}-1)(m_{22}-m_{33})(m_{22}-m_{33}-1)\\
&=(-2m_{11}+2m_{13})c^{\lambda}_{[11;22;0]}(M)\,.
\end{aligned}$$

Hence we obtain the equation (5.3).

We have

$$\begin{aligned}
&(m_{12}-m_{23})\chi_-^{(-1)}(M)c^{\lambda}_{[11;12;0]}(M)+(m_{11}-m_{22})c^{\lambda}_{[11;12;1]}(M)\\
&=-2(m_{12}-m_{23})\chi_-^{(-1)}(M)(m_{13}-m_{12})(m_{13}-m_{12}-1)(m_{22}-m_{33})(m_{22}-m_{33}-1)\\
&\quad+2(m_{11}-m_{22})(m_{13}-m_{12})(m_{22}-m_{33})\left\{\bar{F}\left(M\left(\begin{smallmatrix}-1&0&0\\&0&-1\\&&-1\end{smallmatrix}\right)\right)+\bar{E}(M)\right\}\\
&=2(m_{13}-m_{12})(m_{22}-m_{33})\Big\{-(m_{12}-m_{23})(m_{13}-m_{12}-1)(m_{22}-m_{33}-1)(1-\chi_+(M))\\
&\quad+(m_{11}-m_{22})\big\{-C_1(M)(\bar{C}_1(M)+1)\\
&\quad-\chi_+(M)\{(m_{13}-m_{12}-1)(m_{22}-m_{33}-1)+(m_{13}-m_{33})\delta(M)\}+\bar{E}(M)\big\}\Big\}\\
&=2(m_{13}-m_{12})(m_{22}-m_{33})\Big\{-(m_{12}-m_{23}-\delta(M)\chi_+(M))(m_{13}-m_{12}-1)(m_{22}-m_{33}-1)\\
&\quad-(m_{11}-m_{22})C_1(M)(\bar{C}_1(M)+1)-(m_{11}-m_{22})\chi_+(M)(m_{13}-m_{33})\delta(M)\\
&\quad+(m_{11}-m_{22})\bar{E}(M)\Big\}\\
&=2(m_{13}-m_{12})(m_{22}-m_{33})\Big\{-C_1(M)(m_{13}-m_{12}-1)(m_{22}-m_{33}-1)\\
&\quad-(m_{11}-m_{22})C_1(M)(\bar{C}_1(M)+1)-C_1(M)\chi_+(M)(m_{13}-m_{33})\delta(M)+(m_{11}-m_{22})\bar{E}(M)\Big\}\\
&=2(m_{13}-m_{12})(m_{22}-m_{33})\Big\{-C_1(M)\big\{(m_{13}-m_{12}-1)(m_{22}-m_{33}-1)\\
&\quad+(m_{13}-m_{33})\delta(M)\chi_+(M)+(m_{11}-m_{22})(\bar{C}_1(M)+1)\big\}+(m_{11}-m_{22})\bar{E}(M)\Big\},
\end{aligned}$$

$$\begin{aligned}
&C_1(M)c^{\lambda}_{[11;11;0]}(M)+(m_{33}-m_{22})c^{\lambda}_{[11;11;1]}(M)\\
&=-2C_1(M)(m_{13}-m_{12})(m_{13}-m_{12}-1)(m_{22}-m_{33})(m_{13}-m_{22}+1)\\
&\quad+2(m_{33}-m_{22})(m_{13}-m_{12})\Big\{(m_{22}-m_{33})C_1(M)(\bar{C}_1(M)+1)+(m_{13}-m_{22})\bar{E}(M)\Big\}\\
&=2(m_{13}-m_{12})(m_{22}-m_{33})\Big\{-C_1(M)\big\{(m_{13}-m_{12}-1)(m_{13}-m_{22}+1)\\
&\quad+(m_{22}-m_{33})(\bar{C}_1(M)+1)\big\}-(m_{13}-m_{22})\bar{E}(M)\Big\}.
\end{aligned}$$

Therefore

$$\begin{aligned}
&(m_{12}-m_{23})\chi_-^{(-1)}(M)c^{\lambda}_{[11;12;0]}(M)+(m_{11}-m_{22})c^{\lambda}_{[11;12;1]}(M)\\
&\quad+C_1(M)c^{\lambda}_{[11;11;0]}(M)+(m_{33}-m_{22})c^{\lambda}_{[11;11;1]}(M)\\
&\quad=2(m_{13}-m_{12})(m_{22}-m_{33})\Big\{-C_1(M)\big\{(m_{13}-m_{12}-1)(m_{13}-m_{33})
\end{aligned}$$

$$
\begin{aligned}
&\quad + (m_{13}-m_{33})\delta(M)\chi_+(M) + (m_{11}-m_{33})(\bar{C}_1(M)+1)\big\} + (m_{11}-m_{13})\bar{E}(M)\Big\} \\
&= 2(m_{13}-m_{12})(m_{22}-m_{33})\Big\{ - C_1(M)\big\{(m_{13}-m_{11})(m_{13}-m_{33}) \\
&\quad - (m_{13}-m_{33})(m_{12}-m_{11}-\delta(M)\chi_+(M)+1) + (m_{11}-m_{33})(\bar{C}_1(M)+1)\big\} \\
&\quad + (m_{11}-m_{13})\bar{E}(M)\Big\} \\
&= 2(m_{13}-m_{12})(m_{22}-m_{33})\Big\{(m_{11}-m_{13})C_1(M)(m_{13}-m_{33}-\bar{C}_1(M)-1) \\
&\quad + (m_{11}-m_{13})\bar{E}(M)\Big\} \\
&= 4(m_{11}-m_{13})(m_{13}-m_{12})(m_{22}-m_{33})(\bar{E}(M)-C_1(M)) \\
&= (-2m_{11}+2m_{13})c^{\lambda}_{[11;22;1]}(M)\,.
\end{aligned}
$$

Hence we obtain the equation (5.4). Here we use the relations

$$
\begin{aligned}
(5.8)\quad \bar{F}\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & -1 \\ & & -1 \end{smallmatrix}\right)\right) &= -C_1(M)(\bar{C}_1(M)+1) \\
&\quad - \chi_+(M)\{(m_{13}-m_{12}-1)(m_{22}-m_{33}-1) + (m_{13}-m_{33})\delta(M)\},
\end{aligned}
$$

(4.2) and (4.9).

We have

$$
\begin{aligned}
&(m_{12}-m_{23}-1)\chi_-^{(-1)}(M)c^{\lambda}_{[11;12;1]}(M) + (m_{11}-m_{22}-1)c^{\lambda}_{[11;12;2]}(M) \\
&= 2(m_{12}-m_{23}-1)\chi_-^{(-1)}(M)(m_{13}-m_{12})(m_{22}-m_{33})\Big\{\bar{F}\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & 0 & -1 \\ & & -1 \end{smallmatrix}\right)\right) + \bar{E}(M)\Big\} \\
&\quad - 2(m_{11}-m_{22}-1)\Big\{\bar{E}(M)\bar{F}\left(M\left(\begin{smallmatrix} -1 & 0 & 0 \\ & -1 & 0 \\ & & -1 \end{smallmatrix}\right)\right) \\
&\quad + (m_{13}-m_{12})(m_{22}-m_{33})C_1(M)(\bar{C}_1(M)+1)\chi_+(M)\Big\} \\
&= 2(m_{12}-m_{23}-1)\chi_-^{(-1)}(M)(m_{13}-m_{12})(m_{22}-m_{33})\big\{ - C_1(M)(\bar{C}_1(M)+1) + \bar{E}(M)\big\} \\
&\quad - 2(m_{11}-m_{22}-1)\Big\{\bar{E}(M)\big\{ - (C_1(M)-1)\bar{C}_1(M) - \chi_+(M)\{(m_{13}-m_{12})(m_{22}-m_{33}) \\
&\quad + (m_{13}-m_{33})\delta(M)\}\big\} + (m_{13}-m_{12})(m_{22}-m_{33})C_1(M)(\bar{C}_1(M)+1)\chi_+(M)\Big\} \\
&= 2(C_1(M)-1)\chi_-^{(-1)}(M)(m_{13}-m_{12})(m_{22}-m_{33})\big\{ - C_1(M)(\bar{C}_1(M)+1) + \bar{E}(M)\big\} \\
&\quad - 2\bar{E}(M)\big\{ - (C_1(M)-1)\bar{C}_1(M)(m_{11}-m_{22}-1) \\
&\quad - (C_1(M)-1)\chi_+(M)\{(m_{13}-m_{12})(m_{22}-m_{33}) + (m_{13}-m_{33})\delta(M)\}\big\} \\
&\quad - 2(C_1(M)-1)(m_{13}-m_{12})(m_{22}-m_{33})C_1(M)(\bar{C}_1(M)+1)\chi_+(M) \\
&= -2(C_1(M)-1)\Big\{(m_{13}-m_{12})(m_{22}-m_{33})C_1(M)(\bar{C}_1(M)+1)(\chi_-^{(-1)}(M)+\chi_+(M)) \\
&\quad - \bar{E}(M)\big\{\bar{C}_1(M)(m_{11}-m_{22}-1) + (m_{13}-m_{12})(m_{22}-m_{33})(\chi_-^{(-1)}(M)+\chi_+(M)) \\
&\quad + (m_{13}-m_{33})\delta(M)\chi_+(M)\big\}\Big\} \\
&= -2(C_1(M)-1)\Big\{(m_{13}-m_{12})(m_{22}-m_{33})C_1(M)(\bar{C}_1(M)+1) \\
&\quad - \bar{E}(M)\big\{\bar{C}_1(M)(m_{11}-m_{22}-1) + (m_{13}-m_{12})(m_{22}-m_{33}) \\
&\quad + (m_{13}-m_{33})\delta(M)\chi_+(M)\big\}\Big\},
\end{aligned}
$$

$$
(C_1(M)-1)c^{\lambda}_{[11;11;1]}(M) + (m_{33}-m_{22}-1)c^{\lambda}_{[11;11;2]}(M)
$$

$$\begin{aligned}
&= 2(C_1(M)-1)(m_{13}-m_{12})\{(m_{22}-m_{33})C_1(M)(\bar{C}_1(M)+1)+(m_{13}-m_{22})\bar{E}(M)\}\\
&\quad -2(m_{33}-m_{22}-1)\bar{E}(M)(C_1(M)-1)\bar{C}_1(M)\\
&= -2(C_1(M)-1)\Big\{-(m_{13}-m_{12})(m_{22}-m_{33})C_1(M)(\bar{C}_1(M)+1)\\
&\quad -\bar{E}(M)\{(m_{13}-m_{12})(m_{13}-m_{22})+(m_{22}-m_{33}+1)\bar{C}_1(M)\Big\}.
\end{aligned}$$

Therefore

$$\begin{aligned}
&(m_{12}-m_{23}-1)\chi_-^{(-1)}(M)c^{\lambda}_{[11;12;1]}(M)+(m_{11}-m_{22}-1)c^{\lambda}_{[11;12;2]}(M)\\
&+(C_1(M)-1)\,c^{\lambda}_{[11;11;1]}(M)+(m_{33}-m_{22}-1)c^{\lambda}_{[11;11;2]}(M)\\
&=2(C_1(M)-1)\bar{E}(M)\big\{\bar{C}_1(M)(m_{11}-m_{33})+(m_{13}-m_{12})(m_{13}-m_{33})\\
&\quad +(m_{13}-m_{33})\delta(M)\chi_+(M)\big\}\\
&=2(C_1(M)-1)\bar{E}(M)\big\{\bar{C}_1(M)(m_{11}-m_{33})+(m_{13}-m_{11})(m_{13}-m_{33})\\
&\quad -(m_{13}-m_{33})(m_{12}-m_{11}-\delta(M)\chi_+(M))\big\}\\
&=2(m_{13}-m_{11})\bar{E}(M)(C_1(M)-1)(m_{13}-m_{33}-\bar{C}_1(M))\\
&=(-2m_{11}+2m_{13})c^{\lambda}_{[11;22;2]}(M)\,.
\end{aligned}$$

Hence we obtain the equation (5.5). Here we use the relations

$$\begin{aligned}
\bar{F}\left(M\left(\begin{smallmatrix}-1&0&0\\&-1&0\\&&-1\end{smallmatrix}\right)\right) &= -(C_1(M)-1)\bar{C}_1(M)\\
&\quad -\chi_+(M)\{(m_{13}-m_{12})(m_{22}-m_{33})+(m_{13}-m_{33})\delta(M)\},
\end{aligned}\tag{5.9}$$

(4.17), (4.13), (4.14), (4.10) and (5.8).

We have

$$\begin{aligned}
&(m_{12}-m_{23}-2)\chi_-^{(-1)}(M)c^{\lambda}_{[11;12;2]}(M)+(m_{11}-m_{22}-2)c^{\lambda}_{[11;12;3]}(M)\\
&+(C_1(M)-2)\,c^{\lambda}_{[11;11;2]}(M)\\
&=-2(m_{12}-m_{23}-2)\chi_-^{(-1)}(M)\Big\{\bar{E}(M)\bar{F}\left(M\left(\begin{smallmatrix}-1&0&0\\&-1&0\\&&-1\end{smallmatrix}\right)\right)\\
&\quad +(m_{13}-m_{12})(m_{22}-m_{33})C_1(M)(\bar{C}_1(M)+1)\chi_+(M)\Big\}\\
&\quad +2(m_{11}-m_{22}-2)(C_1(M)-1)\bar{C}_1(M)\bar{E}(M)\chi_+(M)\\
&\quad -2(C_1(M)-2)\bar{E}(M)(C_1(M)-1)\bar{C}_1(M)\\
&=2(C_1(M)-2)\chi_-^{(-1)}(M)\bar{E}(M)(C_1(M)-1)\bar{C}_1(M)\\
&\quad +2(C_1(M)-2)(C_1(M)-1)\bar{C}_1(M)\bar{E}(M)\chi_+(M)\\
&\quad -2(C_1(M)-2)\bar{E}(M)(C_1(M)-1)\bar{C}_1(M)\\
&=2(C_1(M)-2)(C_1(M)-1)\bar{C}_1(M)\bar{E}(M)(\chi_-^{(-1)}(M)+\chi_+(M)-1)=0.
\end{aligned}$$

Hence we obtain the equation (5.6). Here we obtain the relations (4.9), (4.10), (4.13), (4.14) and (5.9).

We have

$$\begin{aligned}
&(m_{12}-m_{23}-3)\chi_-^{(-1)}(M)c^{\lambda}_{[11;12;3]}(M)\\
&=2(m_{12}-m_{23}-3)(C_1(M)-1)\bar{C}_1(M)\bar{E}(M)\chi_-^{(-1)}(M)\chi_+(M)=0.
\end{aligned}$$

Hence we obtain the equation (5.7). Here we use the relation (4.10).

- the proof of the case of the $(i,j)=(2,2)$.

Since $(r_{[22;22]}, r_{[22;12]}, r_{[22;11]}) = (2,2,2)$, we have to confirm the following equations:

(5.10)
$$\begin{aligned}&(m_{11}-m_{22}+1)c^{\lambda}_{[22;12;0]}(M)+(m_{33}-m_{22}+1)c^{\lambda}_{[22;11;0]}(M)\\&=(-2m_{11}+2m_{23}-2)c^{\lambda}_{[22;22;0]}(M),\end{aligned}$$

(5.11)
$$\begin{aligned}&(m_{12}-m_{23}+2)\chi_-^{(1)}(M)c^{\lambda}_{[22;12;0]}(M)+(m_{11}-m_{22})c^{\lambda}_{[22;12;1]}(M)\\&+\left(C_1(M)+\chi_-^{(1)}(M)+\chi_-(M)\right)c^{\lambda}_{[22;11;0]}(M)+(m_{33}-m_{22})c^{\lambda}_{[22;11;1]}(M)\\&=(-2m_{11}+2m_{23}-2)c^{\lambda}_{[22;22;1]}(M),\end{aligned}$$

(5.12)
$$\begin{aligned}&(m_{12}-m_{23}+1)\chi_-^{(1)}(M)c^{\lambda}_{[22;12;1]}(M)+(m_{11}-m_{22}-1)c^{\lambda}_{[22;12;2]}(M)\\&+\left(C_1(M)-1+\chi_-^{(1)}(M)+\chi_-(M)\right)c^{\lambda}_{[22;11;1]}(M)+(m_{33}-m_{22}-1)c^{\lambda}_{[22;11;2]}(M)\\&=(-2m_{11}+2m_{23}-2)c^{\lambda}_{[22;22;2]}(M),\end{aligned}$$

(5.13)
$$(m_{12}-m_{23})\chi_-^{(1)}(M)c^{\lambda}_{[22;12;2]}(M)+\left(C_1(M)-2+\chi_-^{(1)}(M)+\chi_-(M)\right)c^{\lambda}_{[22;11;2]}(M)=0.$$

We have

$$\begin{aligned}&(m_{11}-m_{22}+1)c^{\lambda}_{[22;12;0]}(M)+(m_{33}-m_{22}+1)c^{\lambda}_{[22;11;0]}(M)\\&=-2(m_{11}-m_{22}+1)(m_{22}-m_{33})(m_{22}-m_{33}-1)\\&\quad-2(m_{33}-m_{22}+1)(m_{22}-m_{33})(m_{23}-m_{22})\\&=-2(m_{11}-m_{23}+1)(m_{22}-m_{33})(m_{22}-m_{33}-1)\\&=(-2m_{11}+2m_{23}-2)c^{\lambda}_{[22;22;0]}(M).\end{aligned}$$

Hence we obtain the equation (5.10).

We have

$$\begin{aligned}&(m_{12}-m_{23}+2)\chi_-^{(1)}(M)c^{\lambda}_{[22;12;0]}(M)+(m_{11}-m_{22})c^{\lambda}_{[22;12;1]}(M)\\&+\left(C_1(M)+\chi_-^{(1)}(M)+\chi_-(M)\right)c^{\lambda}_{[22;11;0]}(M)+(m_{33}-m_{22})c^{\lambda}_{[22;11;1]}(M)\\&=-2(m_{12}-m_{23}+2)\chi_-^{(1)}(M)(m_{22}-m_{33})(m_{22}-m_{33}-1)\\&\quad+2(m_{11}-m_{22})(m_{22}-m_{33})\{\bar{C}_1(M)+(\bar{D}(M)+1)\chi_-(M)\}\\&\quad-2\left(C_1(M)+\chi_-^{(1)}(M)+\chi_-(M)\right)(m_{22}-m_{33})(m_{23}-m_{22})\\&\quad+2(m_{33}-m_{22})\{\bar{D}(M)(m_{23}-m_{22}-1)\chi_-(M)-(m_{22}-m_{33})(\bar{C}_1(M)+1)\chi_-^{(1)}(M)\}\\&=2(m_{22}-m_{33})\Big\{-(m_{12}-m_{23}+2)(m_{22}-m_{33}-1)\chi_-^{(1)}(M)\\&\quad+(m_{11}-m_{22})\{m_{23}-m_{22}+(\delta(M)+\bar{D}(M)+1)\chi_-(M)\}\\&\quad-\{m_{11}-m_{22}+(\delta(M)+1)\chi_-(M)+\chi_-^{(1)}(M)\}(m_{23}-m_{22})\\&\quad-\{\bar{D}(M)(m_{23}-m_{22}-1)\chi_-(M)-(m_{22}-m_{33})(m_{12}-m_{11}+1)\chi_-^{(1)}(M)\}\Big\}\end{aligned}$$

$$
\begin{aligned}
&= 2(m_{22}-m_{33})\Big\{ -(m_{11}-m_{23}+1)(m_{22}-m_{33}-1)\chi_-^{(1)}(M) \\
&\quad + (\delta(M)+1)\{(m_{11}-m_{23})\chi_-(M)+\chi_-^{(1)}(M)\} + (m_{11}-m_{23}+1)\bar{D}(M)\chi_-(M)\Big\} \\
&= 2(m_{22}-m_{33})\Big\{(m_{11}-m_{23}+1)(-m_{22}+m_{33}+\delta(M)+2)\chi_-^{(1)}(M) \\
&\quad + (m_{11}-m_{23}+1)\bar{D}(M)\chi_-(M)\Big\} \\
&= 2(m_{11}-m_{23}+1)(m_{22}-m_{33})\{(\bar{D}(M)+2)\chi_-^{(1)}(M)+\bar{D}(M)\chi_-(M)\} \\
&= (-2m_{11}+2m_{23}-2)c^{\lambda}_{[22;22;1]}(M)\,.
\end{aligned}
$$

Hence we obtain the equation (5.11). Here we use the relations (4.1), (4.2), (4.8) and (4.16).

We have

$$
\begin{aligned}
&\big(C_1(M)-1+\chi_-^{(1)}(M)+\chi_-(M)\big)\, c^{\lambda}_{[22;11;1]}(M) \\
&= 2\left(C_1(M)-1+\chi_-^{(1)}(M)+\chi_-(M)\right)\{\bar{D}(M)(m_{23}-m_{22}-1)\chi_-(M) \\
&\quad -(m_{22}-m_{33})(\bar{C}_1(M)+1)\chi_-^{(1)}(M)\} \\
&= 2C_1(M)\bar{D}(M)(m_{23}-m_{22}-1)\chi_-(M) \\
&\quad -2\{(C_1(M)+1)\,(m_{22}-m_{33})(\bar{C}_1(M)+1)-\bar{D}(M)(m_{23}-m_{22}-1)\}\chi_-^{(1)}(M).
\end{aligned}
$$

Therefore

$$
\begin{aligned}
&(m_{12}-m_{23}+1)\chi_-^{(1)}(M)c^{\lambda}_{[22;12;1]}(M)+(m_{11}-m_{22}-1)c^{\lambda}_{[22;12;2]}(M) \\
&+\big(C_1(M)-1+\chi_-^{(1)}(M)+\chi_-(M)\big)\, c^{\lambda}_{[22;11;1]}(M)+(m_{33}-m_{22}-1)c^{\lambda}_{[22;11;2]}(M) \\
&= 2(m_{12}-m_{23}+1)\chi_-^{(1)}(M)(m_{22}-m_{33})\{\bar{C}_1(M)+(\bar{D}(M)+1)\chi_-(M)\} \\
&\quad -2(m_{11}-m_{22}-1)\bar{C}_1(M)\bar{D}(M)\chi_-(M) \\
&\quad +2C_1(M)\bar{D}(M)(m_{23}-m_{22}-1)\chi_-(M) \\
&\quad -2\{(C_1(M)+1)\,(m_{22}-m_{33})(\bar{C}_1(M)+1)-\bar{D}(M)(m_{23}-m_{22}-1)\}\chi_-^{(1)}(M) \\
&\quad +2(m_{33}-m_{22}-1)\bar{C}_1(M)\bar{D}(M)\chi_-^{(1)}(M) \\
&= 2(m_{12}-m_{23}+1)(m_{22}-m_{33})(m_{12}-m_{11}+\bar{D}(M)+1)\chi_-^{(1)}(M) \\
&\quad -2(m_{11}-m_{22}-1)(m_{12}-m_{11})\bar{D}(M)\chi_-(M) \\
&\quad +2(m_{12}-m_{23})\bar{D}(M)(m_{23}-m_{22}-1)\chi_-(M) \\
&\quad -2\{(m_{12}-m_{23}+1)(m_{22}-m_{33})(m_{12}-m_{11}+1)-\bar{D}(M)(m_{23}-m_{22}-1)\}\chi_-^{(1)}(M) \\
&\quad +2(m_{33}-m_{22}-1)(m_{12}-m_{11})\bar{D}(M)\chi_-^{(1)}(M) \\
&= 2(m_{11}-m_{23})(m_{22}-m_{33})\bar{D}(M)\chi_-^{(1)}(M)-2(-m_{22}+m_{33}+\delta(M)+1)\bar{D}(M)\chi_-^{(1)}(M) \\
&\quad -2(m_{11}-m_{23})(\delta(M)+1)\bar{D}(M)\chi_-(M) \\
&= -2(m_{11}-m_{23}+1)(-m_{22}+m_{33}+\delta(M)+1)\bar{D}(M)\chi_-^{(1)}(M) \\
&= (-2m_{11}+2m_{23}-2)c^{\lambda}_{[22;22;2]}(M)\,.
\end{aligned}
$$

Hence we obtain the equation (5.12). Here we use the relations (4.1), (4.2), (4.8), (4.12), (4.16) and (4.14).

We have

$$
(m_{12}-m_{23})\chi_-^{(1)}(M)c^{\lambda}_{[22;12;2]}(M)+\big(C_1(M)-2+\chi_-^{(1)}(M)+\chi_-(M)\big)\, c^{\lambda}_{[22;11;2]}(M)
$$

$$\begin{aligned}&= -2(m_{12}-m_{23})\chi_-^{(1)}(M)\bar{C}_1(M)\bar{D}(M)\chi_-(M)\\&\quad + 2\left(C_1(M)-2+\chi_-^{(1)}(M)+\chi_-(M)\right)\bar{C}_1(M)\bar{D}(M)\chi_-^{(1)}(M)\\&= -2C_1(M)\bar{C}_1(M)\bar{D}(M)\chi_-^{(1)}(M)+2C_1(M)\bar{C}_1(M)\bar{D}(M)\chi_-^{(1)}(M)=0.\end{aligned}$$

Hence we obtain the equation (5.13). Here we use the relations (4.14) and (4.12).

- the proof of the case of the $(i,j)=(3,3)$.

Since $(r_{[33;22]},r_{[33;12]},r_{[33;11]})=(0,1,0)$, we have to confirm the following equations:

$$\begin{aligned}&(m_{11}-m_{22}+1)c^{\lambda}_{[33;12;0]}(M)+(m_{33}-m_{22}-1)c^{\lambda}_{[33;11;0]}(M)\\&\quad=(-2m_{11}+2m_{33}-4)c^{\lambda}_{[33;22;0]}(M),\end{aligned}\tag{5.14}$$

$$\begin{aligned}&(m_{12}-m_{23})\chi_-^{(-1)}(M)c^{\lambda}_{[33;12;0]}(M)+(m_{11}-m_{22})c^{\lambda}_{[33;12;1]}(M)+C_1(M)c^{\lambda}_{[33;11;0]}(M)\\&\quad=(-2m_{11}+2m_{33}-4)c^{\lambda}_{[33;22;1]}(M),\end{aligned}\tag{5.15}$$

$$(m_{12}-m_{23}-1)\chi_-^{(-1)}(M)c^{\lambda}_{[33;12;1]}(M)=0.\tag{5.16}$$

We have

$$\begin{aligned}&(m_{11}-m_{22}+1)c^{\lambda}_{[33;12;0]}(M)+(m_{33}-m_{22}-1)c^{\lambda}_{[33;11;0]}(M)\\&=-2(m_{11}-m_{22}+1)+2(m_{33}-m_{22}-1)\\&=-2m_{11}+2m_{33}-4=(-2m_{11}+2m_{33}-4)c^{\lambda}_{[33;22;0]}(M).\end{aligned}$$

Hence we obtain the equation (5.14).

We have

$$\begin{aligned}&(m_{12}-m_{23})\chi_-^{(-1)}(M)c^{\lambda}_{[33;12;0]}(M)+(m_{11}-m_{22})c^{\lambda}_{[33;12;1]}(M)+C_1(M)c^{\lambda}_{[33;11;0]}(M)\\&=-2(m_{12}-m_{23})\chi_-^{(-1)}(M)-2(m_{11}-m_{22})\chi_+(M)+2C_1(M)\\&=-2(m_{12}-m_{23})(1-\chi_+(M))-2(m_{11}-m_{22})\chi_+(M)+2C_1(M)\\&=-2(m_{12}-m_{23}-\delta(M)\chi_+(M)-C_1(M))=0.\end{aligned}$$

Hence we obtain the equation (5.15). Here we use the relations (4.1) and (4.9).

We have

$$(m_{12}-m_{23}-1)\chi_-^{(-1)}(M)c^{\lambda}_{[33;12;1]}(M)=-2(m_{12}-m_{23}-1)\chi_-^{(-1)}(M)\chi_+(M)=0.$$

Hence we obtain the equation (5.16). Here we use the relation (4.10).

- the proof of the case of the $(i,j)=(1,2)$.

Since $(r_{[12;22]},r_{[12;12]},r_{[12;11]})=(2,2,2)$, we have to confirm the following equations:

$$\begin{aligned}&(m_{11}-m_{22}+1)c^{\lambda}_{[12;12;0]}(M)+(m_{33}-m_{22}+1)c^{\lambda}_{[12;11;0]}(M)\\&\quad=(-2m_{11}+m_{13}+m_{23})c^{\lambda}_{[12;22;0]}(M),\end{aligned}\tag{5.17}$$

$$\begin{aligned}&(m_{12}-m_{23}+1)\chi_-(M)c^{\lambda}_{[12;12;0]}(M)+(m_{11}-m_{22})c^{\lambda}_{[12;12;1]}(M)\\&\quad+\left(C_1(M)+\chi_-(M)\right)c^{\lambda}_{[12;11;0]}(M)+(m_{33}-m_{22})c^{\lambda}_{[12;11;1]}(M)\\&\quad=(-2m_{11}+m_{13}+m_{23})c^{\lambda}_{[12;22;1]}(M),\end{aligned}\tag{5.18}$$

$$\begin{aligned}
&(m_{12}-m_{23})\chi_-(M)c^\lambda_{[12;12;1]}(M)+(m_{11}-m_{22}-1)c^\lambda_{[12;12;2]}(M)\\
&+\left(C_1(M)-1+\chi_-(M)\right)c^\lambda_{[12;11;1]}(M)+(m_{33}-m_{22}-1)c^\lambda_{[12;11;2]}(M)\\
&=(-2m_{11}+m_{13}+m_{23})c^\lambda_{[12;22;2]}(M)\,,
\end{aligned}\tag{5.19}$$

$$(m_{12}-m_{23}-1)\chi_-(M)c^\lambda_{[12;12;2]}(M)+\left(C_1(M)-2+\chi_-(M)\right)c^\lambda_{[12;11;2]}(M)=0.\tag{5.20}$$

We have

$$\begin{aligned}
&(m_{11}-m_{22}+1)c^\lambda_{[12;12;0]}(M)+(m_{33}-m_{22}+1)c^\lambda_{[12;11;0]}(M)\\
&=-2(m_{11}-m_{22}+1)(m_{13}-m_{12})(m_{22}-m_{33})(m_{22}-m_{33}-1)\\
&\quad+(m_{33}-m_{22}+1)(m_{13}-m_{12})(m_{22}-m_{33})(2m_{22}-m_{13}-m_{23}-2)\\
&=-(2m_{11}-m_{13}-m_{23})(m_{13}-m_{12})(m_{22}-m_{33})(m_{22}-m_{33}-1)\\
&=(-2m_{11}+m_{13}+m_{23})c^\lambda_{[12;22;0]}(M)\,.
\end{aligned}$$

Hence we obtain the equation (5.17).

We have

$$\begin{aligned}
&(m_{11}-m_{22})c^\lambda_{[12;12;1]}(M)+(m_{33}-m_{22})c^\lambda_{[12;11;1]}(M)\\
&=(m_{11}-m_{22})(m_{22}-m_{33})\big\{\bar{E}(M)\\
&\quad-C_2(M)-\chi_+(M)\{(m_{13}-m_{12})(m_{22}-m_{33})+(m_{13}-m_{33}+1)\delta(M)\}\\
&\quad+(m_{13}-m_{12})\{\bar{C}_1(M)+1+\bar{D}(M)(1-\chi_+(M))\}\big\}\\
&\quad+(m_{33}-m_{22})\Big\{\bar{E}(M)(m_{23}-m_{22})+C_2(M)(m_{22}-m_{33}+1)\\
&\quad+(m_{13}-m_{12})\chi_-(M)\{\bar{D}(M)(m_{13}-m_{22}+1)-(m_{22}-m_{33})(\bar{C}_1(M)+1)\}\Big\}\\
&=(m_{22}-m_{33})\Big\{(m_{11}-m_{23})\bar{E}(M)\\
&\quad-(m_{11}-m_{33}+1)C_2(M)-(m_{11}-m_{22})(m_{13}-m_{33}+1)\delta(M)\chi_+(M)\\
&\quad+(m_{13}-m_{12})\big\{(m_{11}-m_{22})\{\bar{C}_1(M)+1+\bar{D}(M)(1-\chi_+(M))-(m_{22}-m_{33})\chi_+(M)\}\\
&\quad-\chi_-(M)\{\bar{D}(M)(m_{13}-m_{22}+1)-(m_{22}-m_{33})(\bar{C}_1(M)+1)\}\big\}\Big\}\\
&=(m_{22}-m_{33})\Big\{(2m_{11}-m_{13}-m_{23})\bar{E}(M)+(m_{13}-m_{11})C_1(M)(m_{13}-m_{33}+1-\bar{C}_1(M))\\
&\quad-(m_{11}-m_{33}+1)C_2(M)-C_1(M)(m_{13}-m_{33}+1)\delta(M)\chi_+(M)\\
&\quad+(m_{13}-m_{12})\big\{(m_{11}-m_{22})\{\bar{C}_1(M)+1+\bar{D}(M)-(-m_{22}+m_{33}+\delta(M))\chi_+(M)\\
&\quad-(m_{22}-m_{33})\chi_+(M)\}-\chi_-(M)\{\bar{D}(M)(m_{13}-m_{22}+1)-(m_{22}-m_{33})(\bar{C}_1(M)+1)\}\big\}\Big\}\\
&=(m_{22}-m_{33})\Big\{(2m_{11}-m_{13}-m_{23})\bar{E}(M)\\
&\quad+(m_{13}-m_{11}-\bar{C}_1(M)-\delta(M)\chi_+(M))C_1(M)(m_{13}-m_{33}+1)\\
&\quad+(m_{13}-m_{12})\big\{(m_{11}-m_{22})\{\bar{C}_1(M)+1+\bar{D}(M)-\delta(M)\chi_+(M)\}\\
&\quad-\chi_-(M)\{\bar{D}(M)(m_{13}-m_{22}+1)-(m_{22}-m_{33})(\bar{C}_1(M)+1)\}\big\}\Big\}\\
&=(m_{22}-m_{33})\Big\{(2m_{11}-m_{13}-m_{23})\bar{E}(M)+(m_{13}-m_{12})\big\{C_1(M)(m_{13}-m_{33}+1)
\end{aligned}$$

$$\begin{aligned}
&+ (m_{11} - m_{22})\{\bar{C}_1(M) + 1 + \bar{D}(M) - \delta(M)\chi_+(M)\} \\
&- \chi_-(M)\{\bar{D}(M)(m_{13} - m_{22} + 1) - (m_{22} - m_{33})(\bar{C}_1(M) + 1)\}\}\Big\}.
\end{aligned}$$

Therefore

$$\begin{aligned}
&(m_{12} - m_{23} + 1)\chi_-(M)c^{\lambda}_{[12;12;0]}(M) + (m_{11} - m_{22})c^{\lambda}_{[12;12;1]}(M) \\
&+ (C_1(M) + \chi_-(M))\, c^{\lambda}_{[12;11;0]}(M) + (m_{33} - m_{22})c^{\lambda}_{[12;11;1]}(M) \\
&= -2(m_{12} - m_{23} + 1)\chi_-(M)(m_{13} - m_{12})(m_{22} - m_{33})(m_{22} - m_{33} - 1) \\
&\quad + (C_1(M) + \chi_-(M))\,(m_{13} - m_{12})(m_{22} - m_{33})(2m_{22} - m_{13} - m_{23} - 2) \\
&\quad + (m_{22} - m_{33})\Big\{(2m_{11} - m_{13} - m_{23})\bar{E}(M) + (m_{13} - m_{12})\{C_1(M)(m_{13} - m_{33} + 1) \\
&\quad + (m_{11} - m_{22})\{\bar{C}_1(M) + 1 + \bar{D}(M) - \delta(M)\chi_+(M)\} \\
&\quad - \chi_-(M)\{\bar{D}(M)(m_{13} - m_{22} + 1) - (m_{22} - m_{33})(\bar{C}_1(M) + 1)\}\}\Big\} \\
&= (m_{22} - m_{33})\Big\{(2m_{11} - m_{13} - m_{23})\bar{E}(M) \\
&\quad + (m_{13} - m_{12})\{ -2(m_{12} - m_{23} + 1)(m_{22} - m_{33} - 1)\chi_-(M) \\
&\quad + \{m_{11} - m_{22} + (\delta(M) + 1)\chi_-(M)\}(2m_{22} - m_{13} - m_{23} - 2) \\
&\quad + (m_{11} - m_{22} + \delta(M)\chi_-(M))(m_{13} - m_{33} + 1) + (m_{11} - m_{22}) \\
&\qquad \times \{m_{23} - m_{22} + \delta(M)\chi_-(M) + 1 + (-m_{22} + m_{33} + \delta(M)) - \delta(M)(1 - \chi_-(M))\} \\
&\quad - \chi_-(M)\{(-m_{22} + m_{33} + \delta(M))(m_{13} - m_{22} + 1) - (m_{22} - m_{33})(m_{12} - m_{11} + 1)\}\}\Big\} \\
&= (m_{22} - m_{33})\Big\{(2m_{11} - m_{13} - m_{23})\bar{E}(M) \\
&\quad + (2m_{11} - m_{13} - m_{23})(m_{13} - m_{12})\chi_-(M)(-m_{22} + m_{33} + \delta(M) + 1)\Big\} \\
&= (2m_{11} - m_{13} - m_{23})(m_{22} - m_{33})\{\bar{E}(M) + (m_{13} - m_{12})\chi_-(M)(\bar{D}(M) + 1)\} \\
&= (-2m_{11} + m_{13} + m_{23})c^{\lambda}_{[12;22;1]}(M)\,.
\end{aligned}$$

Hence we obtain the equation (5.18). Here we use the relations (4.1), (4.2) and (4.14).

We have

$$\begin{aligned}
&(m_{12} - m_{23})\chi_-(M)c^{\lambda}_{[12;12;1]}(M) \\
&= (m_{12} - m_{23})\chi_-(M)(m_{22} - m_{33})\{\bar{E}(M) + \bar{F}(M) \\
&\quad + (m_{13} - m_{12})\{\bar{C}_1(M) + 1 + \bar{D}(M)(1 - \chi_+(M))\}\} \\
&= (m_{22} - m_{33})\chi_-(M)\{(m_{12} - m_{23})(\bar{E}(M) - C_2(M)) \\
&\quad + (m_{13} - m_{12})C_1(M)(\bar{C}_1(M) + 1 + \bar{D}(M))\}
\end{aligned}$$

$$\begin{aligned}
&(m_{11} - m_{22} - 1)c^{\lambda}_{[12;12;2]}(M) \\
&= -(m_{11} - m_{22} - 1)\{\bar{C}_1(M)\bar{E}(M) + C_2(M)\left(1 - \bar{D}(M) + \delta(M)\chi_+(M)\right)\} \\
&= -(m_{11} - m_{22} - 1)\{\bar{C}_1(M)\bar{E}(M) + C_2(M)\,(m_{22} - m_{33} + 1 - \delta(M)\chi_-(M))\}
\end{aligned}$$

$$\begin{aligned}
&(C_1(M) - 1 + \chi_-(M))\, c^{\lambda}_{[12;11;1]}(M) \\
&= (C_1(M) - 1 + \chi_-(M))\,\{\bar{E}(M)(m_{23} - m_{22}) + C_2(M)(m_{22} - m_{33} + 1) \\
&\quad + (m_{13} - m_{12})\chi_-(M)\{\bar{D}(M)(m_{13} - m_{22} + 1) - (m_{22} - m_{33})(\bar{C}_1(M) + 1)\}\}
\end{aligned}$$

$$\begin{aligned}
&= (m_{11}-m_{22}-1)\bigl\{\bar{C}_1(M)\bar{E}(M) - \delta(M)\chi_-(M)\bar{E}(M) + C_2(M)(m_{22}-m_{33}+1)\bigr\}\\
&\quad + (\delta(M)+1)\chi_-(M)\bigl\{\bar{E}(M)(m_{23}-m_{22}) + C_2(M)(m_{22}-m_{33}+1)\bigr\}\\
&\quad + (m_{13}-m_{12})C_1(M)\chi_-(M)\{\bar{D}(M)(m_{13}-m_{22}+1) - (m_{22}-m_{33})(\bar{C}_1(M)+1)\}
\end{aligned}$$

$$\begin{aligned}
&(m_{33}-m_{22}-1)c^{\lambda}_{[12;11;2]}(M)\\
&= (m_{33}-m_{22}-1)C_2(M)\chi_-(M)(m_{13}-m_{33}+2-\bar{C}_1(M)-\bar{D}(M))\\
&= -(m_{22}-m_{33}+1)\chi_-(M)\{(m_{12}-m_{11})\bar{E}(M) + (1-\bar{D}(M))C_2(M)\}
\end{aligned}$$

Therefore

$$\begin{aligned}
&(m_{11}-m_{22}-1)c^{\lambda}_{[12;12;2]}(M) + (C_1(M)-1+\chi_-(M))\,c^{\lambda}_{[12;11;1]}(M)\\
&\quad + (m_{12}-m_{23})\chi_-(M)c^{\lambda}_{[12;12;1]}(M) + (m_{33}-m_{22}-1)c^{\lambda}_{[12;11;2]}(M)\\
&= \chi_-(M)\bigl\{-(m_{11}-m_{23})\bar{E}(M)\bar{D}(M) + (m_{11}-m_{33})\delta(M)C_2(M)\\
&\quad + (m_{13}-m_{33}+1)(m_{13}-m_{12})C_1(M)\bar{D}(M) - (m_{12}-m_{23})(m_{22}-m_{33})C_2(M)\\
&\quad + (m_{22}-m_{33}+1)\bar{D}(M)C_2(M)\bigr\}\\
&= \chi_-(M)\bigl\{-(2m_{11}-m_{13}-m_{23})\bar{E}(M)\bar{D}(M)\\
&\quad + (m_{11}-m_{13})(m_{12}-m_{23})(m_{13}-m_{33}+1-m_{12}+m_{11})\bar{D}(M)\\
&\quad + (m_{11}-m_{33})\delta(M)(m_{12}-m_{23})(m_{12}-m_{11})\\
&\quad + (m_{13}-m_{33}+1)(m_{13}-m_{12})(m_{12}-m_{23})\bar{D}(M)\\
&\quad - (m_{12}-m_{23})(m_{22}-m_{33})(m_{12}-m_{23})(m_{12}-m_{11})\\
&\quad + (m_{22}-m_{33}+1)(m_{12}-m_{23})(m_{12}-m_{11})\bar{D}(M)\bigr\}\\
&= \chi_-(M)\bigl\{-(2m_{11}-m_{13}-m_{23})\bar{E}(M)\bar{D}(M) + (m_{12}-m_{23})(m_{12}-m_{11})\{(m_{11}-m_{33})\delta(M)\\
&\quad - (m_{22}-m_{33})(m_{12}-m_{23}) + (m_{22}-m_{11})\bar{D}(M)\}\bigr\}\\
&= \chi_-(M)\bigl\{-(2m_{11}-m_{13}-m_{23})\bar{E}(M)\bar{D}(M) + (m_{12}-m_{23})(m_{12}-m_{11})\{(m_{11}-m_{33})\delta(M)\\
&\quad - (m_{22}-m_{33})(m_{12}-m_{23}) + (m_{22}-m_{11})(-m_{22}+m_{33}-\delta(M))\}\bigr\}\\
&= \chi_-(M) - (2m_{11}-m_{13}-m_{23})\bar{E}(M)\bar{D}(M)\\
&= (-2m_{11}+m_{13}+m_{23})c^{\lambda}_{[12;22;2]}(M)\,.
\end{aligned}$$

Hence we obtain the equation 5.19. Here we use the relations (4.10), (4.12), (4.9), (4.26), (4.16) and (4.14).

We have

$$\begin{aligned}
&(m_{12}-m_{23}-1)\chi_-(M)c^{\lambda}_{[12;12;2]}(M)\\
&= (m_{12}-m_{23}-1)\chi_-(M)\bigl\{-\bar{C}_1(M)\bar{E}(M) - C_2(M)\bigl(1-\bar{D}(M)+\delta(M)\chi_+(M)\bigr)\bigr\}\\
&= -(C_1(M)-1)\chi_-(M)(m_{13}-m_{33}+2-\bar{C}_1(M)-\bar{D}(M)),
\end{aligned}$$

$$\begin{aligned}
&(C_1(M)-2+\chi_-(M))\,c^{\lambda}_{[12;11;2]}(M)\\
&= (C_1(M)-2+\chi_-(M))\,C_2(M)\chi_-(M)(m_{13}-m_{33}+2-\bar{C}_1(M)-\bar{D}(M))\\
&= (C_1(M)-1)C_2(M)\chi_-(M)(m_{13}-m_{33}+2-\bar{C}_1(M)-\bar{D}(M)).
\end{aligned}$$

Therefore

$$(m_{12}-m_{23}-1)\chi_-(M)c^{\lambda}_{[12;12;2]}(M) + (C_1(M)-2+\chi_-(M))\,c^{\lambda}_{[12;11;2]}(M) = 0.$$

Hence we obtain the equation (5.20). Here we use the relations (4.10), (4.12) and (4.14).

- the proof of the case of the $(i, j) = (1, 3)$.

Since $(r_{[13;22]}, r_{[13;12]}, r_{[13;11]}) = (1, 2, 1)$, we have to confirm the following equations:

$$\begin{aligned}(m_{11} - m_{22} + 1)c^{\lambda}_{[13;12;0]}(M) &+ (m_{33} - m_{22})c^{\lambda}_{[13;11;0]}(M) \\ &= (-2m_{11} + m_{13} + m_{33} - 1)c^{\lambda}_{[13;22;0]}(M),\end{aligned} \tag{5.21}$$

$$\begin{aligned}(m_{12} - m_{23})\chi_-^{(-1)}(M)c^{\lambda}_{[13;12;0]}(M) &+ (m_{11} - m_{22})c^{\lambda}_{[13;12;1]}(M) \\ &+ C_1(M)c^{\lambda}_{[13;11;0]}(M) + (m_{33} - m_{22} - 1)c^{\lambda}_{[13;11;1]}(M) \\ &= (-2m_{11} + m_{13} + m_{33} - 1)c^{\lambda}_{[13;22;1]}(M),\end{aligned} \tag{5.22}$$

$$\begin{aligned}(m_{12} - m_{23} - 1)\chi_-^{(-1)}(M)c^{\lambda}_{[13;12;1]}(M) &+ (m_{11} - m_{22} - 1)c^{\lambda}_{[13;12;2]}(M) \\ &+ (C_1(M) - 1)\, c^{\lambda}_{[13;11;1]}(M) = 0,\end{aligned} \tag{5.23}$$

$$(m_{12} - m_{23} - 2)\chi_-^{(-1)}(M)c^{\lambda}_{[13;12;2]}(M) = 0. \tag{5.24}$$

We have

$$\begin{aligned}&(m_{11} - m_{22} + 1)c^{\lambda}_{[13;12;0]}(M) + (m_{33} - m_{22})c^{\lambda}_{[13;11;0]}(M) \\ &= -2(m_{11} - m_{22} + 1)(m_{13} - m_{12})(m_{22} - m_{33}) \\ &\quad + (m_{33} - m_{22})(m_{13} - m_{12})(2m_{22} - m_{13} - m_{23} - 1) \\ &= -(2m_{11} - m_{13} - m_{23} + 1)(m_{13} - m_{12})(m_{22} - m_{33}) \\ &= (-2m_{11} + m_{13} + m_{33} - 1)c^{\lambda}_{[13;22;0]}(M).\end{aligned}$$

Hence we obtain the equation (5.21).

We have

$$\begin{aligned}&(m_{12} - m_{23})\chi_-^{(-1)}(M)c^{\lambda}_{[13;12;0]}(M) + (m_{11} - m_{22})c^{\lambda}_{[13;12;1]}(M) \\ &= -2(m_{12} - m_{23})\chi_-^{(-1)}(M)(m_{13} - m_{12})(m_{22} - m_{33}) \\ &\quad + (m_{11} - m_{22})\{\bar{E}(M) + \bar{F}(M) - (m_{13} - m_{12})(m_{22} - m_{33})\chi_+(M)\} \\ &= -2(m_{12} - m_{23})(m_{13} - m_{12})(m_{22} - m_{33})(1 - \chi_+(M)) + (m_{11} - m_{22})\big\{\bar{E}(M) - C_2(M) \\ &\quad - \chi_+(M)\{2(m_{13} - m_{12})(m_{22} - m_{33}) + (m_{13} - m_{33} + 1)\delta(M)\}\big\} \\ &= -2(m_{13} - m_{12})(m_{22} - m_{33})(m_{12} - m_{23} - \delta(M)\chi_+(M)) \\ &\quad + (m_{11} - m_{22})(\bar{E}(M) - C_2(M)) - (m_{11} - m_{22})(m_{13} - m_{33} + 1)\delta(M)\chi_+(M) \\ &= -2(m_{13} - m_{12})(m_{22} - m_{33})C_1(M) \\ &\quad + (m_{11} - m_{22})(\bar{E}(M) - C_2(M)) - C_1(M)(m_{13} - m_{33} + 1)\delta(M)\chi_+(M) \\ &= -C_1(M)\{(m_{13} - m_{33} + 1)\delta(M)\chi_+(M) + 2(m_{13} - m_{12})(m_{22} - m_{33})\} \\ &\quad + (m_{11} - m_{22})(\bar{E}(M) - C_2(M)),\end{aligned}$$

$$\begin{aligned}&C_1(M)c^{\lambda}_{[13;11;0]}(M) + (m_{33} - m_{22} - 1)c^{\lambda}_{[13;11;1]}(M) \\ &= C_1(M)(m_{13} - m_{12})(2m_{22} - m_{13} - m_{33} - 1) + (m_{33} - m_{22} - 1)(C_2(M) - \bar{E}(M)).\end{aligned}$$

Therefore

$$(m_{12} - m_{23})\chi_-^{(-1)}(M)c^{\lambda}_{[13;12;0]}(M) + (m_{11} - m_{22})c^{\lambda}_{[13;12;1]}(M)$$

$$
\begin{aligned}
&+C_1(M)c^{\lambda}_{[13;11;0]}(M)+(m_{33}-m_{22}-1)c^{\lambda}_{[13;11;1]}(M)\\
&=C_1(M)\{-(m_{13}-m_{33}+1)\delta(M)\chi_+(M)+(m_{13}-m_{12})(m_{33}-m_{13}-1)\}\\
&\quad+(m_{33}-m_{11}-1)(C_2(M)-\bar{E}(M))\\
&=C_1(M)\{-(m_{13}-m_{33}+1)\delta(M)\chi_+(M)+(m_{13}-m_{12})(m_{33}-m_{13}-1)\\
&\quad+(m_{33}-m_{11}-1)\bar{C}_1(M)\}-(m_{33}-m_{11}-1)\bar{E}(M)\\
&=C_1(M)\{(m_{13}-m_{33}+1)(m_{12}-m_{11}-\delta(M)\chi_+(M))+(-m_{13}+m_{11})(m_{13}-m_{33}+1)\\
&\quad+(m_{33}-m_{11}-1)\bar{C}_1(M)\}-(m_{33}-m_{11}-1)\bar{E}(M)\\
&=(-m_{13}+m_{11})C_1(M)(m_{13}-m_{33}+1-\bar{C}_1(M))\}-(m_{33}-m_{11}-1)\bar{E}(M)\\
&=(2m_{11}-m_{13}-m_{33}+1)\bar{E}(M)\\
&=(-2m_{11}+m_{13}+m_{33}-1)c^{\lambda}_{[13;22;1]}(M)\,.
\end{aligned}
$$

Hence we obtain the equation (5.22). Here we use the relations (4.1), (4.2), (4.9) and (4.13).

We have

$$
\begin{aligned}
&(m_{12}-m_{23}-1)\chi_-^{(-1)}(M)c^{\lambda}_{[13;12;1]}(M)\\
&=(m_{12}-m_{23}-1)\chi_-^{(-1)}(M)\{\bar{E}(M)+\bar{F}(M)-(m_{13}-m_{12})(m_{22}-m_{33})\chi_+(M)\}\\
&=(m_{12}-m_{23}-1)(1-\chi_+(M))(\bar{E}(M)-C_2(M)).
\end{aligned}
$$

Therefore

$$
\begin{aligned}
&(m_{12}-m_{23}-1)\chi_-^{(-1)}(M)c^{\lambda}_{[13;12;1]}(M)+(m_{11}-m_{22}-1)c^{\lambda}_{[13;12;2]}(M)\\
&\quad+(C_1(M)-1)\,c^{\lambda}_{[13;11;1]}(M)\\
&=(m_{12}-m_{23}-1)(1-\chi_+(M))(\bar{E}(M)-C_2(M))\\
&\quad+(m_{11}-m_{22}-1)(\bar{E}(M)-C_2(M))\chi_+(M)\\
&\quad+(C_1(M)-1)\,(C_2(M)-\bar{E}(M))\\
&=(m_{12}-m_{23}-\delta(M)\chi_+(M)-1)(\bar{E}(M)-C_2(M))\\
&\quad-(C_1(M)-1)\,(\bar{E}(M)-C_2(M))=0.
\end{aligned}
$$

Hence we obtain the equation (5.23). Here we use the relations (4.1), (4.9), (4.10) and (4.26).

We have

$$
\begin{aligned}
&(m_{12}-m_{23}-2)\chi_-^{(-1)}(M)c^{\lambda}_{[13;12;2]}(M)\\
&=(m_{12}-m_{23}-2)(\bar{E}(M)-C_2(M))\chi_-^{(-1)}(M)\chi_+(M)=0.
\end{aligned}
$$

Hence we obtain the equation (5.24). Here we use the relation (4.10).

- the proof of the case of the $(i,j)=(2,3)$.

Since $(r_{[23;22]},r_{[23;12]},r_{[23;11]})=(1,1,1)$, we have to confirm the following equations:

$$
\begin{aligned}
&(m_{11}-m_{22}+1)c^{\lambda}_{[23;12;0]}(M)+(m_{33}-m_{22})c^{\lambda}_{[23;11;0]}(M)\\
&=(-2m_{11}+m_{23}+m_{33}-2)c^{\lambda}_{[23;22;0]}(M)\,,
\end{aligned}
\tag{5.25}
$$

$$
\begin{aligned}
&(m_{12}-m_{23}+1)\chi_-(M)c^{\lambda}_{[23;12;0]}(M)+(m_{11}-m_{22})c^{\lambda}_{[23;12;1]}(M)\\
&\quad+(C_1(M)+\chi_-(M))\,c^{\lambda}_{[23;11;0]}(M)+(m_{33}-m_{22}-1)c^{\lambda}_{[23;11;1]}(M)\\
&=(-2m_{11}+m_{23}+m_{33}-2)c^{\lambda}_{[23;22;1]}(M)\,,
\end{aligned}
\tag{5.26}
$$

$$(m_{12} - m_{23})\chi_-(M)c^{\lambda}_{[23;12;1]}(M) + (C_1(M) - 1 + \chi_-(M))\, c^{\lambda}_{[23;11;1]}(M) = 0. \tag{5.27}$$

We have

$$\begin{aligned}
&(m_{11} - m_{22} + 1)c^{\lambda}_{[23;12;0]}(M) + (m_{33} - m_{22})c^{\lambda}_{[23;11;0]}(M)\\
&= -2(m_{11} - m_{22} + 1)(m_{22} - m_{33}) + (m_{33} - m_{22})(2m_{22} - m_{23} - m_{33})\\
&= (-2m_{11} + m_{23} + m_{33} - 2)(m_{22} - m_{33})\\
&= (-2m_{11} + m_{23} + m_{33} - 2)c^{\lambda}_{[23;22;0]}(M)\,.
\end{aligned}$$

Hence we obtain the equation (5.25).

We have

$$\begin{aligned}
&(m_{12} - m_{23} + 1)\chi_-(M)c^{\lambda}_{[23;12;0]}(M) + (m_{11} - m_{22})c^{\lambda}_{[23;12;1]}(M)\\
&+ (C_1(M) + \chi_-(M))\, c^{\lambda}_{[23;11;0]}(M) + (m_{33} - m_{22} - 1)c^{\lambda}_{[23;11;1]}(M)\\
&= -2(m_{12} - m_{23} + 1)\chi_-(M)(m_{22} - m_{33})\\
&\quad + (m_{11} - m_{22})\{\bar{C}_1(M) - (m_{22} - m_{33}) + \delta(M)\chi_-(M)\}\\
&\quad + (C_1(M) + \chi_-(M))\,(2m_{22} - m_{23} - m_{33}) - (m_{33} - m_{22} - 1)(\bar{C}_1(M) + \bar{D}(M))\chi_-(M)\\
&= -2(m_{12} - m_{23} + 1)(m_{22} - m_{33})\chi_-(M)\\
&\quad + (m_{11} - m_{22})(-2m_{22} + m_{23} + m_{33} + 2\delta(M)\chi_-(M))\\
&\quad + (m_{11} - m_{22} + (\delta(M) + 1)\chi_-(M))\,(2m_{22} - m_{23} - m_{33})\\
&\quad - (m_{33} - m_{22} - 1)(m_{12} - m_{11} + \bar{D}(M))\chi_-(M)\\
&= (2m_{11} - m_{22} - m_{23} + 1)(-m_{22} + m_{33} + \delta(M))\chi_-(M) + (m_{22} - m_{33} + 1)\bar{D}(M)\chi_-(M)\\
&= (2m_{11} - m_{23} - m_{33} + 2)\bar{D}(M)\chi_-(M)\\
&= (-2m_{11} + m_{23} + m_{33} - 2)c^{\lambda}_{[23;22;1]}(M)\,.
\end{aligned}$$

Hence we obtain the equation (5.26). Here we use the relations (4.1), (4.2) and (4.16).

We have

$$\begin{aligned}
&(m_{12} - m_{23})\chi_-(M)c^{\lambda}_{[23;12;1]}(M) + (C_1(M) - 1 + \chi_-(M))\, c^{\lambda}_{[23;11;1]}(M)\\
&= (m_{12} - m_{23})\chi_-(M)\{\bar{C}_1(M) - (m_{22} - m_{33}) + \delta(M)\chi_-(M)\}\\
&\quad - (C_1(M) - 1 + \chi_-(M))\,(\bar{C}_1(M) + \bar{D}(M))\chi_-(M)\\
&= (m_{12} - m_{23})\chi_-(M)(\bar{C}_1(M) - m_{22} + m_{33} + \delta(M)) - (m_{12} - m_{23})(\bar{C}_1(M) + \bar{D}(M))\chi_-(M)\\
&= 0.
\end{aligned}$$

Hence we obtain the equation (5.27). Here we use the relations (4.16) and (4.12).

□

Theorem 5.6. *For $1 \leq i \leq j \leq 3$, we have an following equation with the matrix representation $R(\Gamma^{\lambda}_{\pm ij}) \in M(d^{\sigma}_{\lambda[\pm ij]}, d^{\sigma}_{\lambda}, \mathbf{C})$ of $\Gamma^{\lambda}_{\pm ij}$ with respect to the induced basis $\{S_{\lambda}(M)\}_{M \in G_{\sigma}(\lambda)}$:*

$$\mathcal{C}^{\lambda}_{\pm ij}\mathbf{S}_{\sigma}(\lambda) = \mathbf{S}_{\sigma}(\lambda[\pm ij]) \cdot R(\Gamma^{\lambda}_{\pm ij}). \tag{5.28}$$

Here $\mathcal{C}^{\lambda}_{\pm ij}\mathbf{S}_{\sigma}(\lambda)$ is a matrix of the size $d_{\lambda[\pm ij]} \times d^{\sigma}_{\lambda}$, whose $l^{\sigma}(M)$-th column is $\mathcal{C}^{\lambda}_{\pm ij}S_{\lambda}(M)$ for $M \in G_{\sigma}(\lambda)$.

The explicit expression of the matrix $R(\Gamma^{\lambda}_{\pm ij})$ is given as follows.

(i) $\mathfrak{p}_+$-side:

$R(\Gamma^{\lambda}_{+ij})$ $\quad(1 \le i \le j \le 3)$ is a matrix of size $d^{\sigma}_{\lambda[+ij]} \times d^{\sigma}_{\lambda}$, whose $l^{\sigma}(M)$-th column is given by

$$\begin{aligned}
&(\nu_1+\rho_1+\gamma_1^M+k_{ij}(M)) \sum_{m=0}^{r_{[ij;22]}} c^{\lambda}_{[ij;22;m]}\left(M\left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\ 0\ 2\\ 2\end{smallmatrix}\right)[m]\right)\mathbf{u}^{\sigma}_{\lambda[+ij]}\left(M\left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\ 0\ 2\\ 2\end{smallmatrix}\right)[m]\right)\\
&+\sum_{m=0}^{r_{[ij;02]}} h_{[ij;m]}(\nu_2,M)\mathbf{u}^{\sigma}_{\lambda[+ij]}\left(M\left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\ 0\ 2\\ 0\end{smallmatrix}\right)[m]\right)\\
&+(\nu_3+\rho_3+\gamma_3^M)\sum_{m=0}^{r_{[ij;00]}} c^{\lambda}_{[ij;00;m]}\left(M\left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\ 0\ 0\\ 0\end{smallmatrix}\right)[m]\right)\mathbf{u}^{\sigma}_{\lambda[+ij]}\left(M\left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\ 0\ 0\\ 0\end{smallmatrix}\right)[m]\right)
\end{aligned}$$

for $M \in G_{\sigma}(\lambda)$. Here

$$\begin{aligned}
h_{[ij;m]}(\nu_2,M) =&(\nu_2+\rho_2+\gamma_2^M)c^{\lambda}_{[ij;02;m]}\left(M\left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\ 0\ 2\\ 0\end{smallmatrix}\right)[m]\right)\\
&+(m_{22}-m_{33}+1+\delta(M))\chi_{+}^{(-1)}(M)c^{\lambda}_{[ij;01;m-1]}\left(M\left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\ 0\ 2\\ 0\end{smallmatrix}\right)[m]\right)\\
&+(m_{22}-m_{33}+1)c^{\lambda}_{[ij;01;m]}\left(M\left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\ 0\ 2\\ 0\end{smallmatrix}\right)[m]\right)
\end{aligned}$$

$$\left(c^{\lambda}_{[ij;kl;m]}\left(M\left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\ 0\ 2\\ 2\end{smallmatrix}\right)[m]\right) = 0 \quad \textit{if} \quad r_{[ij;kl]} < m \ \textit{ or } \quad m<0. \right),$$

and $\mathbf{u}^{\sigma}_{\lambda}(M)$ is a column vector of degree d^{σ}_{λ} which is defined by

$$\mathbf{u}^{\sigma}_{\lambda}(M) = \begin{cases} {}^{t}(\overbrace{0,\cdots,0}^{l^{\sigma}(M)-1},1,\overbrace{0,\cdots,0}^{d^{\sigma}_{\lambda}-l^{\sigma}(M)}) & \textit{if } M \in G_{\sigma}(\lambda),\\ \mathbf{0} & \textit{otherwise .}\end{cases}$$

(ii) $\mathfrak{p}_{-}$-side:
$R(\Gamma^{\lambda}_{-ij})$ $\quad(1 \le i \le j \le 3)$ is a matrix of size $d^{\sigma}_{\lambda[-ij]} \times d^{\sigma}_{\lambda}$, whose $l^{\sigma}(M)$-th column is given by the form

$$\begin{aligned}
&(\nu_1+\rho_1-\gamma_1^M+k_{4-j\,4-i}(\hat{M})) \sum_{m=0}^{r_{[4-j\,4-i;22]}} c^{\widehat{\lambda}}_{[4-j\,4-i;22;m]}\left(\hat{M}\left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\ 0\ 2\\ 2\end{smallmatrix}\right)[m]\right)\mathbf{u}^{\sigma}_{\lambda[-ij]}\left(M\left(\begin{smallmatrix}-\mathbf{e}_i-\mathbf{e}_j\\ -2\ 0\\ -2\end{smallmatrix}\right)[m]\right)\\
&+\sum_{m=0}^{r_{[4-j\,4-i;02]}} h_{[4-j\,4-i;m]}(\nu_2,\hat{M})\mathbf{u}^{\sigma}_{\lambda[-ij]}\left(M\left(\begin{smallmatrix}-\mathbf{e}_i-\mathbf{e}_j\\ -2\ 0\\ 0\end{smallmatrix}\right)[m]\right)\\
&+(\nu_3+\rho_3-\gamma_3^M)\sum_{m=0}^{r_{[4-j\,4-i;00]}} c^{\widehat{\lambda}}_{[4-j\,4-i;00;m]}\left(\hat{M}\left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\ 0\ 0\\ 0\end{smallmatrix}\right)[m]\right)\mathbf{u}^{\sigma}_{\lambda[-ij]}\left(M\left(\begin{smallmatrix}-\mathbf{e}_i-\mathbf{e}_j\\ 0\ 0\\ 0\end{smallmatrix}\right)[m]\right)
\end{aligned}$$

for $M \in G_{\sigma}(\lambda)$.

Proof. For $M, N \in G(\lambda)$, we define

$$\Delta(M,N) = \begin{cases} 1 & \text{if } M = N,\\ 0 & \text{otherwise.}\end{cases}$$

Since

$$s(M,N)(1_6) = \langle f(M)^*, f(N)\rangle = \Delta(M,N), \tag{5.29}$$

we see that the value at $1_6 \in G$ of the vector $S_{\lambda}(M)$ is the $l(M)$-th unit column vector ${}^{t}(0,\cdots,0,1,0,\cdots,0)$ of degree d_{λ}. Therefore, we note that it suffices to evaluate the both of the equation (5.28) at $1_6 \in G$.

First, we compute $\{X_{+pq}s(M,N)\}(1_6)$ for $1 \le p \le q \le 3$. Since $\{s(M,N)\}_{N\in G(\lambda)}$ is the monomial basis of $\langle S_\lambda\,(M)\rangle$, we obtain

$$\{\kappa(E_{pp})s(M,N)\}(1_6) = \gamma_p^N s(M,N)(1_6) = \gamma_p^M \Delta(M,N)\ , \quad 1 \le p \le 3,$$

$$\begin{aligned}
&\{\kappa(E_{21})s(M,N)\}(1_6)\\
&\quad = (n_{11}-n_{22})s(M,N\left(\begin{smallmatrix}0&&0\\&-1&\end{smallmatrix}\right))(1_6) + (n_{12}-n_{23})\chi_-(N)s(M,N\left(\begin{smallmatrix}0&&0\\&-1&\end{smallmatrix}\right)[-1])(1_6)\\
&\quad = (m_{11}-m_{22}+1)\Delta(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right),N) + (m_{12}-m_{23}+1)\chi_-^{(-1)}(M)\Delta(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)[1],N)
\end{aligned}$$

$$\begin{aligned}
&\{\kappa(E_{31})s(M,N)\}(1_6)\\
&\quad = (n_{33}-n_{22})s(M,N\left(\begin{smallmatrix}0&&-1\\&-1&\end{smallmatrix}\right))(1_6) + C_1(N)s(M,N\left(\begin{smallmatrix}0&&-1\\&-1&\end{smallmatrix}\right)[-1])(1_6)\\
&\quad = (m_{33}-m_{22}-1)\Delta(M\left(\begin{smallmatrix}0&&1\\&1&\end{smallmatrix}\right),N) + (C_1(M)+1)\Delta(M\left(\begin{smallmatrix}0&&1\\&1&\end{smallmatrix}\right)[1],N),
\end{aligned}$$

$$\begin{aligned}
&\{\kappa(E_{32})s(M,N)\}(1_6)\\
&\quad = (n_{22}-n_{33})s(M,N\left(\begin{smallmatrix}0&&-1\\&0&\end{smallmatrix}\right))(1_6) + \{n_{22}-n_{33}+\delta(N)\}\chi_+(N)s(M,N\left(\begin{smallmatrix}0&&-1\\&0&\end{smallmatrix}\right)[-1])(1_6)\\
&\quad = (m_{22}-m_{33}+1)\Delta(M\left(\begin{smallmatrix}0&&1\\&0&\end{smallmatrix}\right),N) + \{m_{22}-m_{33}+1+\delta(M)\}\chi_+^{(-1)}(N)\Delta(M\left(\begin{smallmatrix}0&&1\\&0&\end{smallmatrix}\right)[1],N).
\end{aligned}$$

by Proposition 3.1 and the equations (3.2), (3.2). Moreover, we obtain

$$\begin{aligned}
&\{E_\alpha s(M,N)\}(1_6) = 0, && \alpha \in \Sigma^+,\\
&\{H_p s(M,N)\}(1_6) = (\nu_p+\rho_p)s(M,N)(1_6), && 1 \le p \le 3
\end{aligned}$$

by the definition of principal series representation. By above computations and Iwasawa decomposition in Lemma 5.4, we obtain

$$\begin{aligned}
\{X_{+pp}s(M,N)\}(1_6) =&(\nu_p+\rho_p+\gamma_p^M)\Delta(M,N)\ , \quad 1 \le p \le 3,\\
\{X_{+12}s(M,N)\}(1_6) =&(m_{11}-m_{22}+1)\Delta(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right),N)\\
&+ (m_{12}-m_{23}+1)\chi_-^{(-1)}(M)\Delta(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)[1],N),\\
\{X_{+13}s(M,N)\}(1_6) =&(m_{33}-m_{22}-1)\Delta(M\left(\begin{smallmatrix}0&&1\\&1&\end{smallmatrix}\right),N) + (C_1(M)+1)\Delta(M\left(\begin{smallmatrix}0&&1\\&1&\end{smallmatrix}\right)[1],N),\\
\{X_{+23}s(M,N)\}(1_6) =&(m_{22}-m_{33}+1)\Delta(M\left(\begin{smallmatrix}0&&1\\&0&\end{smallmatrix}\right),N)\\
&+ \{m_{22}-m_{33}+1+\delta(M)\}\chi_+^{(-1)}(M)\Delta(M\left(\begin{smallmatrix}0&&1\\&0&\end{smallmatrix}\right)[1],N).
\end{aligned}$$

Let $\mathbf{u}_\lambda(N)$ be a column vector of degree d_λ which is defined by

$$\mathbf{u}_\lambda(N) = \begin{cases} {}^t(\overbrace{0,\cdots,0}^{l(N)-1},1,\overbrace{0,\cdots,0}^{d_\lambda-l(N)}) & \text{if } M \in G(\lambda),\\ \mathbf{0} & \text{otherwise .}\end{cases}$$

We denote by $X_{+pq}S_\lambda\,(M)$ the vector of degree d_λ whose $l(N)$-th component is $X_{+pq}s(M,N)$. Then we obtain

$$\begin{aligned}
\{X_{+pp}S_\lambda\,(M)\}(1_6) =&(\nu_p+\rho_p+\gamma_p^M)\mathbf{u}_\lambda(M) \quad \text{for } \ 1 \le p \le 3,\\
\{X_{+12}S_\lambda\,(M)\}(1_6) =&(m_{12}-m_{23}+1)\chi_-^{(-1)}(M)\mathbf{u}_\lambda\left(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)[1]\right)\\
&+ (m_{11}-m_{22}+1)\mathbf{u}_\lambda\left(M\left(\begin{smallmatrix}0&&0\\&1&\end{smallmatrix}\right)\right),\\
\{X_{+13}S_\lambda\,(M)\}(1_6) =&(C_1(M)+1)\mathbf{u}_\lambda\left(M\left(\begin{smallmatrix}1&&0\\&1&\end{smallmatrix}\right)[1]\right) + (m_{33}-m_{22}-1)\mathbf{u}_\lambda\left(M\left(\begin{smallmatrix}1&&0\\&1&\end{smallmatrix}\right)\right),\\
\{X_{+23}S_\lambda\,(M)\}(1_6) =&\,(m_{22}-m_{33}+1+\delta(M))\,\chi_+^{(-1)}(M)\mathbf{u}_\lambda\left(M\left(\begin{smallmatrix}1&&0\\&0&\end{smallmatrix}\right)[1]\right)\\
&+ (m_{22}-m_{33}+1)\mathbf{u}_\lambda\left(M\left(\begin{smallmatrix}1&&0\\&0&\end{smallmatrix}\right)\right).
\end{aligned}$$

Let us compute $\{\mathcal{C}_{+ij}^\lambda S_\lambda\,(M)\}(1_6)$.

Since

$$L^{\lambda}_{+ij}\left(\left(\begin{smallmatrix} 0\;-l \\ -k \end{smallmatrix}\right)[-m]\right)\cdot \mathbf{u}_{\lambda}(N) = c^{\lambda}_{[ij;kl;m]}\left(N\left(\begin{smallmatrix} \mathbf{e}_i+\mathbf{e}_j \\ 0\;l \\ k \end{smallmatrix}\right)[m]\right)\mathbf{u}_{\lambda[+ij]}\left(N\left(\begin{smallmatrix} \mathbf{e}_i+\mathbf{e}_j \\ 0\;l \\ k \end{smallmatrix}\right)[m]\right)$$

for $N \in G(\lambda)$, we obtain

(5.30)

$$\begin{aligned}
&\left\{\left(\sum_{m=0}^{r_{[ij;22]}} L^{\lambda}_{+ij}\left(\left(\begin{smallmatrix} 0\;-2 \\ -2 \end{smallmatrix}\right)[-m]\right)\otimes X_{+11}\right) S_{\lambda}(M)\right\}(1_6) \\
&= \left(\sum_{m=0}^{r_{[ij;22]}} L^{\lambda}_{+ij}\left(\left(\begin{smallmatrix} 0\;-2 \\ -2 \end{smallmatrix}\right)[-m]\right)\right)\cdot\{X_{+11}S_{\lambda}(M)\}(1_6) \\
&= (\nu_1+\rho_1+\gamma_1^M)\sum_{m=0}^{r_{[ij;22]}} c^{\lambda}_{[ij;22;m]}\left(M\left(\begin{smallmatrix} \mathbf{e}_i+\mathbf{e}_j \\ 0\;2 \\ 2 \end{smallmatrix}\right)[m]\right)\mathbf{u}_{\lambda[+ij]}\left(M\left(\begin{smallmatrix} \mathbf{e}_i+\mathbf{e}_j \\ 0\;2 \\ 2 \end{smallmatrix}\right)[m]\right),
\end{aligned}$$

(5.31)

$$\begin{aligned}
&\left\{\left(\sum_{m=0}^{r_{[ij;12]}} L^{\lambda}_{+ij}\left(\left(\begin{smallmatrix} 0\;-2 \\ -1 \end{smallmatrix}\right)[-m]\right)\otimes X_{+12}\right) S_{\lambda}(M)\right\}(1_6) \\
&= \left(\sum_{m=0}^{r_{[ij;12]}} L^{\lambda}_{+ij}\left(\left(\begin{smallmatrix} 0\;-2 \\ -1 \end{smallmatrix}\right)[-m]\right)\right)\cdot\{X_{+12}S_{\lambda}(M)\}(1_6) \\
&= (m_{12}-m_{23}+1)\chi_{-}^{(-1)}(M)\sum_{m=0}^{r_{[ij;12]}} c^{\lambda}_{[ij;12;m]}\left(M\left(\begin{smallmatrix} \mathbf{e}_i+\mathbf{e}_j \\ 0\;2 \\ 2 \end{smallmatrix}\right)[m+1]\right)\mathbf{u}_{\lambda}\left(M\left(\begin{smallmatrix} \mathbf{e}_i+\mathbf{e}_j \\ 0\;2 \\ 2 \end{smallmatrix}\right)[m+1]\right) \\
&\qquad + (m_{11}-m_{22}+1)\sum_{m=0}^{r_{[ij;12]}} c^{\lambda}_{[ij;12;m]}\left(M\left(\begin{smallmatrix} \mathbf{e}_i+\mathbf{e}_j \\ 0\;2 \\ 2 \end{smallmatrix}\right)[m]\right)\mathbf{u}_{\lambda}\left(M\left(\begin{smallmatrix} \mathbf{e}_i+\mathbf{e}_j \\ 0\;2 \\ 2 \end{smallmatrix}\right)[m]\right), \\
&= \sum_{m=0}^{r_{[ij;12]}+1}\Bigg\{(m_{12}-m_{23}+1)\chi_{-}^{(-1)}(M)c^{\lambda}_{[ij;12;m-1]}\left(M\left(\begin{smallmatrix} \mathbf{e}_i+\mathbf{e}_j \\ 0\;2 \\ 2 \end{smallmatrix}\right)[m]\right)\mathbf{u}_{\lambda}\left(M\left(\begin{smallmatrix} \mathbf{e}_i+\mathbf{e}_j \\ 0\;2 \\ 2 \end{smallmatrix}\right)[m]\right) \\
&\qquad + (m_{11}-m_{22}+1)c^{\lambda}_{[ij;12;m]}\left(M\left(\begin{smallmatrix} \mathbf{e}_i+\mathbf{e}_j \\ 0\;2 \\ 2 \end{smallmatrix}\right)[m]\right)\Bigg\}\mathbf{u}_{\lambda}\left(M\left(\begin{smallmatrix} \mathbf{e}_i+\mathbf{e}_j \\ 0\;2 \\ 2 \end{smallmatrix}\right)[m]\right),
\end{aligned}$$

(5.32)

$$\begin{aligned}
&\left\{\left(\sum_{m=0}^{r_{[ij;11]}} L^{\lambda}_{+ij}\left(\left(\begin{smallmatrix} 0\;-1 \\ -1 \end{smallmatrix}\right)[-m]\right)\otimes X_{+13}\right) S_{\lambda}(M)\right\}(1_6) \\
&= \left(\sum_{m=0}^{r_{[ij;11]}} L^{\lambda}_{+ij}\left(\left(\begin{smallmatrix} 0\;-1 \\ -1 \end{smallmatrix}\right)[-m]\right)\right)\cdot\{X_{+13}S_{\lambda}(M)\}(1_6) \\
&= (C_1(M)+1)\sum_{m=0}^{r_{[ij;11]}} c^{\lambda}_{[ij;11;m]}\left(M\left(\begin{smallmatrix} \mathbf{e}_i+\mathbf{e}_j \\ 0\;2 \\ 2 \end{smallmatrix}\right)[m+1]\right)\mathbf{u}_{\lambda}\left(M\left(\begin{smallmatrix} \mathbf{e}_i+\mathbf{e}_j \\ 0\;2 \\ 2 \end{smallmatrix}\right)[m+1]\right) \\
&\qquad + (m_{33}-m_{22}-1)\sum_{m=0}^{r_{[ij;11]}} c^{\lambda}_{[ij;12;m]}\left(M\left(\begin{smallmatrix} \mathbf{e}_i+\mathbf{e}_j \\ 0\;2 \\ 2 \end{smallmatrix}\right)[m]\right)\mathbf{u}_{\lambda}\left(M\left(\begin{smallmatrix} \mathbf{e}_i+\mathbf{e}_j \\ 0\;2 \\ 2 \end{smallmatrix}\right)[m]\right)
\end{aligned}$$

$$\begin{aligned}
&= \sum_{m=0}^{r_{[ij;11]}+1} \Big\{ (C_1(M)+1) c^{\lambda}_{[ij;11;m-1]} \left(M \left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\ 0\;2\\ 2\end{smallmatrix}\right)[m] \right) \\
&\quad + (m_{33}-m_{22}-1) c^{\lambda}_{[ij;12;m]} \left(M \left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\ 0\;2\\ 2\end{smallmatrix}\right)[m] \right) \Big\} \mathbf{u}_{\lambda} \left(M \left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\ 0\;2\\ 2\end{smallmatrix}\right)[m] \right).
\end{aligned}$$

By the equations (5.30), (5.31), (5.32) and Lemma 5.5, we obtain

$$\begin{aligned}
&\Bigg\{ \Bigg(\sum_{m=0}^{r_{[ij;22]}} L^{\lambda}_{+ij} \left(\left(\begin{smallmatrix}0&-2\\&-2\end{smallmatrix}\right)[-m] \right) \otimes X_{+11} + \sum_{m=0}^{r_{[ij;12]}} L^{\lambda}_{+ij} \left(\left(\begin{smallmatrix}0&-2\\&-1\end{smallmatrix}\right)[-m] \right) \otimes X_{+12} \\
&\quad + \sum_{m=0}^{r_{[ij;11]}} L^{\lambda}_{+ij} \left(\left(\begin{smallmatrix}0&-1\\&-1\end{smallmatrix}\right)[-m] \right) \otimes X_{+13} \Bigg) S_{\lambda}(M) \Bigg\} (1_6) \\
&= (\nu_1+\rho_1+\gamma^M_1+k_{ij}(M)) \sum_{m=0}^{r_{[ij;22]}} c^{\lambda}_{[ij;22;m]} \left(M \left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\ 0\;2\\ 2\end{smallmatrix}\right)[m] \right) \mathbf{u}_{\lambda[+ij]} \left(M \left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\ 0\;2\\ 2\end{smallmatrix}\right)[m] \right).
\end{aligned} \tag{5.33}$$

Similarly,

$$\begin{aligned}
&\Bigg\{ \Bigg(\sum_{m=0}^{4} L^{\lambda}_{+11} \left(\left(\begin{smallmatrix}0&-2\\&0\end{smallmatrix}\right)[-m] \right) \otimes X_{+22} \Bigg) S_{\lambda}(M) \Bigg\} (1_6) \\
&= \Bigg(\sum_{m=0}^{r_{[ij;02]}} L^{\lambda}_{+ij} \left(\left(\begin{smallmatrix}0&-2\\&0\end{smallmatrix}\right)[-m] \right) \Bigg) \cdot \{ X_{+22} S_{\lambda}(M) \} (1_6) \\
&= (\nu_2+\rho_2+\gamma^M_2) \sum_{m=0}^{r_{[ij;02]}} c^{\lambda}_{[ij;02;m]} \left(M \left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\ 0\;2\\ 0\end{smallmatrix}\right)[m] \right) \mathbf{u}_{\lambda[+ij]} \left(M \left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\ 0\;2\\ 0\end{smallmatrix}\right)[m] \right),
\end{aligned} \tag{5.34}$$

$$\begin{aligned}
&\Bigg\{ \Bigg(\sum_{m=0}^{r_{[ij;01]}} L^{\lambda}_{+ij} \left(\left(\begin{smallmatrix}0&-1\\&0\end{smallmatrix}\right)[-m] \right) \otimes X_{+23} \Bigg) S_{\lambda}(M) \Bigg\} (1_6) \\
&= \Bigg(\sum_{m=0}^{r_{[ij;01]}} L^{\lambda}_{+ij} \left(\left(\begin{smallmatrix}0&-1\\&0\end{smallmatrix}\right)[-m] \right) \Bigg) \cdot \{ X_{+23} S_{\lambda}(M) \} (1_6) \\
&= (m_{22}-m_{33}+1+\delta(M)) \chi^{(-1)}_{+}(M) \sum_{m=0}^{r_{[ij;01]}} c^{\lambda}_{[ij;01;m]} \left(M \left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\ 0\;2\\ 0\end{smallmatrix}\right)[m+1] \right) \\
&\qquad \times \mathbf{u}_{\lambda[+ij]} \left(M \left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\ 0\;2\\ 0\end{smallmatrix}\right)[m+1] \right) \\
&\qquad + (m_{22}-m_{33}+1) \sum_{m=0}^{r_{[ij;01]}} c^{\lambda}_{[ij;01;m]} \left(M \left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\ 0\;2\\ 0\end{smallmatrix}\right)[m] \right) \mathbf{u}_{\lambda[+ij]} \left(M \left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\ 0\;2\\ 0\end{smallmatrix}\right)[m] \right) \\
&= \sum_{m=0}^{r_{[ij;01]}+1} \Big\{ (m_{22}-m_{33}+1+\delta(M)) \chi^{(-1)}_{+}(M) c^{\lambda}_{[ij;01;m-1]} \left(M \left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\ 0\;2\\ 0\end{smallmatrix}\right)[m] \right) \\
&\qquad + (m_{22}-m_{33}+1) c^{\lambda}_{[ij;01;m]} \left(M \left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\ 0\;2\\ 0\end{smallmatrix}\right)[m] \right) \Big\} \mathbf{u}_{\lambda[+ij]} \left(M \left(\begin{smallmatrix}\mathbf{e}_i+\mathbf{e}_j\\ 0\;2\\ 0\end{smallmatrix}\right)[m] \right).
\end{aligned} \tag{5.35}$$

By direct computation from the equations (5.34) and (5.35), we obtain

$$\Bigg\{ \Bigg(\sum_{m=0}^{r_{[ij;02]}} L^{\lambda}_{+ij} \left(\left(\begin{smallmatrix}0&-2\\&0\end{smallmatrix}\right)[-m] \right) \otimes X_{+22} + \sum_{m=0}^{r_{[ij;01]}} L^{\lambda}_{+ij} \left(\left(\begin{smallmatrix}0&-1\\&0\end{smallmatrix}\right)[-m] \right) \otimes X_{+23} \Bigg) S_{\lambda}(M) \Bigg\} (1_6) \tag{5.36}$$

$$= \sum_{m=0}^{r_{[ij;02]}} h_{[ij;m]}(\nu_2, M)\mathbf{u}_{\lambda[+ij]}\left(M\left(\substack{\mathbf{e}_i+\mathbf{e}_j\\ 0\,2\\ 0}\right)[m]\right).$$

Similarly,

$$\left\{\left(\sum_{m=0}^{r_{[ij;00]}} L^{\lambda}_{+ij}\left(\left(\substack{0\,0\\ 0}\right)[-m]\right)\otimes X_{+33}\right)S_{\lambda}(M)\right\}(1_6) \tag{5.37}$$
$$= \left(\sum_{m=0}^{r_{[ij;00]}} L^{\lambda}_{+ij}\left(\left(\substack{0\,0\\ 0}\right)[-m]\right)\right)\cdot\{X_{+33}S_{\lambda}(M)\}(1_6)$$
$$= (\nu_3+\rho_3+\gamma^M_3)\sum_{m=0}^{r_{[ij;00]}} c^{\lambda}_{[ij;00;m]}\left(M\left(\substack{\mathbf{e}_i+\mathbf{e}_j\\ 0\,0\\ 0}\right)[m]\right)\mathbf{u}_{\lambda[+ij]}\left(M\left(\substack{\mathbf{e}_i+\mathbf{e}_j\\ 0\,0\\ 0}\right)[m]\right).$$

Summing up the equations (5.33), (5.36) and (5.37), we obtain

$$\{\mathcal{C}^{\lambda}_{11}S_{\lambda}(M)\}(1_6) \tag{5.38}$$
$$= (\nu_1+\rho_1+\gamma^M_1+k_{ij}(M))\sum_{m=0}^{r_{[ij;22]}} c^{\lambda}_{[ij;22;m]}\left(M\left(\substack{\mathbf{e}_i+\mathbf{e}_j\\ 0\,2\\ 2}\right)[m]\right)\mathbf{u}_{\lambda[+ij]}\left(M\left(\substack{\mathbf{e}_i+\mathbf{e}_j\\ 0\,2\\ 2}\right)[m]\right)$$
$$+\sum_{m=0}^{r_{[ij;02]}} h_{[ij;m]}(\nu_2, M)\mathbf{u}_{\lambda[+ij]}\left(M\left(\substack{\mathbf{e}_i+\mathbf{e}_j\\ 0\,2\\ 0}\right)[m]\right)$$
$$+(\nu_3+\rho_3+\gamma^M_3)\sum_{m=0}^{r_{[ij;00]}} c^{\lambda}_{[ij;00;m]}\left(M\left(\substack{\mathbf{e}_i+\mathbf{e}_j\\ 0\,0\\ 0}\right)[m]\right)\mathbf{u}_{\lambda[+ij]}\left(M\left(\substack{\mathbf{e}_i+\mathbf{e}_j\\ 0\,0\\ 0}\right)[m]\right).$$

By the remark at the beginning of this proof, we obtain the assertion of the case of (i). The case of (ii) is treated similarly. □

6. EXAMPLES OF CONTIGUOUS RELATIONS AND THEIR APPLICATIONS

Here are some examples of the contiguous relations at the peripheral K-types. Moreover, as their applications, we determine the holonomic systems, whose solutions are Whittaker functions.

6.1. **Whittaker functions.**

For a unitary character ξ of $N_{\min}$, we denote the derivative of ξ by the same letter. Since

$$\mathfrak{n}/[\mathfrak{n},\mathfrak{n}] \simeq \mathfrak{g}_{e_1-e_2}\oplus\mathfrak{g}_{e_2-e_3}\oplus\mathfrak{g}_{2e_3},$$

ξ is specified by three real numbers c_{12}, c_{23} and c_3 such that

$$\xi(E_{e_1-e_2}) = 2\pi\sqrt{-1}c_{12},\ \xi(E_{e_2-e_3}) = 2\pi\sqrt{-1}c_{23} \text{ and } \xi(E_{2e_3}) = 2\pi\sqrt{-1}c_3.$$

When $c_{12}c_{23}c_3 \neq 0$, a unitary character ξ of $N_{\min}$ is called *non-degenerate.*

For a finite dimensional representation (τ, V) of K and a non-degenerate unitary character ξ of $N_{\min}$, we consider the space $C^{\infty}_{\xi,\tau}(N_{\min}\backslash G/K)$ of smooth functions $\varphi\colon G\to V_{\tau}$ with the property

$$\varphi(ngk) = \xi(n)\tau(k)^{-1}\varphi(g),\ (n,g,k)\in N_{\min}\times G\times K.$$

Here we remark that any functions $\varphi\in C^{\infty}_{\xi,\tau}(N_{\min}\backslash G/K)$ is determined by its restriction $\varphi|_{A_{\min}}$ to $A_{\min}$ from the Iwasawa decomposition $G = N_{\min}A_{\min}K$ of G. $\varphi|_{A_{\min}}$ is called the $A_{\min}$-radial part of φ. Also let $C^{\infty}\operatorname{Ind}^{G}_{N_{\min}}(\xi)$ be the C^{∞}-induced representation from ξ with the representation space

$$C^{\infty}_{\xi}(N_{\min}\backslash G) = \{\varphi\in C^{\infty}(G)\mid\varphi(ng) = \xi(n)\varphi(g),\ (n,g)\in N_{\min}\times G\},$$

on which G acts by right translation. Then the space $C^\infty_{\xi,\tau}(N_{\min}\backslash G/K)$ is isomorphic to $\mathrm{Hom}_K(V^*, C^\infty_\xi(N_{\min}\backslash G))$ via the correspondence between $\iota \in \mathrm{Hom}_K(V^*, C^\infty_\xi(N_{\min}\backslash G))$ and $F^{[\iota]} \in C^\infty_{\xi,\tau}(N_{\min}\backslash G/K)$ given by the relation $\iota(v^*)(g) = \langle v^*, F^{[\iota]}(g)\rangle$ for $v^* \in V^*$ and $g \in G$ with canonical pairing $\langle , \rangle$ on $V^* \times V$.

Let (π, H_π) be an irreducible admissible representation of G, and take a K-type (τ^*, V^*) of π with an injective K-homomorphism $\eta\colon V^* \to H_\pi$. Then, for each element T in the intertwining space $\mathcal{I}_{\xi,\pi} = \mathrm{Hom}_{(\mathfrak{g}_{\mathbf{C}},K)}(H_{\pi,K}, C^\infty_\xi(N_{\min}\backslash G))$, the relation $T(\eta(v^*))(g) = \langle v^*, \Phi(T,\eta)(g)\rangle$ $(v^* \in V^*,\ g \in G)$ determines an element $\Phi(T,\eta) \in C^\infty_{\xi,\tau}(N_{\min}\backslash G/K)$. Here $H_{\pi,K}$ is a subspace of H_π, consisting of all K-finite vectors. Now we put

$$\mathrm{Wh}(\pi,\xi,\tau) = \bigcup_{\eta \in \mathrm{Hom}_K(V^*, H_{\pi,K})} \{\Phi(T,\eta) \in C^\infty_{\xi,\tau}(N_{\min}\backslash G/K) \mid T \in \mathcal{I}_{\xi,\pi}\}$$

and call $\mathrm{Wh}(\pi,\xi,\tau)$ *the space of Whittaker functions* for (π,ξ,τ). We consider the Whittaker functions for the irreducible principal series representation $\pi = \pi_{(\sigma,\nu)}$.

6.2. **The $\pm$-chilarity matrices.** We define the $\pm$-chilarity matrices as follows.

Definition 6.1. *T*he $\pm$-chilarity matrices $m_i(C_\pm)$ for $1 \le i \le 3$ are defined by

$$m_1(C_\pm) = \begin{pmatrix} X_{\pm 11} & X_{\pm 12} & X_{\pm 13} \\ X_{\pm 12} & X_{\pm 22} & X_{\pm 23} \\ X_{\pm 13} & X_{\pm 23} & X_{\pm 33} \end{pmatrix}, \qquad m_2(C_\pm) = \begin{pmatrix} M_{\pm 11} & -M_{\pm 12} & M_{\pm 13} \\ -M_{\pm 12} & M_{\pm 22} & -M_{\pm 23} \\ M_{\pm 13} & -M_{\pm 23} & M_{\pm 33} \end{pmatrix},$$

and $m_3(C_\pm) = \det(m_1(C_\pm))$. Here $M_{\pm ij}$ is (i,j)-minor of the matrix $m_1(C_\pm)$ for each $1 \le i \le j \le 3$, that is,

$$M_{\pm 11} = \begin{vmatrix} X_{\pm 22} & X_{\pm 23} \\ X_{\pm 23} & X_{\pm 33} \end{vmatrix}, \qquad M_{\pm 22} = \begin{vmatrix} X_{\pm 11} & X_{\pm 13} \\ X_{\pm 13} & X_{\pm 33} \end{vmatrix}, \qquad M_{\pm 33} = \begin{vmatrix} X_{\pm 11} & X_{\pm 12} \\ X_{\pm 12} & X_{\pm 22} \end{vmatrix},$$

$$M_{\pm 12} = \begin{vmatrix} X_{\pm 12} & X_{\pm 23} \\ X_{\pm 13} & X_{\pm 33} \end{vmatrix}, \qquad M_{\pm 13} = \begin{vmatrix} X_{\pm 12} & X_{\pm 22} \\ X_{\pm 13} & X_{\pm 23} \end{vmatrix}, \qquad M_{\pm 23} = \begin{vmatrix} X_{\pm 11} & X_{\pm 12} \\ X_{\pm 13} & X_{\pm 23} \end{vmatrix}.$$

Then we can find the following lemma immediately from the definition of the chilarity matrices.

Lemma 6.2. *For each $1 \le i \le 3$, the element $C_{2i} = \mathrm{Tr}(m_i(C_+)m_i(C_-))$ in $U(\mathfrak{g}_{\mathbf{C}})$ is invariant under the adjoint action of K, that is,*

$$C_{2i} \in U(\mathfrak{g}_{\mathbf{C}})^K = \{X \in U(\mathfrak{g}_{\mathbf{C}}) \mid \mathrm{Ad}(k)X = X,\ k \in K\}.$$

Remark 6.3. *In the case of $Sp(n,\mathbf{R})$, we can define C_{2i} for each $1 \le i \le n$ belonging to $U(\mathfrak{g}_{\mathbf{C}})^K$ similarly. The operator C_{2i} is essentially same as the so-called Maass shift operator in the classical literature* [9]. *Also, the chilarity matrices are used to construct the Capelli elements for a symmetric pair in* [8], *recently.*

Now we consider a system of differential equations which are satisfied by the $A_{\min}$-radial part of each element in $\mathrm{Wh}(\pi,\xi,\tau)$ when τ^* is a multiplicity one K-type of $\pi_{(\sigma,\lambda)}$. Let λ be the highest weight of τ^*. The elements C_2, C_4 and C_6 in $U(\mathfrak{g}_{\mathbf{C}})^K$ defined in Lemma 6.2 are acting on the space $C^\infty_\xi(N_{\min}\backslash G)$ as differential operators and acting on the space $H_{(\sigma,\nu)}$ as scalar operators. Therefore, each element Φ in $\mathrm{Wh}(\pi,\xi,\tau)$ satisfies the following system of differential equations

$$C_{2i}\Phi = \chi_{2i,\sigma,\nu,\lambda}\Phi \quad (1 \le i \le 3),$$

where $\chi_{2i,\sigma,\nu,\lambda}$ is the scalar value for the action of the operator C_{2i}. In the later subsections, we compute those values for peripheral K-types.

6.3. **The chilarity operators in the cases of** $(\sigma_1,\sigma_2,\sigma_3)=(0,0,0),\ (1,1,1)$**.** In this subsection, we consider the cases of $(\sigma_1,\sigma_2,\sigma_3)=(0,0,0),\ (1,1,1)$. We set $M_l=\left(\begin{smallmatrix} l\,l\,l \\ l\,l \\ l \end{smallmatrix}\right)\in G((l,l,l))$ for $l\in\mathbf{Z}$. Let ε_σ be the integer defined by ε

$$\varepsilon_\sigma=\begin{cases} 0 & \text{if } (\sigma_1,\sigma_2,\sigma_3)=(0,0,0),\\ 1 & \text{if } (\sigma_1,\sigma_2,\sigma_3)=(1,1,1).\end{cases}$$

Since $(\sigma_1,\sigma_2,\sigma_3)=(0,0,0),\ (1,1,1)$, there are multiplicity one K-types $\tau_{(l,l,l)}$ ($l\equiv\varepsilon_\sigma \bmod 2$) of $\pi_{(\sigma,\nu)}$. Let us compute the values of $\chi_{2i,\sigma,\nu,(l,l,l)}$ $(1\leq i\leq 3)$.

Lemma 6.4. *(i) Let σ be a character of $M_{\min}$ such that $(\sigma_1,\sigma_2,\sigma_3)=(0,0,0),\ (1,1,1)$. For $l\equiv\varepsilon_\sigma \bmod 2$, the following equations hold:*

$$\begin{aligned}
\mathcal{C}^{(l-2,l-2,l-2)}_{+11}\mathbf{S}_\sigma((l-2,l-2,l-2)) &=\mathbf{S}_\sigma((l,l-2,l-2))\cdot R(\Gamma^{(l-2,l-2,l-2)}_{+11}),\\
\mathcal{C}^{(l,l-2,l-2)}_{+22}\mathbf{S}_\sigma((l,l-2,l-2)) &=\mathbf{S}_\sigma((l,l,l-2))\cdot R(\Gamma^{(l,l-2,l-2)}_{+22}),\\
\mathcal{C}^{(l,l,l-2)}_{+33}\mathbf{S}_\sigma((l,l,l-2)) &=\mathbf{S}_\sigma((l,l,l))\cdot R(\Gamma^{(l,l,l-2)}_{+33}),\\
\mathcal{C}^{(l,l,l)}_{-33}\mathbf{S}_\sigma((l,l,l)) &=\mathbf{S}_\sigma((l,l,l-2))\cdot R(\Gamma^{(l,l,l)}_{-33}),\\
\mathcal{C}^{(l,l,l-2)}_{-22}\mathbf{S}_\sigma((l,l,l-2)) &=\mathbf{S}_\sigma((l,l-2,l-2))\cdot R(\Gamma^{(l,l,l-2)}_{-22}),\\
\mathcal{C}^{(l,l-2,l-2)}_{-11}\mathbf{S}_\sigma((l,l-2,l-2)) &=\mathbf{S}_\sigma((l-2,l-2,l-2))\cdot R(\Gamma^{(l,l-2,l-2)}_{-11}).
\end{aligned}$$

Here

$$\mathcal{C}^{(l-2,l-2,l-2)}_{+11}=12\begin{pmatrix} X_{+11}\\ X_{+12}\\ X_{+13}\\ X_{+22}\\ X_{+23}\\ X_{+33}\end{pmatrix},\quad R(\Gamma^{(l-2,l-2,l-2)}_{+11})=12\begin{pmatrix}\nu_1+l+1\\ \nu_2+l\\ \nu_3+l-1\end{pmatrix},$$

$$\mathcal{C}^{(l,l-2,l-2)}_{+22}=2\begin{pmatrix}
X_{+22} & -2X_{+12} & 0 & X_{+11} & 0 & 0\\
X_{+23} & -X_{+13} & -X_{+12} & 0 & X_{+11} & 0\\
X_{+33} & 0 & -2X_{+13} & 0 & 0 & X_{+11}\\
0 & X_{+23} & -X_{+22} & -X_{+13} & X_{+12} & 0\\
0 & X_{+33} & -X_{+23} & 0 & -X_{+13} & X_{+12}\\
0 & 0 & 0 & X_{+33} & -2X_{+23} & X_{+22}
\end{pmatrix},$$

$$R(\Gamma^{(l,l-2,l-2)}_{+22})=2\begin{pmatrix}
\nu_2+l & \nu_1+l-1 & 0\\
\nu_3+l-1 & 0 & \nu_1+l-1\\
0 & \nu_3+l-1 & \nu_2+l-2
\end{pmatrix},$$

$$\begin{aligned}
&\mathcal{C}^{(l,l,l-2)}_{+33}=\begin{pmatrix} X_{+33} & -2X_{+23} & X_{+22} & 2X_{+13} & -2X_{+12} & X_{+11}\end{pmatrix},\\
&R(\Gamma^{(l,l,l-2)}_{+33})=\begin{pmatrix}\nu_3+l-1 & \nu_2+l-2 & \nu_1+l-3\end{pmatrix},
\end{aligned}$$

$$\mathcal{C}^{(l,l,l)}_{-33}=12\begin{pmatrix} X_{-33}\\ -X_{-23}\\ X_{-22}\\ X_{-13}\\ -X_{-12}\\ X_{-11}\end{pmatrix},\quad R(\Gamma^{(l,l,l)}_{-33})=12\begin{pmatrix}\nu_3-l+1\\ \nu_2-l+2\\ \nu_1-l+3\end{pmatrix},$$

$$\mathcal{C}_{-22}^{(l,l,l-2)} = 2\begin{pmatrix} X_{-22} & 2X_{-23} & 0 & X_{-33} & 0 & 0 \\ -X_{-12} & -X_{-13} & 0 & X_{-23} & X_{-33} & 0 \\ 0 & -X_{-12} & -X_{-13} & -X_{-22} & X_{-33} & 0 \\ X_{-11} & 0 & 0 & -2X_{-13} & 0 & X_{-33} \\ 0 & X_{-11} & 0 & X_{-12} & -X_{-13} & X_{-23} \\ 0 & 0 & X_{-11} & 0 & 2X_{-12} & X_{-22} \end{pmatrix},$$

$$R(\Gamma_{-22}^{(l,l,l-2)}) = 2\begin{pmatrix} \nu_2 - l & \nu_3 - l + 1 & 0 \\ \nu_1 - l + 1 & 0 & \nu_3 - l + 1 \\ 0 & \nu_1 - l + 1 & \nu_2 - l + 2 \end{pmatrix},$$

$$\mathcal{C}_{-11}^{(l,l-2,l-2)} = \begin{pmatrix} X_{-11} & 2X_{-12} & 2X_{-13} & X_{-22} & 2X_{-23} & X_{-33} \end{pmatrix},$$

$$R(\Gamma_{-11}^{(l,l-2,l-2)}) = \begin{pmatrix} \nu_1 - l - 1 & \nu_2 - l & \nu_3 - l + 1 \end{pmatrix}.$$

(ii) The elements C_{2i} $(i = 1, 2, 3)$ are represented by the $\mathfrak{p}_\pm$-matrices $\mathcal{C}_{\pm kl}^{\lambda}$ as follows:

$$C_2 = \frac{1}{12}\mathcal{C}_{+33}^{(l,l,l-2)} \cdot \mathcal{C}_{-33}^{(l,l,l)}, \quad C_4 = \frac{1}{192}\mathcal{C}_{+33}^{(l,l,l-2)} \cdot \mathcal{C}_{+22}^{(l,l-2,l-2)} \cdot \mathcal{C}_{-22}^{(l,l,l-2)} \cdot \mathcal{C}_{-33}^{(l,l,l)},$$

$$C_6 = \frac{1}{20736}\mathcal{C}_{+33}^{(l,l,l-2)} \cdot \mathcal{C}_{+22}^{(l,l-2,l-2)} \cdot \mathcal{C}_{+11}^{(l-2,l-2,l-2)} \cdot \mathcal{C}_{-11}^{(l,l-2,l-2)} \cdot \mathcal{C}_{-22}^{(l,l,l-2)} \cdot \mathcal{C}_{-33}^{(l,l,l)}.$$

Proof. From Theorem 5.6 in the case of $(\sigma_1, \sigma_2, \sigma_3) = (0, 0, 0)$, $(1, 1, 1)$, we obtain the assertion by direct computation. □

From above lemma, we obtain differential equations which Whittaker functions satisfy.

Proposition 6.5. *Let σ be a character of $M_{\min}$ such that $(\sigma_1, \sigma_2, \sigma_3) = (0, 0, 0)$ or $(1, 1, 1)$, and T be an element of the space $\mathcal{I}_{\xi,\pi_{(\sigma,\nu)}}$. For $l \equiv \varepsilon_\sigma \mod 2$, we define a function $\phi_{T,l} \in C_\xi^\infty(N_{\min}\backslash G)$ by the equation $\Phi(T, S_{(l,l,l)}(M_l)) = \phi_{T,l} \otimes f(M_l)^*$. Then $\phi_{T,l}$ satisfies following differential equations:*

$$C_{2i}\phi_{T,l} = \chi_{2i,\sigma,\nu,(l,l,l)}\phi_{T,l} \quad (i = 1, 2, 3).$$

Here

$$\chi_{2,\sigma,\nu,(l,l,l)} = \{\nu_1^2 - (l-3)^2\} + \{\nu_2^2 - (l-2)^2\} + \{\nu_3^2 - (l-1)^2\}, \tag{6.1}$$

$$\begin{aligned} \chi_{4,\sigma,\nu,(l,l,l)} = &\{\nu_1^2 - (l-2)^2\}\{\nu_2^2 - (l-2)^2\} + \{\nu_1^2 - (l-2)^2\}\{\nu_3^2 - (l-1)^2\} \\ &+ \{\nu_2^2 - (l-1)^2\}\{\nu_3^2 - (l-1)^2\}, \end{aligned} \tag{6.2}$$

$$\chi_{6,\sigma,\nu,(l,l,l)} = \{\nu_1^2 - (l-1)^2\}\{\nu_2^2 - (l-1)^2\}\{\nu_3^2 - (l-1)^2\}. \tag{6.3}$$

Proof. Let $\chi_{2i,\sigma,\nu,(l,l,l)} (i = 1, 2, 3)$ be the complex numbers defined by the equations (6.1), (6.2) and (6.3). From Lemma 6.4 and $S_{(l,l,l)}(M_l) = s(M_l, M_l)$, we see that

$$C_2 s(M_l, M_l) = \frac{1}{12}s(M_l, M_l)R(\Gamma_{+33}^{(l,l,l-2)}) \cdot R(\Gamma_{-33}^{(l,l,l)}) = \chi_{2,\sigma,\nu,(l,l,l)}s(M_l, M_l).$$

Similarly, we obtain $C_{2i}s(M_l, M_l) = \chi_{2i,\sigma,\nu,(l,l,l)}s(M_l, M_l)$ $(i = 2, 3)$. Since $\phi_{T,l} = T(s(M_l, M_l))$, we obtain the assertion from these equations. □

6.4. **The chilarity operators in the case of** $(\sigma_1, \sigma_2, \sigma_3) \neq (0, 0, 0)$, $(1, 1, 1)$**.** In this subsection, we consider the cases of $(\sigma_1, \sigma_2, \sigma_3) \neq (0, 0, 0)$, $(1, 1, 1)$. Let ε_σ, $\delta_{\sigma;i} \in \{0, 1\}$ $(i = 1, 2, 3)$ be the integers defined by

$$\varepsilon_\sigma = \begin{cases} 0 & \text{if } (\sigma_1, \sigma_2, \sigma_3) = (1, 0, 0),\ (0, 1, 0),\ (0, 0, 1), \\ 1 & \text{if } (\sigma_1, \sigma_2, \sigma_3) = (1, 1, 0),\ (1, 0, 1),\ (0, 1, 1), \end{cases} \quad \text{and } \delta_{\sigma;i} = \begin{cases} 0 & \text{if } \varepsilon_\sigma - \sigma_i = 0, \\ 1 & \text{otherwise}. \end{cases}$$

In the cases of $(\sigma_1, \sigma_2, \sigma_3) \neq (0, 0, 0)$, $(1, 1, 1)$, there are multiplicity one K-types $\tau_{(l+1,l,l)}$ and $\tau_{(l,l,l-1)}$ $(l \equiv \varepsilon_\sigma \mod 2)$ of $\pi_{(\sigma,\nu)}$. Let us compute the values $\chi_{2i,\sigma,\nu,(l+,l,l)}$, $\chi_{2i,\sigma,\nu,(l,l,l-1)}$ $(1 \leq i \leq 3)$.

Lemma 6.6. *(i) Let σ be a character of $M_{\min}$ such that $(\sigma_1, \sigma_2, \sigma_3) \neq (0,0,0),\ (1,1,1)$. For $l \equiv \varepsilon_\sigma \bmod 2$, the following equations hold:*

$$\begin{aligned}
\mathcal{C}_{+22}^{(l+1,l-2,-2)} \mathbf{S}_\sigma((l+1,l-2,-2)) =& \mathbf{S}_\sigma((l+1,l,l-2)) \cdot R(\Gamma_{+22}^{(l+1,l-2,-2)}),\\
\mathcal{C}_{+33}^{(l+1,l,l-2)} \mathbf{S}_\sigma((l+1,l,l-2)) =& \mathbf{S}_\sigma((l+1,l,l)) \cdot R(\Gamma_{+33}^{(l+1,l,l-2)}),\\
\mathcal{C}_{+33}^{(l,l,l-3)} \mathbf{S}_\sigma((l,l,l-3)) =& \mathbf{S}_\sigma((l,l,l-1)) \cdot R(\Gamma_{+33}^{(l,l,l-3)}),\\
\mathcal{C}_{+12}^{(l-1,l-2,l-2)} \mathbf{S}_\sigma((l-1,l-2,l-2)) =& \mathbf{S}_\sigma((l,l-1,l-2)) \cdot R(\Gamma_{+12}^{(l-1,l-2,l-2)}),\\
\mathcal{C}_{+13}^{(l,l,l-1)} \mathbf{S}_\sigma((l,l,l-1)) =& \mathbf{S}_\sigma((l+1,l,l)) \cdot R(\Gamma_{+13}^{(l,l,l-1)}),\\
\mathcal{C}_{+13}^{(l-2,l-2,l-3)} \mathbf{S}_\sigma((l-2,l-2,l-3)) =& \mathbf{S}_\sigma((l-1,l-2,l-2)) \cdot R(\Gamma_{+13}^{(l-2,l-2,l-3)}),\\
\mathcal{C}_{+23}^{(l,l-1,l-2)} \mathbf{S}_\sigma((l,l-1,l-2)) =& \mathbf{S}_\sigma((l,l,l-1)) \cdot R(\Gamma_{+23}^{(l,l-1,l-2)}),\\
\mathcal{C}_{+23}^{(l,l-2,l-3)} \mathbf{S}_\sigma((l,l-2,l-3)) =& \mathbf{S}_\sigma((l,l-1,l-2)) \cdot R(\Gamma_{+23}^{(l,l-2,l-3)}),\\
\mathcal{C}_{-22}^{(l+1,l,l-2)} \mathbf{S}_\sigma((l+1,l,l-2)) =& \mathbf{S}_\sigma((l+1,l-2,-2)) \cdot R(\Gamma_{-22}^{(l+1,l,l-2)}),\\
\mathcal{C}_{-33}^{(l+1,l,l)} \mathbf{S}_\sigma((l+1,l,l)) =& \mathbf{S}_\sigma((l+1,l,l-2)) \cdot R(\Gamma_{-33}^{(l+1,l,l)}),\\
\mathcal{C}_{-33}^{(l,l,l-1)} \mathbf{S}_\sigma((l,l,l-1)) =& \mathbf{S}_\sigma((l,l,l-3)) \cdot R(\Gamma_{-33}^{(l,l,l-1)}),\\
\mathcal{C}_{-12}^{(l,l-1,l-2)} \mathbf{S}_\sigma((l,l-1,l-2)) =& \mathbf{S}_\sigma((l-1,l-2,l-2)) \cdot R(\Gamma_{-12}^{(l,l-1,l-2)}),\\
\mathcal{C}_{-13}^{(l+1,l,l)} \mathbf{S}_\sigma((l+1,l,l)) =& \mathbf{S}_\sigma((l,l,l-1)) \cdot R(\Gamma_{-13}^{(l+1,l,l)}),\\
\mathcal{C}_{-13}^{(l-1,l-2,l-2)} \mathbf{S}_\sigma((l-1,l-2,l-2)) =& \mathbf{S}_\sigma((l-2,l-2,l-3)) \cdot R(\Gamma_{-13}^{(l-1,l-2,l-2)}),\\
\mathcal{C}_{-23}^{(l,l,l-1)} \mathbf{S}_\sigma((l,l,l-1)) =& \mathbf{S}_\sigma((l,l-1,l-2)) \cdot R(\Gamma_{-23}^{(l,l,l-1)}),\\
\mathcal{C}_{-23}^{(l,l-1,l-2)} \mathbf{S}_\sigma((l,l-1,l-2)) =& \mathbf{S}_\sigma((l,l-2,l-3)) \cdot R(\Gamma_{-23}^{(l,l-1,l-2)}).
\end{aligned}$$

The explicit expressions of $\mathcal{C}_{\pm ij}^{\lambda}$ and $R(\Gamma_{\pm ij}^{\lambda})$ in the above equations are given as follows:

$$\mathcal{C}_{+22}^{(l+1,l-2,-2)} =$$
$$2\begin{pmatrix}
X_{+22} & -2X_{+12} & 0 & X_{+11} & 0 & 0 & 0 & 0 & 0 & 0\\
X_{+23} & -X_{+13} & -X_{+12} & 0 & X_{+11} & 0 & 0 & 0 & 0 & 0\\
X_{+33} & 0 & -2X_{+13} & 0 & 0 & X_{+11} & 0 & 0 & 0 & 0\\
0 & X_{+22} & 0 & -2X_{+12} & 0 & 0 & X_{+11} & 0 & 0 & 0\\
0 & X_{+23} & -X_{+22} & -X_{+13} & X_{+12} & 0 & 0 & 0 & 0 & 0\\
0 & 0 & X_{+22} & 0 & -2X_{+12} & 0 & 0 & X_{+11} & 0 & 0\\
0 & X_{+33} & -X_{+23} & 0 & -X_{+13} & X_{+12} & 0 & 0 & 0 & 0\\
0 & 0 & X_{+23} & 0 & -X_{+13} & -X_{+12} & 0 & 0 & X_{+11} & 0\\
0 & 0 & X_{+33} & 0 & 0 & -2X_{+13} & 0 & 0 & 0 & X_{+11}\\
0 & 0 & 0 & X_{+23} & -X_{+22} & 0 & -X_{+13} & X_{+12} & 0 & 0\\
0 & 0 & 0 & X_{+33} & -2X_{+23} & X_{+22} & 0 & 0 & 0 & 0\\
0 & 0 & 0 & 0 & X_{+23} & -X_{+22} & 0 & -X_{+13} & X_{+12} & 0\\
0 & 0 & 0 & 0 & X_{+33} & -X_{+23} & 0 & 0 & -X_{+13} & X_{+12}\\
0 & 0 & 0 & 0 & 0 & 0 & X_{+33} & -2X_{+23} & X_{+22} & 0\\
0 & 0 & 0 & 0 & 0 & 0 & 0 & X_{+33} & -2X_{+23} & X_{+22}
\end{pmatrix},$$

$$\mathcal{C}_{+33}^{(l+1,l,l-2)} = {}^t\begin{pmatrix} X_{+33} & 0 & 0 \\ -2X_{+23} & 0 & 0 \\ X_{+22} & 0 & 0 \\ 0 & X_{+33} & 0 \\ 2X_{+13} & -2X_{+23} & 0 \\ 0 & -2X_{+23} & X_{+33} \\ -2X_{+12} & X_{+22} & 0 \\ 0 & X_{+22} & -2X_{+23} \\ 0 & 0 & X_{+22} \\ 0 & 2X_{+13} & 0 \\ X_{+11} & -2X_{+12} & 0 \\ 0 & -2X_{+12} & 2X_{+13} \\ 0 & 0 & -2X_{+12} \\ 0 & X_{+11} & 0 \\ 0 & 0 & X_{+11} \end{pmatrix},$$

$$\mathcal{C}_{+33}^{(l,l,l-3)} = \begin{pmatrix} X_{+33} & -2X_{+23} & X_{+22} & 0 & 2X_{+13} & -2X_{+12} & 0 & X_{+11} & 0 & 0 \\ 0 & X_{+33} & -2X_{+23} & X_{+22} & 0 & 2X_{+13} & -2X_{+12} & 0 & X_{+11} & 0 \\ 0 & 0 & 0 & 0 & X_{+33} & -2X_{+23} & X_{+22} & 2X_{+13} & -2X_{+12} & X_{+11} \end{pmatrix},$$

$$\mathcal{C}_{+12}^{(l-1,l-2,l-2)} = 3\begin{pmatrix} X_{+12} & -X_{+11} & 0 \\ X_{+13} & 0 & -X_{+11} \\ X_{+22} & -X_{+12} & 0 \\ 0 & X_{+13} & -X_{+12} \\ X_{+23} & -X_{+13} & 0 \\ X_{+33} & 0 & -X_{+13} \\ 0 & X_{+23} & -X_{+22} \\ 0 & X_{+33} & -X_{+23} \end{pmatrix},$$

$$\mathcal{C}_{+13}^{(l,l,l-1)} = \mathcal{C}_{+13}^{(l-2,l-2,l-3)} = 2\begin{pmatrix} -X_{+13} & X_{+12} & -X_{+11} \\ -X_{+23} & X_{+22} & -X_{+12} \\ -X_{+33} & X_{+23} & -X_{+13} \end{pmatrix},$$

$$\mathcal{C}_{+23}^{(l,l-1,l-2)} = \begin{pmatrix} -X_{+23} & X_{+22} & X_{+13} & -2X_{+12} & -X_{+12} & 0 & X_{+11} & 0 \\ -X_{+33} & X_{+23} & 0 & -X_{+13} & X_{+13} & -X_{+12} & 0 & X_{+11} \\ 0 & 0 & -X_{+33} & X_{+23} & 2X_{+23} & -X_{+22} & -X_{+13} & X_{+12} \end{pmatrix},$$

$$\mathcal{C}_{+23}^{(l,l-2,l-3)} = {}^t\begin{pmatrix} -X_{+23} & -X_{+33} & 0 & 0 & 0 & 0 & 0 & 0 \\ X_{+22} & X_{+23} & 0 & 0 & 0 & 0 & 0 & 0 \\ X_{+13} & 0 & -X_{+23} & -X_{+33} & 0 & 0 & 0 & 0 \\ -2X_{+12} & -X_{+13} & X_{+22} & X_{+23} & 0 & 0 & 0 & 0 \\ -X_{+12} & X_{+13} & X_{+22} & 2X_{+23} & -X_{+23} & -X_{+33} & 0 & 0 \\ 0 & -X_{+12} & 0 & -X_{+22} & X_{+22} & X_{+23} & 0 & 0 \\ 0 & 0 & X_{+13} & 0 & 0 & 0 & -X_{+33} & 0 \\ X_{+11} & 0 & -2X_{+12} & -X_{+13} & 0 & 0 & X_{+23} & 0 \\ 0 & 0 & -X_{+12} & 0 & X_{+13} & 0 & 2X_{+23} & -X_{+33} \\ 0 & X_{+11} & 0 & X_{+12} & -2X_{+12} & -X_{+13} & -X_{+22} & X_{+23} \\ 0 & 0 & 0 & 0 & -X_{+12} & X_{+13} & -X_{+22} & 2X_{+23} \\ 0 & 0 & 0 & 0 & 0 & -X_{+12} & 0 & -X_{+22} \\ 0 & 0 & X_{+11} & 0 & 0 & 0 & -X_{+13} & 0 \\ 0 & 0 & 0 & 0 & X_{+11} & 0 & X_{+12} & -X_{+13} \\ 0 & 0 & 0 & 0 & 0 & X_{+11} & 0 & X_{+12} \end{pmatrix},$$

$$\mathcal{C}_{-22}^{(l+1,l,l-2)} =$$

$$2\ {}^t\!\begin{pmatrix} 3X_{-22} & -2X_{-12} & 0 & X_{-11} & 0 & 0 & 0 & 0 & 0 & 0 \\ 6X_{-23} & -2X_{-13} & -2X_{-12} & 0 & X_{-11} & 0 & 0 & 0 & 0 & 0 \\ 3X_{-33} & 0 & -2X_{-13} & 0 & 0 & X_{-11} & 0 & 0 & 0 & 0 \\ 0 & X_{-22} & 0 & -2X_{-12} & 0 & 0 & 3X_{-11} & 0 & 0 & 0 \\ 0 & 4X_{-23} & -2X_{-22} & -4X_{-13} & 0 & 0 & 0 & 2X_{-11} & 0 & 0 \\ 0 & 2X_{-23} & X_{-22} & -2X_{-13} & -2X_{-12} & 0 & 0 & 3X_{-11} & 0 & 0 \\ 0 & 3X_{-33} & -2X_{-23} & 0 & -2X_{-13} & 2X_{-12} & 0 & 0 & X_{-11} & 0 \\ 0 & X_{-33} & 2X_{-23} & 0 & -2X_{-13} & -2X_{-12} & 0 & 0 & 3X_{-11} & 0 \\ 0 & 0 & X_{-33} & 0 & 0 & -2X_{-13} & 0 & 0 & 0 & 3X_{-11} \\ 0 & 0 & 0 & 2X_{-23} & -1X_{-22} & 0 & -6X_{-13} & 2X_{-12} & 0 & 0 \\ 0 & 0 & 0 & 3X_{-33} & -2X_{-23} & X_{-22} & 0 & -2X_{-13} & 2X_{-12} & 0 \\ 0 & 0 & 0 & 2X_{-33} & 0 & -2X_{-22} & 0 & -4X_{-13} & 4X_{-12} & 0 \\ 0 & 0 & 0 & 0 & X_{-33} & -2X_{-23} & 0 & 0 & -2X_{-13} & 6X_{-12} \\ 0 & 0 & 0 & 0 & 0 & 0 & 3X_{-33} & -2X_{-23} & X_{-22} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & X_{-33} & -2X_{-23} & 3X_{-22} \end{pmatrix},$$

$$\mathcal{C}_{-33}^{(l+1,l,l)} = 6\begin{pmatrix} 4X_{-33} & 0 & 0 \\ -4X_{-23} & 0 & 0 \\ 4X_{-22} & 0 & 0 \\ 0 & 4X_{-33} & 0 \\ 3X_{-13} & -X_{-23} & -X_{-33} \\ -2X_{-13} & -2X_{-23} & 2X_{-33} \\ -3X_{-12} & X_{-22} & X_{-23} \\ X_{-12} & X_{-22} & -3X_{-23} \\ 0 & 0 & 4X_{-22} \\ 0 & 4X_{-13} & 0 \\ 2X_{-11} & -2X_{-12} & -2X_{-13} \\ -X_{-11} & -X_{-12} & 3X_{-13} \\ 0 & 0 & -4X_{-12} \\ 0 & 4X_{-11} & 0 \\ 0 & 0 & 4X_{-11} \end{pmatrix},$$

$$\mathcal{C}_{-33}^{(l,l,l-1)} = 24\begin{pmatrix} 3X_{-33} & 0 & 0 \\ -2X_{-23} & X_{-33} & 0 \\ X_{-22} & -2X_{-23} & 0 \\ 0 & 3X_{-22} & 0 \\ 2X_{-13} & 0 & X_{-33} \\ -X_{-12} & X_{-13} & -X_{-23} \\ 0 & -2X_{-12} & X_{-22} \\ X_{-11} & 0 & 2X_{-13} \\ 0 & X_{-11} & -2X_{-12} \\ 0 & 0 & 3X_{-11} \end{pmatrix},$$

$$\mathcal{C}_{-12}^{(l,l-1,l-2)} = \begin{pmatrix} -X_{-12} & -X_{-13} & -X_{-22} & -X_{-23} & -2X_{-23} & -X_{-33} & 0 & 0 \\ X_{-11} & 0 & X_{-12} & -X_{-13} & X_{-13} & 0 & -X_{-23} & -X_{-33} \\ 0 & X_{-11} & 0 & 2X_{-12} & X_{-12} & X_{-13} & X_{-22} & X_{-23} \end{pmatrix},$$

$$\mathcal{C}_{-13}^{(l+1,l,l)} = \mathcal{C}_{-13}^{(l-1,l-2,l-2)} = 2\begin{pmatrix} -X_{-13} & -X_{-23} & -X_{-33} \\ X_{-12} & X_{-22} & X_{-23} \\ -X_{-11} & -X_{-12} & -X_{-13} \end{pmatrix},$$

$$\mathcal{C}_{-23}^{(l,l,l-1)} = 3\begin{pmatrix} X_{-23} & X_{-33} & 0 \\ -X_{-22} & -X_{-23} & 0 \\ -X_{-13} & 0 & X_{-33} \\ X_{-12} & X_{-13} & 0 \\ 0 & -X_{-13} & -X_{-23} \\ 0 & X_{-12} & X_{-22} \\ -X_{-11} & 0 & X_{-13} \\ 0 & -X_{-11} & -X_{-12} \end{pmatrix},$$

$$\mathcal{C}_{-23}^{(l,l-1,l-2)} =$$

$$\begin{pmatrix} 8X_{-23} & 8X_{-33} & 0 & 0 & 0 & 0 & 0 & 0 \\ -8X_{-22} & -8X_{-23} & 0 & 0 & 0 & 0 & 0 & 0 \\ -4X_{-13} & 0 & 4X_{-23} & 8X_{-33} & 4X_{-33} & 0 & 0 & 0 \\ 6X_{-12} & 6X_{-13} & -2X_{-22} & -2X_{-23} & -4X_{-23} & -2X_{-33} & 0 & 0 \\ -X_{-12} & -5X_{-13} & -X_{-22} & -5X_{-23} & 2X_{-23} & 3X_{-33} & 0 & 0 \\ 0 & 4X_{-12} & 0 & 4X_{-22} & -4X_{-22} & -4X_{-23} & 0 & 0 \\ 0 & 0 & -8X_{-13} & 0 & 0 & 0 & 8X_{-33} & 0 \\ -3X_{-11} & 0 & 5X_{-12} & 7X_{-13} & 5X_{-13} & 0 & -X_{-23} & -X_{-33} \\ X_{-11} & 0 & X_{-12} & -5X_{-13} & -7X_{-13} & 0 & -5X_{-23} & 3X_{-33} \\ 0 & -3X_{-11} & 0 & -2X_{-12} & 5X_{-12} & 5X_{-13} & X_{-22} & X_{-23} \\ 0 & 2X_{-11} & 0 & 4X_{-12} & 2X_{-12} & -6X_{-13} & 2X_{-22} & -6X_{-23} \\ 0 & 0 & 0 & 0 & 0 & 8X_{-12} & 0 & 8X_{-22} \\ 0 & 0 & -8X_{-11} & 0 & 0 & 0 & 8X_{-13} & 0 \\ 0 & 0 & 0 & -4X_{-11} & -8X_{-11} & 0 & -4X_{-12} & 4X_{-13} \\ 0 & 0 & 0 & 0 & 0 & -8X_{-11} & 0 & -8X_{-12} \end{pmatrix}.$$

- *the case of* $(\sigma_1,\sigma_2,\sigma_3) = (1,0,0),\ (0,1,1)$.

$$R(\Gamma_{+22}^{(l+1,l-2,-2)}) = 2\begin{pmatrix} \nu_2+l & \nu_1+l-2 & 0 \\ \nu_3+l-1 & 0 & \nu_1+l-2 \\ 0 & \nu_3+l-1 & \nu_2+l-2 \\ 0 & 0 & -(\nu_2+l-1) \end{pmatrix},$$

$$R(\Gamma_{+33}^{(l+1,l,l-2)}) = \begin{pmatrix} \nu_3+l-1 & \nu_2+l-2 & \nu_1+l-4 & 0 \end{pmatrix},$$

$$R(\Gamma_{+33}^{(l,l,l-3)}) = \begin{pmatrix} \nu_3+l-1 & \nu_2+l-2 & \nu_1+l-4 \end{pmatrix},$$

$$R(\Gamma_{+12}^{(l-1,l-2,l-2)}) = 3\begin{pmatrix} \nu_2+l \\ \nu_3+l-1 \end{pmatrix}, \quad R(\Gamma_{+13}^{(l,l,l-1)}) = -2(\nu_1+l),$$

$$R(\Gamma_{+13}^{(l-2,l-2,l-3)}) = -2(\nu_1+l-2), \quad R(\Gamma_{+23}^{(l,l-1,l-2)}) = -\begin{pmatrix} \nu_3+l-1 & \nu_2+l-2 \end{pmatrix},$$

$$R(\Gamma_{+23}^{(l,l-2,l-3)}) = \begin{pmatrix} \nu_2+l-1 & \nu_2+l-2 & \nu_1+l-3 & 0 \\ 0 & -(\nu_3+l-1) & 0 & \nu_1+l-3 \end{pmatrix},$$

$$R(\Gamma_{-22}^{(l+1,l,l-2)}) = 2\begin{pmatrix} 3(\nu_2-l) & 3(\nu_3-l+1) & 0 & 0 \\ \nu_1-l+2 & 0 & 3(\nu_3-l+1) & 2(\nu_3-l+1) \\ 0 & \nu_1-l+2 & \nu_2-l+4 & -2(\nu_2-l) \end{pmatrix},$$

$$R(\Gamma_{-33}^{(l+1,l,l)}) = 6\begin{pmatrix} 4(\nu_3-l+1) \\ 4(\nu_2-l+2) \\ 2(\nu_1-l+4) \\ -(\nu_1-l+4) \end{pmatrix}, \quad R(\Gamma_{-33}^{(l,l,l-1)}) = 24\begin{pmatrix} \nu_3-l+1 \\ \nu_2-l+2 \\ 3(\nu_r-l+4) \end{pmatrix},$$

$$R(\Gamma_{-12}^{(l,l-1,l-2)}) = -\begin{pmatrix} \nu_2-l & \nu_3-l+1 \end{pmatrix}, \quad R(\Gamma_{-13}^{(l+1,l,l)}) = -2(\nu_1-l),$$

$$R(\Gamma^{(l-1,l-2,l-2)}_{-13}) = -2(\nu_1 - l + 2), \quad R(\Gamma^{(l,l,l-1)}_{-23}) = 3\begin{pmatrix} \nu_3 - l + 1 \\ \nu_2 - l + 2 \end{pmatrix},$$

$$R(\Gamma^{(l,l-1,l-2)}_{-23}) = \begin{pmatrix} -2(\nu_2 - l) & -2(\nu_3 - l + 1) \\ -(\nu_2 - l + 4) & 3(\nu_3 - l + 1) \\ -8(\nu_1 - l + 3) & 0 \\ 0 & -8(\nu_1 - l + 3) \end{pmatrix}.$$

- *the case of* $(\sigma_1, \sigma_2, \sigma_3) = (0,1,0),\ (1,0,1)$.

$$R(\Gamma^{(l+1,l-2,-2)}_{+22}) = 2\begin{pmatrix} \nu_2 + l + 1 & \nu_1 + l - 1 & 0 \\ \nu_3 + l - 1 & 0 & 0 \\ 0 & 0 & \nu_1 + l - 1 \\ 0 & \nu_3 + l - 1 & \nu_2 + l - 3 \end{pmatrix},$$

$$R(\Gamma^{(l+1,l,l-2)}_{+33}) = \begin{pmatrix} \nu_3 + l - 1 & \nu_2 + l - 1 & \nu_2 + l - 3 & \nu_1 + l - 3 \end{pmatrix},$$

$$R(\Gamma^{(l,l,l-3)}_{+33}) = \begin{pmatrix} \nu_3 + l - 1 & \nu_2 + l - 3 & \nu_1 + l - 5 \end{pmatrix},$$

$$R(\Gamma^{(l-1,l-2,l-2)}_{+12}) = 3\begin{pmatrix} -(\nu_1 + l) \\ \nu_3 + l - 1 \end{pmatrix}, \quad R(\Gamma^{(l,l,l-1)}_{+13}) = 2(\nu_2 + l),$$

$$R(\Gamma^{(l-2,l-2,l-3)}_{+13}) = 2(\nu_2 + l - 2), \quad R(\Gamma^{(l,l-1,l-2)}_{+23}) = \begin{pmatrix} -(\nu_3 + l - 1) & \nu_1 + l - 2 \end{pmatrix},$$

$$R(\Gamma^{(l,l-2,l-3)}_{+23}) = \begin{pmatrix} \nu_2 + l - 2 & \nu_1 + l - 4 & 0 & 0 \\ 0 & 0 & -(\nu_3 + l - 1) & -(\nu_2 + l - 3) \end{pmatrix},$$

$$R(\Gamma^{(l+1,l,l-2)}_{-22}) = 2\begin{pmatrix} \nu_2 - l - 1 & 3(\nu_3 - l + 1) & \nu_3 - l + 1 & 0 \\ 3(\nu_1 - l + 1) & 0 & 0 & 3(\nu_3 - l + 1) \\ 0 & \nu_1 - l + 1 & 3(\nu_1 - l + 1) & (\nu_2 - l + 3) \end{pmatrix},$$

$$R(\Gamma^{(l+1,l,l)}_{-33}) = 6\begin{pmatrix} 4(\nu_3 - l + 1) \\ (\nu_2 - l) \\ (\nu_2 - l + 4) \\ 4(\nu_1 - l + 3) \end{pmatrix}, \quad R(\Gamma^{(l,l,l-1)}_{-33}) = 24\begin{pmatrix} \nu_3 - l + 1 \\ 3(\nu_2 - l + 3) \\ \nu_1 - l + 5 \end{pmatrix},$$

$$R(\Gamma^{(l,l-1,l-2)}_{-12}) = \begin{pmatrix} \nu_1 - l & -(\nu_3 - l + 1) \end{pmatrix}, \quad R(\Gamma^{(l+1,l,l)}_{-13}) = 2(\nu_2 - l),$$

$$R(\Gamma^{(l-1,l-2,l-2)}_{-13}) = 2(\nu_2 - l + 2), \quad R(\Gamma^{(l,l,l-1)}_{-23}) = 3\begin{pmatrix} \nu_3 - l + 1 \\ -(\nu_1 - l + 2) \end{pmatrix},$$

$$R(\Gamma^{(l,l-1,l-2)}_{-23}) = \begin{pmatrix} -8(\nu_2 - l + 2) & 0 \\ -3(\nu_1 - l + 4) & -(\nu_3 - l + 1) \\ \nu_1 - l + 4 & 3(\nu_3 - l + 1) \\ 0 & 8(\nu_2 - l + 3) \end{pmatrix}.$$

- *the case of* $(\sigma_1, \sigma_2, \sigma_3) = (0,0,1),\ (1,1,0)$.

$$R(\Gamma^{(l+1,l-2,-2)}_{+22}) = 2\begin{pmatrix} -(\nu_2 + l - 1) & 0 & 0 \\ \nu_2 + l & \nu_1 + l - 1 & 0 \\ \nu_3 + l & 0 & \nu_1 + l - 1 \\ 0 & \nu_3 + l & \nu_2 + l - 2 \end{pmatrix},$$

$$R(\Gamma^{(l+1,l,l-2)}_{+33}) = \begin{pmatrix} 0 & \nu_3 + l & \nu_2 + l - 2 & \nu_1 + l - 3 \end{pmatrix},$$

$$R(\Gamma^{(l,l,l-3)}_{+33}) = \begin{pmatrix} \nu_3 + l - 2 & \nu_2 + l - 4 & \nu_1 + l - 5 \end{pmatrix},$$

$$R(\Gamma^{(l-1,l-2,l-2)}_{+12}) = -3\begin{pmatrix} \nu_1 + l \\ \nu_2 + l - 1 \end{pmatrix}, \quad R(\Gamma^{(l,l,l-1)}_{+13}) = -2(\nu_3 + l),$$

$$R(\Gamma^{(l-2,l-2,l-3)}_{+13}) = -2(\nu_3 + l - 2), \quad R(\Gamma^{(l,l-1,l-2)}_{+23}) = \begin{pmatrix} \nu_2 + l - 1 & \nu_1 + l - 2 \end{pmatrix},$$

$$R(\Gamma_{+23}^{(l,l-2,l-3)}) = \begin{pmatrix} -(\nu_3+l-2) & 0 & \nu_1+l-4 & 0 \\ 0 & -(\nu_3+l-2) & -(\nu_2+l-3) & -(\nu_2+l-4) \end{pmatrix},$$

$$R(\Gamma_{-22}^{(l+1,l,l-2)}) = 2\begin{pmatrix} -2(\nu_2-l+2) & \nu_2-l-2 & \nu_3-l & 0 \\ 2(\nu_1-l+1) & 3(\nu_1-l+1) & 0 & \nu_3-l \\ 0 & 0 & 3(\nu_1-l+1) & 3(\nu_2-l+2) \end{pmatrix},$$

$$R(\Gamma_{-33}^{(l+1,l,l)}) = 6\begin{pmatrix} -(\nu_3-l) \\ 2(\nu_3-l) \\ 4(\nu_2-l+2) \\ 4(\nu_1-l+3) \end{pmatrix}, \quad R(\Gamma_{-33}^{(l,l,l-1)}) = 24\begin{pmatrix} 3(\nu_3-l+2) \\ \nu_2-l+4 \\ \nu_1-l+5 \end{pmatrix},$$

$$R(\Gamma_{-12}^{(l,l-1,l-2)}) = \begin{pmatrix} \nu_1-l & \nu_2-l+1 \end{pmatrix}, \quad R(\Gamma_{-13}^{(l+1,l,l)}) = -2(\nu_3-l),$$

$$R(\Gamma_{-13}^{(l-1,l-2,l-2)}) = -2(\nu_3-l+2), \quad R(\Gamma_{-23}^{(l,l,l-1)}) = -3\begin{pmatrix} \nu_2-l+1 \\ \nu_1-l+2 \end{pmatrix},$$

$$R(\Gamma_{-23}^{(l,l-1,l-2)}) = \begin{pmatrix} 8(\nu_3-l+2) & 0 \\ 0 & 8(\nu_3-l+2) \\ -3(\nu_1-l+4) & \nu_2-l+1 \\ 2(\nu_1-l+4) & 2(\nu_2-l+5) \end{pmatrix}.$$

(ii) The elements C_{2i} $(i=1,2,3)$ are represented by the $\mathfrak{p}_\pm$-matrices $\mathcal{C}_{\pm kl}^{\lambda}$ as follows:

$$\begin{aligned}
\begin{pmatrix} C_2 & 0 & 0 \\ 0 & C_2 & 0 \\ 0 & 0 & C_2 \end{pmatrix} &= \frac{1}{24}\Big\{\mathcal{C}_{+33}^{(l+1,l,l-2)}\cdot\mathcal{C}_{-33}^{(l+1,l,l)} + 3\,\mathcal{C}_{+13}^{(l,l,l-1)}\cdot\mathcal{C}_{-13}^{(l+1,l,l)}\Big\} \\
&= \frac{1}{72}\Big\{\mathcal{C}_{+33}^{(l,l,l-3)}\cdot\mathcal{C}_{-33}^{(l,l,l-1)} - 16\,\mathcal{C}_{+23}^{(l,l-1,l-2)}\cdot\mathcal{C}_{-23}^{(l,l,l-1)}\Big\},
\end{aligned}$$

$$\begin{aligned}
\begin{pmatrix} C_4 & 0 & 0 \\ 0 & C_4 & 0 \\ 0 & 0 & C_4 \end{pmatrix} &= \frac{1}{1152}\Big\{\mathcal{C}_{+33}^{(l+1,l,l-2)}\cdot\mathcal{C}_{+22}^{(l+1,l-2,-2)}\cdot\mathcal{C}_{-22}^{(l+1,l,l-2)}\cdot\mathcal{C}_{-33}^{(l+1,l,l)} \\
&\qquad - 64\,\mathcal{C}_{+13}^{(l,l,l-1)}\cdot\mathcal{C}_{+23}^{(l,l-1,l-2)}\cdot\mathcal{C}_{-23}^{(l,l,l-1)}\cdot\mathcal{C}_{-13}^{(l+1,l,l)}\Big\} \\
&= \frac{1}{72}\Big\{\mathcal{C}_{+23}^{(l,l-1,l-2)}\cdot\mathcal{C}_{+12}^{(l-1,l-2,l-2)}\cdot\mathcal{C}_{-12}^{(l,l-1,l-2)}\cdot\mathcal{C}_{-23}^{(l,l,l-1)} \\
&\qquad + 3\,\mathcal{C}_{+23}^{(l,l-1,l-2)}\cdot\mathcal{C}_{+23}^{(l,l-2,l-3)}\cdot\mathcal{C}_{-23}^{(l,l-1,l-2)}\cdot\mathcal{C}_{-23}^{(l,l,l-1)}\Big\},
\end{aligned}$$

$$\begin{aligned}
\begin{pmatrix} C_6 & 0 & 0 \\ 0 & C_6 & 0 \\ 0 & 0 & C_6 \end{pmatrix} &= \frac{1}{144}\Big\{\mathcal{C}_{+13}^{(l,l,l-1)}\cdot\mathcal{C}_{+23}^{(l,l-1,l-2)}\cdot\mathcal{C}_{+12}^{(l-1,l-2,l-2)}\cdot\mathcal{C}_{-12}^{(l,l-1,l-2)}\cdot\mathcal{C}_{-23}^{(l,l,l-1)}\cdot\mathcal{C}_{-13}^{(l+1,l,l)}\Big\} \\
&= \frac{1}{144}\Big\{\mathcal{C}_{+23}^{(l,l-1,l-2)}\cdot\mathcal{C}_{+12}^{(l-1,l-2,l-2)}\cdot\mathcal{C}_{+13}^{(l-2,l-2,l-3)} \\
&\qquad \times\mathcal{C}_{-13}^{(l-1,l-2,l-2)}\cdot\mathcal{C}_{-12}^{(l,l-1,l-2)}\cdot\mathcal{C}_{-23}^{(l,l,l-1)}\Big\}.
\end{aligned}$$

Proof. From Theorem 5.6 in the case of $(\sigma_1,\sigma_2,\sigma_3)\neq(0,0,0),\ (1,1,1)$, we obtain the assertion by direct computation. □

From above lemma, we obtain differential equations which Whittaker functions satisfy. For $l\in\mathbf{Z}$, we set

$$M_{l;1}^{(1)} = M_l\left(\begin{smallmatrix}1&0&0\\&1&0\\&&1\end{smallmatrix}\right), \qquad M_{l;2}^{(1)} = M_l\left(\begin{smallmatrix}1&0&0\\&1&0\\&&0\end{smallmatrix}\right), \qquad M_{l;3}^{(1)} = M_l\left(\begin{smallmatrix}1&0&0\\&0&0\\&&0\end{smallmatrix}\right),$$

$$M_{l;1}^{(2)} = M_l\left(\begin{smallmatrix}0&0&-1\\&0&0\\&&0\end{smallmatrix}\right), \qquad M_{l;2}^{(2)} = M_l\left(\begin{smallmatrix}0&0&-1\\&0&-1\\&&0\end{smallmatrix}\right), \qquad M_{l;3}^{(2)} = M_l\left(\begin{smallmatrix}0&0&-1\\&0&-1\\&&-1\end{smallmatrix}\right).$$

Then $G((l+1,l,l)) = \{M^{(1)}_{l;j} \mid j=1,2,3\}$ and $G((l,l,l-1)) = \{M^{(2)}_{l;j} \mid j=1,2,3\}$. For $l \equiv \varepsilon_\sigma \mod 2$, let $M_l^{(\sigma;1)}$ and $M_l^{(\sigma;2)}$ be the unique element of $G_\sigma((l+1,l,l))$ and $G_\sigma((l,l,l-1))$, respectively.

Proposition 6.7. *Let σ be a character of $M_{\min}$ such that $(\sigma_1,\sigma_2,\sigma_3)$ is neither $(0,0,0)$ nor $(1,1,1)$, and T be an element of the space $\mathcal{I}_{\xi,\pi_{(\sigma,\nu)}}$.*

(i) For $l \equiv \varepsilon_\sigma \mod 2$, we define functions $\phi^{(1)}_{T,l;j} \in C^\infty_\xi(N_{\min}\backslash G)$ $(j=1,2,3)$ by the equation $\Phi(T, S_{(l+1,l,l)}(M_l^{(\sigma;1)})) = \sum_{j=1}^3 \phi^{(1)}_{T,l;j} \otimes f(M^{(1)}_{l;j})^*$. Then $\phi^{(1)}_{T,l;1}$, $\phi^{(1)}_{T,l;2}$ and $\phi^{(1)}_{T,l;3}$ satisfy following differential equations:

$$\begin{aligned}
&C_{2i}\phi^{(1)}_{T,l;j} = \chi_{2i,\sigma,\nu,(l+1,l,l)}\phi^{(1)}_{T,l;j},\\
&D^{(+,-)}_{i1}\phi^{(1)}_{T,l;1} + D^{(+,-)}_{i2}\phi^{(1)}_{T,l;2} + D^{(+,-)}_{i3}\phi^{(1)}_{T,l;3} = \tilde{\chi}_{2,\sigma,\nu,(l+1,l,l)}\phi^{(1)}_{T,l;i}
\end{aligned}$$

for $i,j=1,2,3$. Here

$$\begin{pmatrix} D^{(+,-)}_{11} & D^{(+,-)}_{12} & D^{(+,-)}_{13} \\ D^{(+,-)}_{21} & D^{(+,-)}_{22} & D^{(+,-)}_{23} \\ D^{(+,-)}_{31} & D^{(+,-)}_{32} & D^{(+,-)}_{33} \end{pmatrix} = m_1(C_+)m_1(C_-),$$

$$\begin{aligned}
\chi_{2,\sigma,\nu,(l+1,l,l)} =& \{\nu_1^2-(l-3)^2\} + \{\nu_2^2-(l-2)^2\} + \{\nu_3^2-(l-1)^2\} - 2l + 1,\\
\chi_{4,\sigma,\nu,(l+1,l,l)} =& \{\nu_1^2-(l-2)^2-(2l-1)\delta_{\sigma;1}\}\{\nu_2^2-(l-2)^2-(2l-1)\delta_{\sigma;2}\}\\
&+ \{\nu_1^2-(l-2)^2-(2l-1)\delta_{\sigma;1}\}\{\nu_3^2-(l-1)^2-(2l-1)\delta_{\sigma;3}\}\\
&+ \{\nu_2^2-(l-1)^2-(2l-1)\delta_{\sigma;2}\}\{\nu_3^2-(l-1)^2-(2l-1)\delta_{\sigma;3}\},\\
\chi_{6,\sigma,\nu,(l+1,l,l)} =& \{\nu_1^2-(l-1)^2-(2l-1)\delta_{\sigma;1}\}\{\nu_2^2-(l-1)^2-(2l-1)\delta_{\sigma;2}\}\\
&\times \{\nu_3^2-(l-1)^2-(2l-1)\delta_{\sigma;3}\},\\
\tilde{\chi}_{2,\sigma,\nu,(l+1,l,l)} =& \nu_1^2\delta_{\sigma;1} + \nu_2^2\delta_{\sigma;2} + \nu_3^2\delta_{\sigma;3} - l^2.
\end{aligned}$$

(ii) For $l \equiv \varepsilon_\sigma \mod 2$, we define functions $\phi^{(2)}_{T,l;j} \in C^\infty_\xi(N_{\min}\backslash G)$ $(j=1,2,3)$ by the equation $\Phi(T, S_{(l,l,l-1)}(M_l^{(\sigma;2)})) = \sum_{j=1}^3 \phi^{(2)}_{T,l;j} \otimes f(M^{(2)}_{l;j})^*$. Then $\phi^{(2)}_{T,l;1}$, $\phi^{(2)}_{T,l;2}$ and $\phi^{(2)}_{T,l;3}$ satisfy following differential equations:

$$\begin{aligned}
&C_{2i}\phi^{(2)}_{T,l;j} = \chi_{2i,\sigma,\nu,(l,l,l-1)}\phi^{(2)}_{T,l;j},\\
&-D^{(-,+)}_{i3}\phi^{(2)}_{T,l;1} + D^{(-,+)}_{i2}\phi^{(2)}_{T,l;2} - D^{(-,+)}_{i1}\phi^{(2)}_{T,l;3} = (-1)^i\tilde{\chi}_{2,\sigma,\nu,(l,l,l-1)}\phi^{(2)}_{T,l;4-i}
\end{aligned}$$

for $i,j=1,2,3$. Here

$$\begin{pmatrix} D^{(-,+)}_{11} & D^{(-,+)}_{12} & D^{(-,+)}_{13} \\ D^{(-,+)}_{21} & D^{(-,+)}_{22} & D^{(-,+)}_{23} \\ D^{(-,+)}_{31} & D^{(-,+)}_{32} & D^{(-,+)}_{33} \end{pmatrix} = m_1(C_-)m_1(C_+),$$

$$\begin{aligned}
\chi_{2,\sigma,\nu,(l,l,l-1)} =& \{\nu_1^2-(l-3)^2\} + \{\nu_2^2-(l-2)^2\} + \{\nu_3^2-(l-1)^2\} + 2l - 7,\\
\chi_{4,\sigma,\nu,(l,l,l-1)} =& \{\nu_1^2-(l-2)^2+(2l-5)\delta_{\sigma;1}\}\{\nu_2^2-(l-2)^2+(2l-5)\delta_{\sigma;2}\}\\
&+ \{\nu_1^2-(l-2)^2+(2l-5)\delta_{\sigma;1}\}\{\nu_3^2-(l-1)^2+(2l-5)\delta_{\sigma;3}\}\\
&+ \{\nu_2^2-(l-1)^2+(2l-5)\delta_{\sigma;2}\}\{\nu_3^2-(l-1)^2+(2l-5)\delta_{\sigma;3}\},\\
\chi_{6,\sigma,\nu,(l,l,l-1)} =& \{\nu_1^2-(l-1)^2+(2l-3)\delta_{\sigma;1}\}\{\nu_2^2-(l-1)^2+(2l-3)\delta_{\sigma;2}\}\\
&\times \{\nu_3^2-(l-1)^2+(2l-3)\delta_{\sigma;3}\},\\
\tilde{\chi}_{2,\sigma,\nu,(l,l,l-1)} =& \nu_1^2\delta_{\sigma;1} + \nu_2^2\delta_{\sigma;2} + \nu_3^2\delta_{\sigma;3} - l^2.
\end{aligned}$$

Proof. From the Lemma 6.6 and the equations

$$m_1(C_+)m_1(C_-)=\frac{1}{4}\mathcal{C}^{(l,l,l-1)}_{+13}\mathcal{C}^{(l+1,l,l)}_{-13},$$

$$m_1(C_-)m_1(C_+)\begin{pmatrix}0&0&-1\\0&1&0\\-1&0&0\end{pmatrix}=\frac{1}{4}\begin{pmatrix}0&0&-1\\0&1&0\\-1&0&0\end{pmatrix}\mathcal{C}^{(l+1,l,l)}_{-13}\mathcal{C}^{(l,l,l-1)}_{+13},$$

$$S_{(l+1,l,l)}(M^{(\sigma;1)}_l)=\begin{pmatrix}s(M^{(\sigma;1)}_l,M^{(1)}_{l;1})\\ s(M^{(\sigma;1)}_l,M^{(1)}_{l;2})\\ s(M^{(\sigma;1)}_l,M^{(1)}_{l;3})\end{pmatrix},\quad S_{(l,l,l-1)}(M^{(\sigma;2)}_l)=\begin{pmatrix}s(M^{(\sigma;2)}_l,M^{(2)}_{l;1})\\ s(M^{(\sigma;2)}_l,M^{(2)}_{l;2})\\ s(M^{(\sigma;2)}_l,M^{(2)}_{l;3})\end{pmatrix},$$

we obtain

$$C_{2i}\phi^{(1)}_{T,l;j}=\chi_{2i,\sigma,\nu,(l+1,l,l)}\phi^{(1)}_{T,l;j},\quad C_{2i}\phi^{(2)}_{T,l;j}=\chi_{2i,\sigma,\nu,(l,l,l-1)}\phi^{(2)}_{T,l;j}\quad(i,j=1,2,3)$$

$$\begin{pmatrix}D^{(+,-)}_{11}&D^{(+,-)}_{12}&D^{(+,-)}_{13}\\D^{(+,-)}_{21}&D^{(+,-)}_{22}&D^{(+,-)}_{23}\\D^{(+,-)}_{31}&D^{(+,-)}_{32}&D^{(+,-)}_{33}\end{pmatrix}\begin{pmatrix}s(M^{(\sigma;1)}_l,M^{(1)}_{l;1})\\ s(M^{(\sigma;1)}_l,M^{(1)}_{l;2})\\ s(M^{(\sigma;1)}_l,M^{(1)}_{l;3})\end{pmatrix}=\tilde{\chi}_{2,\sigma,\nu,(l+1,l,l)}\begin{pmatrix}s(M^{(\sigma;1)}_l,M^{(1)}_{l;1})\\ s(M^{(\sigma;1)}_l,M^{(1)}_{l;2})\\ s(M^{(\sigma;1)}_l,M^{(1)}_{l;3})\end{pmatrix},$$

$$\begin{pmatrix}D^{(-,+)}_{11}&D^{(-,+)}_{12}&D^{(-,+)}_{13}\\D^{(-,+)}_{21}&D^{(-,+)}_{22}&D^{(-,+)}_{23}\\D^{(-,+)}_{31}&D^{(-,+)}_{32}&D^{(-,+)}_{33}\end{pmatrix}\begin{pmatrix}-s(M^{(\sigma;2)}_l,M^{(2)}_{l;3})\\ s(M^{(\sigma;2)}_l,M^{(2)}_{l;2})\\ -s(M^{(\sigma;2)}_l,M^{(2)}_{l;1})\end{pmatrix}=\tilde{\chi}_{2,\sigma,\nu,(l,l,l-1)}\begin{pmatrix}-s(M^{(\sigma;2)}_l,M^{(2)}_{l;3})\\ s(M^{(\sigma;2)}_l,M^{(2)}_{l;2})\\ -s(M^{(\sigma;2)}_l,M^{(2)}_{l;1})\end{pmatrix}.$$

From these equations, we obtain the assertion. □

Remark 6.8. *Since $V_{(l+1,l,l)}$ and $V_{(l,l,l-1)}$ are three dimensional, the differential equations obtained from C_{2i} $(i=1,2,3)$ do not suffice to characterize the Whittaker functions.*

6.5. **Differential equations.** To obtain the explicit actions of the operators C_{2i}, $D^{(\pm,\mp)}_{jk}$ $(1\le i,j,k\le 3)$, we may express these operators in the normal order modulo $[\mathfrak{n},\mathfrak{n}]$ with respect to the Iwasawa decomposition of $\mathfrak{g}$, according to the following lemma.

Lemma 6.9. *(i) Let $\phi_{T,l}$, $\phi^{(1)}_{T,l;i}$, $\phi^{(2)}_{T,l;i}$ $(i=1,2,3)$ be the elements of $C^\infty_\xi(N_{\min}\backslash G)$ defined in Proposition 6.5 and 6.7. The explicit expressions of the action of $\mathfrak{k}_{\mathbf{C}}$ on these functions given as follows.*

- *the case of $(\sigma_1,\sigma_2,\sigma_3)=(0,0,0),(1,1,1)$.*

$$\kappa(E_{ii})\phi_{T,l}=l\phi_{T,l}\qquad(1\le i\le 3),$$
$$\kappa(E_{jk})\phi_{T,l}=0\qquad(1\le j\ne k\le 3).$$

- *the case of $(\sigma_1,\sigma_2,\sigma_3)\ne(0,0,0),(1,1,1)$.*

$$\kappa(E_{jj})\phi^{(1)}_{T,l;i}=(l+\delta_{ij})\phi^{(1)}_{T,l;i}\qquad(1\le i,j\le 3),$$
$$\kappa(E_{mn})\phi^{(1)}_{T,l;k}=\delta_{nk}\phi^{(1)}_{T,l;m}\qquad(1\le k\le 3,\ 1\le m\ne n\le 3),$$

and

$$\kappa(E_{jj})\phi^{(2)}_{T,l;i}=(l-\delta_{4-i\,j})\phi^{(2)}_{T,l;i}\qquad(1\le i,j\le 3),$$
$$\kappa(E_{mn})\phi^{(2)}_{T,l;k}=(-1)^{m+n+1}\delta_{4-m\,k}\phi^{(2)}_{T,l;4-n}\qquad(1\le k\le 3,\ 1\le m\ne n\le 3).$$

(ii) Let $\phi\in C^\infty_\xi(N_{\min}\backslash G)$. For $X\in U(\mathfrak{n}_{\mathbf{C}})$, $Y\in U(\mathfrak{a}_{\mathbf{C}})$ and $a\in A_{\min}$, we have the equation $(\mathrm{Ad}(a^{-1})X)Y\phi(a)=\xi(X)(Y\phi)(a)$. In particular, for $a=\mathrm{diag}(a_1,a_2,a_3,a_1^{-1},a_2^{-1},a_3^{-1})\in A_{\min}$, we have

$$H_i\phi(a)=a_i\frac{\partial}{\partial a_i}\phi(a)\quad(1\le i\le 3),\qquad E_{e_1-e_2}\phi(a)=2\pi\sqrt{-1}c_{12}\frac{a_1}{a_2}\phi(a),$$

$$E_{e_2-e_3}\phi(a)=2\pi\sqrt{-1}c_{23}\frac{a_2}{a_3}\phi(a), \qquad E_{2e_3}\phi(a)=2\pi\sqrt{-1}c_3a_3^2\phi(a),$$

and $E_\alpha\phi(a)=0$ *for* $\alpha\in\Sigma^+\backslash\{e_1-e_2,\ e_2-e_3,\ 2e_3\}$.

Proof. From the definition of $\phi_{T,l},\ \phi^{(1)}_{T,l;i},\ \phi^{(2)}_{T,l;i}$ $(i=1,2,3)$ and Lemma 3.1, we obtain the statement (i). The statement (ii) is obvious from the definition of $C^\infty_\xi(N_{\min}\backslash G)$. □

Moreover, we have the following lemma which is required to get the expressions of the elements in $U(\mathfrak{g}_{\mathbf{C}})$ in normal order. In the following, we denote $X\equiv Y$ for two elements X and Y in $U(\mathfrak{g}_{\mathbf{C}})$ when $X-Y\in[\mathfrak{n},\mathfrak{n}]U(\mathfrak{g}_{\mathbf{C}})$.

Lemma 6.10. *(i) The root vectors* $X_{\pm ij}$ *in* $\mathfrak{p}_\pm$ *have the following expressions:*

$$X_{+ij}\equiv\begin{cases}H_i+\kappa(E_{ii}) & (i=j=1,2),\\ E_{e_i-e_j}+\kappa(E_{ji}) & ((i,j)=(1,2),(2,3)),\end{cases}$$
$$X_{-ij}\equiv\begin{cases}H_i-\kappa(E_{ii}) & (i=j=1,2),\\ E_{e_i-e_j}-\kappa(E_{ij}) & ((i,j)=(1,2),(2,3)),\end{cases}$$

and $X_{\pm33}\equiv\pm2\sqrt{-1}E_{2e_3}+H_3\pm\kappa(E_{33}),\ X_{+13}\equiv\kappa(E_{31}),\ X_{-13}\equiv-\kappa(E_{13})$.
(ii) Each (i,j)*-minor* M_{+ij} *in the matrix* $m_2(C_+)$ *has the following expression.*

$$\begin{aligned}
M_{+11}\equiv&(H_2-1)X_{+33}+X_{+33}\kappa(E_{22})-E_{e_2-e_3}X_{+23}-X_{+23}\kappa(E_{32}),\\
M_{+22}\equiv&(H_1-1)X_{+33}+X_{+33}\kappa(E_{11})-X_{+13}\kappa(E_{31}),\\
M_{+33}\equiv&(H_1-1)X_{+22}+X_{+22}\kappa(E_{11})-E_{e_1-e_2}X_{+12}-X_{+12}\kappa(E_{21}),\\
M_{+12}\equiv&E_{e_1-e_2}X_{+33}+X_{+33}\kappa(E_{21})-X_{+23}\kappa(E_{31}),\\
M_{+23}\equiv&(H_1-1)X_{+23}+X_{+23}\kappa(E_{11})-X_{+12}\kappa(E_{31}),\\
M_{+13}\equiv&E_{e_1-e_2}X_{+23}+X_{+23}\kappa(E_{21})-X_{+22}\kappa(E_{31}).
\end{aligned}$$

(iii) Each (i,j)*-minor* M_{-ij} *in the matrix* $m_2(C_-)$ *has the following expression.*

$$\begin{aligned}
M_{-11}\equiv&(H_2-1)X_{-33}-X_{-33}\kappa(E_{22})-E_{e_2-e_3}X_{-23}+X_{-23}\kappa(E_{23}),\\
M_{-22}\equiv&(H_1-1)X_{-33}-X_{-33}\kappa(E_{11})+X_{-13}\kappa(E_{13}),\\
M_{-33}\equiv&(H_1-1)X_{-22}-X_{-22}\kappa(E_{11})-E_{e_1-e_2}X_{-12}+X_{-12}\kappa(E_{12}),\\
M_{-12}\equiv&E_{e_1-e_2}X_{-33}-X_{-33}\kappa(E_{12})+X_{-23}\kappa(E_{13}),\\
M_{-23}\equiv&(H_1-1)X_{-23}-X_{-23}\kappa(E_{11})+X_{-12}\kappa(E_{13}),\\
M_{-13}\equiv&E_{e_1-e_2}X_{-23}-X_{-23}\kappa(E_{12})+X_{-22}\kappa(E_{13}).
\end{aligned}$$

Proof. The statement (i) is obvious from Lemma 3.4. The statements (ii), (iii) are obtain by direct computation using tables in the proof of Lemma 3.4. □

By using above lemma and tables in the proof of Lemma 3.4, we have the following expressions of the elements $D^{(\pm,\mp)}_{ij}$ $(1\le i,j\le3)$, C_{2k} $(1\le k\le2)$, $m_3(C_\pm)$ in normal order:
For $1\le i\le3$, we have

$$\begin{aligned}
D^{(+,-)}_{1i}\equiv&(H_1-4)X_{-1i}+X_{-1i}\kappa(E_{11})+E_{e_1-e_2}X_{-2i}+X_{-2i}\kappa(E_{21})+X_{-3i}\kappa(E_{31}),\\
D^{(+,-)}_{2i}\equiv&E_{e_1-e_2}X_{-1i}+X_{-1i}\kappa(E_{21})+(H_2-3+\delta_{1i})X_{-2i}+X_{-2i}\kappa(E_{22})\\
&+E_{e_2-e_3}X_{-3i}+X_{-3i}\kappa(E_{32})-\delta_{2i}X_{11},\\
D^{(+,-)}_{3i}\equiv&X_{-1i}\kappa(E_{31})+E_{e_2-e_3}X_{-2i}+X_{-2i}\kappa(E_{32})\\
&+(H_3-1-\delta_{3i}+2\sqrt{-1}E_{2e_3})X_{-3i}+X_{-3i}\kappa(E_{33})-\delta_{3i}(X_{-11}+X_{-22}),
\end{aligned}$$

and

$$D^{(-,+)}_{1i}\equiv(H_1-4)X_{+1i}-X_{+1i}\kappa(E_{11})+E_{e_1-e_2}X_{+2i}-X_{+2i}\kappa(E_{12})-X_{+3i}\kappa(E_{13}),$$

$$\begin{aligned}
D_{2i}^{(-,+)} \equiv& E_{e_1-e_2}X_{+1i} - X_{+1i}\kappa(E_{12}) + (H_2-3+\delta_{1i})X_{+2i} - X_{+2i}\kappa(E_{22})\\
&+ E_{e_2-e_3}X_{+3i} - X_{+3i}\kappa(E_{23}) - \delta_{2i}X_{+11},\\
D_{3i}^{(-,+)} \equiv& - X_{+1i}\kappa(E_{13}) + E_{e_2-e_3}X_{+2i} - X_{+2i}\kappa(E_{23})\\
&+ (H_3-1-\delta_{3i}-2\sqrt{-1}E_{2e_3})X_{+3i} - X_{+3i}\kappa(E_{33}) - \delta_{3i}(X_{+11}+X_{+22}).
\end{aligned}$$

Here we denote $X_{\pm ji} = X_{\pm ij}$ $(1\le i<j\le 3)$.

$$\begin{aligned}
C_2 \equiv\ & (H_1-6)X_{-11} + X_{-11}\kappa(E_{11}) + (H_2-4)X_{-22} + X_{-22}\kappa(E_{22})\\
&+ (H_3+2\sqrt{-1}E_{2e_3}-2)X_{-33} + X_{-33}\kappa(E_{33}) + 2E_{e_1-e_2}X_{-12}\\
&+ 2X_{-12}\kappa(E_{21}) + 2E_{e_2-e_3}X_{-23} + 2X_{-23}\kappa(E_{32}) + 2X_{-13}\kappa(E_{31}).\\
C_4 \equiv\ & (H_2-1)\big\{(2\sqrt{-1}E_{2e_3}+H_3)M_{-11} + M_{-11}(\kappa(E_{33})-2)\big\}\\
&+ \big\{(2\sqrt{-1}E_{2e_3}+H_3)M_{-11} + M_{-11}(\kappa(E_{33})-2)\big\}(\kappa(E_{22})-2)\\
&- E_{e_2-e_3}^2M_{-11} - 2E_{e_2-e_3}M_{-11}\kappa(E_{32}) - M_{-11}\kappa(E_{32})^2\\
&+ (H_1-1)\big\{(2\sqrt{-1}E_{2e_3}+H_3)M_{-22} + M_{-22}(\kappa(E_{33})-2)\big\}\\
&+ \big\{(2\sqrt{-1}E_{2e_3}+H_3)M_{-22} + M_{-22}(\kappa(E_{33})-2)\big\}(\kappa(E_{11})-2) - M_{-22}\kappa(E_{31})^2\\
&+ (H_1-1)\big\{H_2M_{-33} + M_{-33}(\kappa(E_{22})-2)\big\}\\
&+ \big\{H_2M_{-33} + M_{-33}(\kappa(E_{22})-2)\big\}(\kappa(E_{11})-2)\\
&- E_{e_1-e_2}^2M_{-33} - 2E_{e_1-e_2}M_{-33}\kappa(E_{21}) - M_{-33}\kappa(E_{21})^2\\
&+ 2E_{e_1-e_2}\big\{(2\sqrt{-1}E_{2e_3}+H_3)M_{-12} + M_{-12}(\kappa(E_{33})-2)\big\}\\
&+ 2\big\{(2\sqrt{-1}E_{2e_3}+H_3)M_{-12} + M_{-12}(\kappa(E_{33})-2)\big\}\kappa(E_{21})\\
&- 2\big\{(2\sqrt{-1}E_{2e_3}+H_3)M_{-22} + M_{-22}(\kappa(E_{33})-2)\big\} - 2E_{e_2-e_3}M_{-23}\\
&- 2(E_{e_2-e_3}M_{-12} + M_{-12}\kappa(E_{32}) - M_{-13})\kappa(E_{31}) + 2M_{-33} - 2M_{-23}\kappa(E_{32})\\
&+ 2(H_1-1)(E_{e_2-e_3}M_{-23} + M_{-23}\kappa(E_{32}) - M_{-33})\\
&+ 2(E_{e_2-e_3}M_{-23} + M_{-23}\kappa(E_{32}) - M_{-33})(\kappa(E_{11})-2)\\
&- 2(E_{e_1-e_2}M_{-23} + M_{-23}\kappa(E_{21}))\kappa(E_{31})\\
&+ 2E_{e_1-e_2}(E_{e_2-e_3}M_{-13} + M_{-13}\kappa(E_{32})) - 2E_{e_2-e_3}M_{-23}\\
&- 2(H_2-3)M_{-33} - 2M_{-33}\kappa(E_{22}) - 2M_{-23}\kappa(E_{32})\\
&+ 2(E_{e_2-e_3}M_{-13} + M_{-13}\kappa(E_{32}))\kappa(E_{21})\\
&- 2\big\{(H_2-2)M_{-13} + M_{-13}\kappa(E_{22})\big\}\kappa(E_{31}),\\
m_3(C_+) \equiv& (H_1-2)M_{+11} + M_{+11}\kappa(E_{11}) - E_{e_1-e_2}M_{+12} - M_{+12}\kappa(E_{21}) + M_{+13}\kappa(E_{31}),\\
m_3(C_-) \equiv& (H_1-2)M_{-11} - M_{-11}\kappa(E_{11}) - E_{e_1-e_2}M_{-12} + M_{-12}\kappa(E_{12}) - M_{-13}\kappa(E_{13}).
\end{aligned}$$

From above expressions and Lemma 6.9(i), We can summarize the explicit actions of the operators C_{2i} $(1\le i\le 3)$ and $D_{jk}^{(\pm,\mp)}$ $(1\le j,k\le 3)$ on the functions in Proposition 6.5, 6.7.

Proposition 6.11. *The operators C_{2i} $(1\le i\le 3)$ and $D_{jk}^{(\pm,\mp)}$ $(1\le j,k\le 3)$ acting on the functions $\phi_{T,l}$, $\phi_{T,l;i}^{(1)}$, $\phi_{T,l;i}^{(2)}$ $(i=1,2,3)$ in Proposition 6.5, 6.7 are given as follows.*

- *the case of $(\sigma_1,\sigma_2,\sigma_3)=(0,0,0),(1,1,1)$.*

$$\begin{aligned}
C_2\phi_{T,l} =& \{(H_1+l-6)(H_1-l) + (H_2+l-4)(H_2-l)\\
&+ (H_3+l-2+2\sqrt{-1}E_{2e_3})(H_3-l-2\sqrt{-1}E_{2e_3}) + 2E_{e_1-e_2}^2 + 2E_{e_2-e_3}^2\}\phi_{T,l},
\end{aligned}$$

$$\begin{aligned}
C_4\phi_{T,l} =&\Big\{\{(H_2+l-3)(H_3+l-2+2\sqrt{-1}E_{2e_3})-E^2_{e_2-e_3}\}\\
&\quad\times\{(H_2-l-1)(H_3-l-2\sqrt{-1}E_{2e_3})-E^2_{e_2-e_3}\}\\
&+(H_1+l-5)(H_3+l-2+2\sqrt{-1}E_{2e_3})(H_1-l-1)(H_3-l-2\sqrt{-1}E_{2e_3})\\
&+\{(H_1+l-5)(H_2+l-4)-E^2_{e_1-e_2}\}\{(H_1-l-1)(H_2-l)-E^2_{e_1-e_2}\}\\
&+2E^2_{e_1-e_2}(H_3+l-2+2\sqrt{-1}E_{2e_3})(H_3-l-2\sqrt{-1}E_{2e_3})\\
&+2E^2_{e_1-e_2}E^2_{e_2-e_3}+2E^2_{e_2-e_3}(H_1+l-5)(H_1-l-1)\Big\}\phi_{T,l},\\
C_6\phi_{T,l} =&\{(H_1+l-4)(H_2+l-3)(H_3+l-2+2\sqrt{-1}E_{2e_3})\\
&-E^2_{e_2-e_3}(H_1+l-4)-E^2_{e_1-e_2}(H_3+l-2+2\sqrt{-1}E_{2e_3})\}\\
&\times\{(H_1-l-2)(H_2-l-1)(H_3-l-2\sqrt{-1}E_{2e_3})\\
&-E^2_{e_2-e_3}(H_1-l-2)-E^2_{e_1-e_2}(H_3-l-2\sqrt{-1}E_{2e_3})\}\phi_{T,l}.
\end{aligned}$$

- *the case of* $(\sigma_1,\sigma_2,\sigma_3)\neq(0,0,0),(1,1,1)$.

$$\begin{aligned}
&D^{(+,-)}_{11}\phi^{(1)}_{T,l;1}+D^{(+,-)}_{12}\phi^{(1)}_{T,l;2}+D^{(+,-)}_{13}\phi^{(1)}_{T,l;3}\\
&=\{(H_1+l-3)(H_1-l-3)+E^2_{e_1-e_2}\}\phi^{(1)}_{T,l;1}+E_{e_1-e_2}(H_1+H_2-4)\phi^{(1)}_{T,l;2}\\
&\quad+E_{e_1-e_2}E_{e_2-e_3}\phi^{(1)}_{T,l;3},\\
&D^{(+,-)}_{21}\phi^{(1)}_{T,l;1}+D^{(+,-)}_{22}\phi^{(1)}_{T,l;2}+D^{(+,-)}_{23}\phi^{(1)}_{T,l;3}\\
&=E_{e_1-e_2}(H_1+H_2-6)\phi^{(1)}_{T,l;1}+\{(H_2+l-2)(H_2-l-2)+E^2_{e_1-e_2}+E^2_{e_2-e_3}\}\phi^{(1)}_{T,l;2}\\
&\quad+E_{e_2-e_3}(H_2+H_3-2-2\sqrt{-1}E_{2e_3})\phi^{(1)}_{T,l;3},\\
&D^{(+,-)}_{31}\phi^{(1)}_{T,l;1}+D^{(+,-)}_{32}\phi^{(1)}_{T,l;2}+D^{(+,-)}_{33}\phi^{(1)}_{T,l;3}\\
&=+E_{e_1-e_2}E_{e_2-e_3}\phi^{(1)}_{T,l;1}+E_{e_2-e_3}(H_2+H_3-4+2\sqrt{-1}E_{2e_3})\phi^{(1)}_{T,l;2}\\
&\quad+\{(H_3+l-1+2\sqrt{-1}E_{2e_3})(H_3-l-1-2\sqrt{-1}E_{2e_3})+E^2_{e_2-e_3}\}\phi^{(1)}_{T,l;3},
\end{aligned}$$

$$\begin{aligned}
C_2\phi^{(1)}_{T,l;i} =&\{(H_1+l-6+\delta_{1i})(H_1-l-\delta_{1i})+(H_2+l-4+\delta_{2i})(H_2-l-\delta_{2i})\\
&+(H_3+l-2+\delta_{3i}+2\sqrt{-1}E_{2e_3})(H_3-l-\delta_{3i}-2\sqrt{-1}E_{2e_3})\\
&+2E^2_{e_1-e_2}+2E^2_{e_2-e_3}\}\phi^{(1)}_{T,l;i}\\
&-2\delta_{1i}\{2\phi^{(1)}_{T,l;1}-E_{e_1-e_2}\phi^{(1)}_{T,l;2}\}-2\delta_{2i}\{E_{e_1-e_2}\phi^{(1)}_{T,l;1}+\phi^{(1)}_{T,l;2}-E_{e_2-e_3}\phi^{(1)}_{T,l;3}\}\\
&-2\delta_{3i}E_{e_2-e_3}\phi^{(1)}_{T,l;2},\\
C_4\phi^{(1)}_{T,l;i} =&\Big\{\{(H_2+l-3+\delta_{2i})(H_3+l-2+\delta_{3i}+2\sqrt{-1}E_{2e_3})-E^2_{e_2-e_3}\}\\
&\quad\times\{(H_2-l-1-\delta_{2i})(H_3-l-\delta_{3i}-2\sqrt{-1}E_{2e_3})-E^2_{e_2-e_3}\}\\
&+(H_1+l-5+\delta_{1i})(H_3+l-2+\delta_{3i}+2\sqrt{-1}E_{2e_3})\\
&\quad\times(H_1-l-1-\delta_{1i})(H_3-l-\delta_{3i}-2\sqrt{-1}E_{2e_3})\\
&+\{(H_1+l-5+\delta_{1i})(H_2+l-4+\delta_{2i})-E^2_{e_1-e_2}\}\\
&\quad\times\{(H_1-l-1-\delta_{1i})(H_2-l-\delta_{2i})-E^2_{e_1-e_2}\}\\
&+2E^2_{e_1-e_2}(H_3+l-2+\delta_{3i}+2\sqrt{-1}E_{2e_3})(H_3-l-\delta_{3i}-2\sqrt{-1}E_{2e_3})\\
&+2E^2_{e_1-e_2}E^2_{e_2-e_3}+2E^2_{e_2-e_3}(H_1+l-5+\delta_{1i})(H_1-l-1-\delta_{1i})\Big\}\phi^{(1)}_{T,l;i}
\end{aligned}$$

$$
\begin{aligned}
&+ 2\delta_{1i}\Big\{ - \{(H_2 + l - 4)(H_2 - l) \\
&\quad + (H_3 + l - 2 + 2\sqrt{-1}E_{2e_3})(H_3 - l - 2\sqrt{-1}E_{2e_3}) + 3E^2_{e_1-e_2} + 2E^2_{e_2-e_3}\}\phi^{(1)}_{T,l;1} \\
&\quad + E_{e_1-e_2}\{(H_3 + l - 2 + 2\sqrt{-1}E_{2e_3})(H_3 - l - 2\sqrt{-1}E_{2e_3}) \\
&\quad - (H_1 - l - 1)(H_2 - l - 1) + H_1 + H_2 - 6 + E^2_{e_1-e_2} + E^2_{e_2-e_3}\}\phi^{(1)}_{T,l;2} \\
&\quad - E_{e_1-e_2}E_{e_2-e_3}(H_1 + H_2 + H_3 - l - 6 - 2\sqrt{-1}E_{2e_3})\phi^{(1)}_{T,l;3}\Big\} \\
&+ 2\delta_{2i}\Big\{ - E_{e_1-e_2}\{(H_3 + l - 2 + 2\sqrt{-1}E_{2e_3})(H_3 - l - 2\sqrt{-1}E_{2e_3}) \\
&\quad - (H_1 + l - 4)(H_2 + l - 4) - (H_1 + H_2 - 4) + E^2_{e_1-e_2} + E^2_{e_2-e_3}\}\phi^{(1)}_{T,l;1} \\
&\quad - \{(H_1 + l - 5)(H_1 - l - 1) + E^2_{e_1-e_2} + 2E^2_{e_2-e_3}\}\phi^{(1)}_{T,l;2} \\
&\quad + E_{e_2-e_3}\{(H_1 + l - 5)(H_1 - l - 1) \\
&\quad - (H_2 - l - 1)(H_3 - l - 1 - 2\sqrt{-1}E_{2e_3}) + E^2_{e_1-e_2} + E^2_{e_2-e_3}\}\phi^{(1)}_{T,l;3}\Big\} \\
&+ 2\delta_{3i}\Big\{E_{e_1-e_2}E_{e_2-e_3}(H_1 + H_2 + H_3 + l - 7 + 2\sqrt{-1}E_{2e_3})\phi^{(1)}_{T,l;1} \\
&\quad + E_{e_2-e_3}\{(H_2 + l - 2)(H_3 + l - 2 + 2\sqrt{-1}E_{2e_3}) \\
&\quad - (H_1 + l - 5)(H_1 - l - 1) - E^2_{e_1-e_2} - E^2_{e_2-e_3}\}\phi^{(1)}_{T,l;2}\Big\},
\end{aligned}
$$

$$
\begin{aligned}
C_6\phi^{(1)}_{T,l;i} =& \{(H_1 + l - 4 + \delta_{1i})(H_2 + l - 3 + \delta_{2i})(H_3 + l - 2 + \delta_{3i} + 2\sqrt{-1}E_{2e_3}) \\
&\quad - E^2_{e_2-e_3}(H_1 + l - 4 + \delta_{1i}) - E^2_{e_1-e_2}(H_3 + l - 2 + \delta_{3i} + 2\sqrt{-1}E_{2e_3})\} \\
&\quad \times \{(H_1 - l - 2 - \delta_{1i})(H_2 - l - 1 - \delta_{2i})(H_3 - l - \delta_{3i} - 2\sqrt{-1}E_{2e_3}) \\
&- E^2_{e_2-e_3}(H_1 - l - 2 - \delta_{1i}) - E^2_{e_1-e_2}(H_3 - l - \delta_{3i} - 2\sqrt{-1}E_{2e_3})\}\phi^{(1)}_{T,l;i} \\
&+ 2\delta_{1i}\Big\{ - 2E^2_{e_1-e_2}\{(H_3 + l - 2 + 2\sqrt{-1}E_{2e_3})(H_3 - l - 2\sqrt{-1}E_{2e_3}) + E^2_{e_2-e_3}\}\phi^{(1)}_{T,l;1} \\
&\quad - E_{e_1-e_2}\{(H_3 + l - 2 + 2\sqrt{-1}E_{2e_3})(H_1 - l - 2)(H_2 - l - 2)(H_3 - l - 2\sqrt{-1}E_{2e_3}) \\
&\quad - E^2_{e_2-e_3}(H_3 + l - 2 + 2\sqrt{-1}E_{2e_3})(H_1 - l - 2) \\
&\quad - E^2_{e_1-e_2}(H_3 + l - 2 + 2\sqrt{-1}E_{2e_3})(H_3 - l - 2\sqrt{-1}E_{2e_3})\}\phi^{(1)}_{T,l;2} \\
&\quad + E_{e_1-e_2}E_{e_2-e_3}\{(H_1 - l - 2)(H_2 - l - 1)(H_3 - l - 1 - 2\sqrt{-1}E_{2e_3}) \\
&\quad - E^2_{e_2-e_3}(H_1 - l - 2) - E^2_{e_1-e_2}(H_3 - l - 1 - 2\sqrt{-1}E_{2e_3})\}\phi^{(1)}_{T,l;3}\Big\} \\
&+ 2\delta_{2i}\Big\{E_{e_1-e_2}\{(H_1 + l - 3)(H_2 + l - 3) \\
&\quad \times (H_3 + l - 2 + 2\sqrt{-1}E_{2e_3})(H_3 - l - 2\sqrt{-1}E_{2e_3}) \\
&\quad - E^2_{e_2-e_3}(H_1 + l - 3)(H_3 - l - 2 - 2\sqrt{-1}E_{2e_3}) \\
&\quad - E^2_{e_1-e_2}(H_3 + l - 2 + 2\sqrt{-1}E_{2e_3})(H_3 - l - 2\sqrt{-1}E_{2e_3})\}\phi^{(1)}_{T,l;1} \\
&\quad - 2E^2_{e_2-e_3}(H_1 + l - 4)(H_1 - l - 2)\phi^{(1)}_{T,l;2} \\
&\quad - E_{e_2-e_3}(H_1 + l - 4)\{(H_1 - l - 2)(H_2 - l - 1)(H_3 - l - 1 - 2\sqrt{-1}E_{2e_3}) \\
&\quad - E^2_{e_2-e_3}(H_1 - l - 2) - E^2_{e_1-e_2}(H_3 - l - 1 - 2\sqrt{-1}E_{2e_3})\}\phi^{(1)}_{T,l;3}\Big\} \\
&+ 2\delta_{3i}\Big\{ - E_{e_1-e_2}E_{e_2-e_3}\{(H_1 + l - 3)(H_2 + l - 3)(H_3 + l - 2 + 2\sqrt{-1}E_{2e_3})
\end{aligned}
$$

$$
\begin{aligned}
&- E_{e_2-e_3}^2(H_1+l-3) - E_{e_1-e_2}^2(H_3+l-2+2\sqrt{-1}E_{2e_3})\}\phi_{T,l;1}^{(1)} \\
&+ E_{e_2-e_3}\{(H_1+l-4)(H_2+l-2)(H_3+l-2+2\sqrt{-1}E_{2e_3}) \\
&- E_{e_2-e_3}^2(H_1+l-4) - E_{e_1-e_2}^2(H_3+l-2+2\sqrt{-1}E_{2e_3})\}(H_1-l-2)\phi_{T,l;2}^{(1)}\Big\},
\end{aligned}
$$

and

$$
\begin{aligned}
&- D_{11}^{(-,+)}\phi_{T,l;3}^{(2)} + D_{12}^{(-,+)}\phi_{T,l;2}^{(2)} - D_{13}^{(-,+)}\phi_{T,l;1}^{(2)} \\
&= - E_{e_1-e_2}E_{e_2-e_3}\phi_{T,l;1}^{(2)} + E_{e_1-e_2}(H_1+H_2-4)\phi_{T,l;2}^{(2)} \\
&\quad - \{(H_1-l-3)(H_1+l-3) + E_{e_1-e_2}^2\}\phi_{T,l;3}^{(2)}, \\
&- D_{21}^{(-,+)}\phi_{T,l;3}^{(2)} + D_{22}^{(-,+)}\phi_{T,l;2}^{(2)} - D_{23}^{(-,+)}\phi_{T,l;1}^{(2)} \\
&= - E_{e_2-e_3}(H_2+H_3-2+2\sqrt{-1}E_{2e_3})\phi_{T,l;1}^{(2)} \\
&\quad + \{(H_2-l-2)(H_2+l-2) + E_{e_1-e_2}^2 + E_{e_2-e_3}^2\}\phi_{T,l;2}^{(2)} - E_{e_1-e_2}(H_1+H_2-6)\phi_{T,l;3}^{(2)}, \\
&- D_{31}^{(-,+)}\phi_{T,l;3}^{(2)} + D_{32}^{(-,+)}\phi_{T,l;2}^{(2)} - D_{33}^{(-,+)}\phi_{T,l;1}^{(2)} \\
&= - \{(H_3-l-1-2\sqrt{-1}E_{2e_3})(H_3+l-1+2\sqrt{-1}E_{2e_3}) + E_{e_2-e_3}^2\}\phi_{T,l;1}^{(2)} \\
&\quad + E_{e_2-e_3}(H_2+H_3-4-2\sqrt{-1}E_{2e_3})\phi_{T,l;2}^{(2)} - E_{e_1-e_2}E_{e_2-e_3}\phi_{T,l;3}^{(2)},
\end{aligned}
$$

$$
\begin{aligned}
C_2\phi_{T,l;i}^{(2)} =& \{(H_1+l-6-\delta_{3i})(H_1-l+\delta_{3i}) + (H_2+l-4-\delta_{2i})(H_2-l+\delta_{2i}) \\
&\quad + (H_3+l-2-\delta_{1i}+2\sqrt{-1}E_{2e_3})(H_3-l+\delta_{1i}-2\sqrt{-1}E_{2e_3}) \\
&\quad + 2E_{e_1-e_2}^2 + 2E_{e_2-e_3}^2\}\phi_{T,l;i}^{(2)} \\
&- 2\delta_{1i}\{2\delta_{1i}\phi_{T,l;1}^{(2)} - E_{e_2-e_3}\delta_{1i}\phi_{T,l;2}^{(2)}\} - 2\delta_{2i}\{E_{e_2-e_3}\phi_{T,l;1}^{(2)} + \phi_{T,l;2}^{(2)} - E_{e_1-e_2}\phi_{T,l;3}^{(2)}\} \\
&- 2\delta_{3i}E_{e_1-e_2}\phi_{T,l;2}^{(2)}, \\
C_4\phi_{T,l;i}^{(2)} =& \Big\{\{(H_2+l-3-\delta_{2i})(H_3+l-2-\delta_{1i}+2\sqrt{-1}E_{2e_3}) - E_{e_2-e_3}^2\} \\
&\quad \times \{(H_2-l-1+\delta_{2i})(H_3-l+\delta_{1i}-2\sqrt{-1}E_{2e_3}) - E_{e_2-e_3}^2\} \\
&\quad + (H_1+l-5-\delta_{3i})(H_3+l-2-\delta_{1i}+2\sqrt{-1}E_{2e_3}) \\
&\quad \times (H_1-l-1+\delta_{3i})(H_3-l+\delta_{1i}-2\sqrt{-1}E_{2e_3}) \\
&\quad + \{(H_1+l-5-\delta_{3i})(H_2+l-4-\delta_{2i}) - E_{e_1-e_2}^2\} \\
&\quad \times \{(H_1-l-1+\delta_{3i})(H_2-l+\delta_{2i}) - E_{e_1-e_2}^2\} \\
&\quad + 2E_{e_1-e_2}^2(H_3+l-2-\delta_{1i}+2\sqrt{-1}E_{2e_3})(H_3-l+\delta_{1i}-2\sqrt{-1}E_{2e_3}) \\
&\quad + 2E_{e_2-e_3}^2\{(H_1+l-5-\delta_{3i})(H_1-l-1+\delta_{3i}) + E_{e_1-e_2}^2\}\Big\}\phi_{T,l;i}^{(2)} \\
&+ 2\delta_{1i}\Big\{ - \{(H_1+l-5)(H_1-l-1) + (H_2+l-3)(H_2-l-1) \\
&\quad + 2E_{e_1-e_2}^2 + 3E_{e_2-e_3}^2\}\phi_{T,l;1}^{(2)} \\
&\quad + E_{e_2-e_3}\{(H_1+l-5)(H_1-l-1) \\
&\quad - (H_2-l)(H_3-l-2\sqrt{-1}E_{2e_3}) \\
&\quad - (H_2+H_3-2-2\sqrt{-1}E_{2e_3}) + E_{e_1-e_2}^2 + E_{e_2-e_3}^2\}\phi_{T,l;2}^{(2)} \\
&\quad + E_{e_1-e_2}E_{e_2-e_3}(H_1+H_2+H_3-l-3-2\sqrt{-1}E_{2e_3})\phi_{T,l;3}^{(2)}\Big\}
\end{aligned}
$$

$$\begin{aligned}
&+2\delta_{2i}\Big\{-E_{e_2-e_3}\{(H_1+l-5)(H_1-l-1)\\
&\quad-(H_2+l-3)(H_3+l-3+2\sqrt{-1}E_{2e_3})\\
&\quad+H_2+H_3-4+2\sqrt{-1}E_{2e_3}+E^2_{e_1-e_2}+E^2_{e_2-e_3}\}\phi^{(2)}_{T,l;1}\\
&\quad-\{(H_3+l-2+2\sqrt{-1}E_{2e_3})(H_3-l-2\sqrt{-1}E_{2e_3})\\
&\quad+2E^2_{e_1-e_2}+E^2_{e_2-e_3}\}\phi^{(2)}_{T,l;2}\\
&\quad+E_{e_1-e_2}\{-(H_1-l)(H_2-l)\\
&\quad+(H_3+l-2+2\sqrt{-1}E_{2e_3})(H_3-l-2\sqrt{-1}E_{2e_3})\\
&\quad+E^2_{e_1-e_2}+E^2_{e_2-e_3}\}\phi^{(2)}_{T,l;3}\Big\}\\
&+2\delta_{3i}\Big\{-E_{e_1-e_2}E_{e_2-e_3}(H_1+H_2+H_3+l-9+2\sqrt{-1}E_{2e_3})\phi^{(2)}_{T,l;1}\\
&\quad+E_{e_1-e_2}\{(H_1+l-5)(H_2+l-5)\\
&\quad-(H_3+l-2+2\sqrt{-1}E_{2e_3})(H_3-l-2\sqrt{-1}E_{2e_3})\\
&\quad-E^2_{e_1-e_2}-E^2_{e_2-e_3}\}\phi^{(2)}_{T,l;2}\Big\},
\end{aligned}$$

$$\begin{aligned}
C_6\phi^{(2)}_{T,l;i}=&\{(H_1+l-4-\delta_{3i})(H_2+l-3-\delta_{2i})(H_3+l-2-\delta_{1i}+2\sqrt{-1}E_{2e_3})\\
&\quad-E^2_{e_2-e_3}(H_1+l-4-\delta_{3i})-E^2_{e_1-e_2}(H_3+l-2-\delta_{1i}+2\sqrt{-1}E_{2e_3})\}\\
&\quad\times\{(H_1-l-2+\delta_{3i})(H_2-l-1+\delta_{2i})(H_3-l+\delta_{1i}-2\sqrt{-1}E_{2e_3})\\
&\quad-E^2_{e_1-e_2}(H_3-l+\delta_{1i}-2\sqrt{-1}E_{2e_3})-E^2_{e_2-e_3}(H_1-l-2+\delta_{3i})\}\phi^{(2)}_{T,l;i}\\
&+2\delta_{1i}\Big\{-2E^2_{e_2-e_3}\{(H_1+l-4)(H_1-l-2)+E^2_{e_1-e_2}\}\phi^{(2)}_{T,l;1}\\
&\quad-E_{e_2-e_3}\{(H_1+l-4)(H_1-l-2)\\
&\quad\times(H_2-l)(H_3-l-2\sqrt{-1}E_{2e_3})\\
&\quad-E^2_{e_1-e_2}(H_1+l-4)(H_3-l-2\sqrt{-1}E_{2e_3})\\
&\quad-E^2_{e_2-e_3}(H_1+l-4)(H_1-l-2)\}\phi^{(2)}_{T,l;2}\\
&\quad-E_{e_1-e_2}E_{e_2-e_3}\{(H_1-l-1)(H_2-l-1)(H_3-l-2\sqrt{-1}E_{2e_3})\\
&\quad-E^2_{e_1-e_2}(H_3-l-2\sqrt{-1}E_{2e_3})-E^2_{e_2-e_3}(H_1-l-1)\}\phi^{(2)}_{T,l;3}\Big\}\\
&+2\delta_{2i}\Big\{E_{e_2-e_3}\{(H_1+l-4)(H_2+l-3)\\
&\quad\times(H_3+l-3+2\sqrt{-1}E_{2e_3})(H_1-l-2)\\
&\quad-E^2_{e_2-e_3}(H_1+l-4)(H_1-l-2)\\
&\quad-E^2_{e_1-e_2}(H_3+l-3+2\sqrt{-1}E_{2e_3})(H_1-l)\}\phi^{(2)}_{T,l;1}\\
&\quad-2E^2_{e_1-e_2}(H_3+l-2+2\sqrt{-1}E_{2e_3})(H_3-l-2\sqrt{-1}E_{2e_3})\phi^{(2)}_{T,l;2}\\
&\quad-E_{e_1-e_2}(H_3+l-2+2\sqrt{-1}E_{2e_3})\\
&\quad\times\{(H_1-l-1)(H_2-l-1)(H_3-l-2\sqrt{-1}E_{2e_3})\\
&\quad-E^2_{e_1-e_2}(H_3-l-2\sqrt{-1}E_{2e_3})-E^2_{e_2-e_3}(H_1-l-1)\}\phi^{(2)}_{T,l;3}\Big\}\\
&+2\delta_{3i}\Big\{E_{e_1-e_2}E_{e_2-e_3}\{(H_1+l-4)(H_2+l-3)(H_3+l-3+2\sqrt{-1}E_{2e_3})
\end{aligned}$$

$$
\begin{aligned}
&- E_{e_2-e_3}^2(H_1+l-4) - E_{e_1-e_2}^2(H_3+l-3+2\sqrt{-1}E_{2e_3})\}\phi^{(2)}_{T,l;1}\\
&+ E_{e_1-e_2}\{(H_1+l-4)(H_2+l-4)(H_3+l-2+2\sqrt{-1}E_{2e_3})\\
&- E_{e_2-e_3}^2(H_1+l-4) - E_{e_1-e_2}^2(H_3+l-2+2\sqrt{-1}E_{2e_3})\}\\
&\times (H_3-l-2\sqrt{-1}E_{2e_3})\phi^{(2)}_{T,l;2}\Big\}.
\end{aligned}
$$

Proof. The actions of the operators C_{2i} $(1 \le i \le 2)$, $D^{(\pm,\mp)}_{jk}$ $(1 \le j,k \le 3)$ and $m_3(C_\pm)$ are obtained by direct computation from the expressions in normal order and Lemma 6.9(i).

We can summarize the explicit actions of the operators C_{2i} $(1 \le i \le 3)$ and $D^{(\pm,\mp)}_{jk}$ $(1 \le j,k \le 3)$ on the functions in Proposition 6.5, 6.7. The action of $C_6 = m_3(C_+)m_3(C_-)$ is obtained by the composite of following actions of $m_3(C_+)$ and $m_3(C_-)$:

- the case of $(\sigma_1,\sigma_2,\sigma_3) = (0,0,0),(1,1,1)$.

$$
\begin{aligned}
m_3(C_+)\phi_{T,l-2} =&\{(H_1+l-4)(H_2+l-3)(H_3+l-2+2\sqrt{-1}E_{2e_3})\\
&- E_{e_2-e_3}^2(H_1+l-4) - E_{e_1-e_2}^2(H_3+l-2+2\sqrt{-1}E_{2e_3})\}\phi_{T,l-2},\\
m_3(C_-)\phi_{T,l} =&\{(H_1-l-2)(H_2-l-1)(H_3-l-2\sqrt{-1}E_{2e_3})\\
&- E_{e_2-e_3}^2(H_1-l-2) - E_{e_1-e_2}^2(H_3-l-2\sqrt{-1}E_{2e_3})\}\phi_{T,l}.
\end{aligned}
$$

- the case of $(\sigma_1,\sigma_2,\sigma_3) \neq (0,0,0),(1,1,1)$.

$$
\begin{aligned}
m_3(C_+)\phi^{(1)}_{T,l-2;i} =&\{(H_1+l-4+\delta_{1i})(H_2+l-3+\delta_{2i})(H_3+l-2+\delta_{3i}+2\sqrt{-1}E_{2e_3})\\
&- E_{e_2-e_3}^2(H_1+l-4+\delta_{1i}) - E_{e_1-e_2}^2(H_3+l-2+\delta_{3i}\\
&+2\sqrt{-1}E_{2e_3})\}\phi^{(1)}_{T,l-2;i}\\
&-2\delta_{1i}\{E_{e_1-e_2}(H_3+l-2+2\sqrt{-1}E_{2e_3})\phi^{(1)}_{T,l-2;2} - E_{e_1-e_2}E_{e_2-e_3}\phi^{(1)}_{T,l-2;3}\}\\
&-2\delta_{2i}E_{e_2-e_3}(H_1+l-4)\phi^{(1)}_{T,l-2;3},
\end{aligned}
$$

$$
\begin{aligned}
m_3(C_-)\phi^{(1)}_{T,l;i} =&\{(H_1-l-2-\delta_{1i})(H_2-l-1-\delta_{2i})(H_3-l-\delta_{3i}-2\sqrt{-1}E_{2e_3})\\
&- E_{e_2-e_3}^2(H_1-l-2-\delta_{1i}) - E_{e_1-e_2}^2(H_3-l-\delta_{3i}-2\sqrt{-1}E_{2e_3})\}\phi^{(1)}_{T,l;i}\\
&+2E_{e_1-e_2}(H_3-l-2\sqrt{-1}E_{2e_3})\delta_{2i}\phi^{(1)}_{T,l;1}\\
&-2\delta_{3i}\{E_{e_1-e_2}E_{e_2-e_3}\phi^{(1)}_{T,l;1} - E_{e_2-e_3}(H_1-l-2)\phi^{(1)}_{T,l;2}\},
\end{aligned}
$$

and

$$
\begin{aligned}
m_3(C_+)\phi^{(2)}_{T,l-2;i} =&\{(H_1+l-4-\delta_{3i})(H_2+l-3-\delta_{2i})(H_3+l-2-\delta_{1i}+2\sqrt{-1}E_{2e_3})\\
&- E_{e_2-e_3}^2(H_1+l-4-\delta_{3i})\\
&- E_{e_1-e_2}^2(H_3+l-2-\delta_{1i}+2\sqrt{-1}E_{2e_3})\}\phi^{(2)}_{T,l-2;i}\\
&-2\delta_{1i}\{E_{e_2-e_3}(H_1+l-4)\phi^{(2)}_{T,l-2;2} + E_{e_1-e_2}E_{e_2-e_3}\phi^{(2)}_{T,l-2;3}\}\\
&-2\delta_{2i}E_{e_1-e_2}(H_3+l-2+2\sqrt{-1}E_{2e_3})\phi^{(2)}_{T,l-2;3},\\
m_3(C_-)\phi^{(2)}_{T,l;i} =&\{(H_1-l-2+\delta_{3i})(H_2-l-1+\delta_{2i})(H_3-l+1-1+\delta_{1i}-2\sqrt{-1}E_{2e_3})\\
&- E_{e_1-e_2}^2(H_3-l+\delta_{1i}-2\sqrt{-1}E_{2e_3}) - E_{e_2-e_3}^2(H_1-l-2+\delta_{3i})\}\phi^{(2)}_{T,l;i}\\
&+2\delta_{2i}E_{e_2-e_3}(H_1-l-2)\phi^{(2)}_{T,l;1}\\
&+2\delta_{3i}\{E_{e_1-e_2}E_{e_2-e_3}\phi^{(2)}_{T,l;1} + E_{e_1-e_2}(H_3-l-2\sqrt{-1}E_{2e_3})\phi^{(2)}_{T,l;2}\}.
\end{aligned}
$$

$\square$

To state an explicit form of a holonomic system of partial differential equations satisfied by the A-radial part of each element in $\mathrm{Wh}(\pi_{(\sigma,\nu)}, \xi, \tau)$ for peripheral K-type τ^*, we introduce the coordinates $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$ on A defined by

$$\begin{aligned} &x_1 = \left(\pi c_{12}\frac{a_1}{a_2}\right)^2, \quad x_2 = \left(\pi c_{23}\frac{a_2}{a_3}\right)^2, \quad x_3 = 4\pi c_3 a_3^2, \\ &y_1 = 2\pi c_{12}\frac{a_1}{a_2}, \quad y_2 = 2\pi c_{23}\frac{a_2}{a_3}, \quad y_3 = 4\pi c_3 a_3^2, \end{aligned}$$

for $\mathrm{diag}(a_1, a_2, a_3, a_1^{-1}, a_2^{-1}, a_3^{-1}) \in A$. We use the coordinate x for the case of $(\sigma_1, \sigma_2, \sigma_3) = (0, 0, 0), (1, 1, 1)$, and use the coordinate y for the case of $(\sigma_1, \sigma_2, \sigma_3) \neq (0, 0, 0), (1, 1, 1)$. Then we have following theorem.

Theorem 6.12. *Let T be an element of the space $\mathcal{I}_{\xi, \pi_{(\sigma,\nu)}}$.*
(i) If σ is a character of $M_{\min}$ such that $(\sigma_1, \sigma_2, \sigma_3) = (0, 0, 0)$ or $(1, 1, 1)$, then there exist multiplicity one K-types $\tau_{(l,l,l)}$ ($l \equiv \varepsilon_\sigma \mod 2$). For $l \equiv \varepsilon_\sigma \mod 2$, the Whittaker function $\Phi(T, S_{(l,l,l)}(M_l)) = \phi_{T,l} \otimes f(M_l)^*$ satisfies the following holonomic system of partial differential equations of rank 48:

$$\begin{aligned} &\{(2\partial_{x_1} + l - 6)(2\partial_{x_1} - l) + (-2\partial_{x_1} + 2\partial_{x_2} + l - 4)(-2\partial_{x_1} + 2\partial_{x_2} - l) \\ &+ (-2\partial_{x_2} + 2\partial_{x_3} + l - 2 - x_3)(-2\partial_{x_2} + 2\partial_{x_3} - l + x_3) \\ &- 8x_1 - 8x_2 - \chi_{2,\sigma,\nu,(l,l,l)}\}\phi_{T,l} = 0, \end{aligned}$$

$$\begin{aligned} &\Big\{\{(-2\partial_{x_1} + 2\partial_{x_2} + l - 3)(-2\partial_{x_2} + 2\partial_{x_3} + l - 2 - x_3) + 4x_2\} \\ &\qquad \times \{(-2\partial_{x_1} + 2\partial_{x_2} - l - 1)(-2\partial_{x_2} + 2\partial_{x_3} - l + x_3) + 4x_2\} \\ &\quad + (2\partial_{x_1} + l - 5)(-2\partial_{x_2} + 2\partial_{x_3} + l - 2 - x_3) \\ &\qquad \times (2\partial_{x_1} - l - 1)(-2\partial_{x_2} + 2\partial_{x_3} - l + x_3) \\ &\quad + \{(2\partial_{x_1} + l - 5)(-2\partial_{x_1} + 2\partial_{x_2} + l - 4) + 4x_1\} \\ &\qquad \times \{(2\partial_{x_1} - l - 1)(-2\partial_{x_1} + 2\partial_{x_2} - l) + 4x_1\} \\ &\quad - 8x_1(-2\partial_{x_2} + 2\partial_{x_3} + l - 2 - x_3)(-2\partial_{x_2} + 2\partial_{x_3} - l + x_3) \\ &\quad + 32x_1x_2 - 8x_2(2\partial_{x_1} + l - 5)(2\partial_{x_1} - l - 1) - \chi_{4,\sigma,\nu,(l,l,l)}\Big\}\phi_{T,l} = 0, \end{aligned}$$

$$\begin{aligned} &\Big\{\{(2\partial_{x_1} + l - 4)(-2\partial_{x_1} + 2\partial_{x_2} + l - 3)(-2\partial_{x_2} + 2\partial_{x_3} + l - 2 - x_3) \\ &\quad + 4x_2(2\partial_{x_1} + l - 4) + 4x_1(-2\partial_{x_2} + 2\partial_{x_3} + l - 2 - x_3)\} \\ &\times \{(2\partial_{x_1} - l - 2)(-2\partial_{x_1} + 2\partial_{x_2} - l - 1)(-2\partial_{x_2} + 2\partial_{x_3} - l + x_3) \\ &\quad + 4x_2(2\partial_{x_1} - l - 2) + 4x_1(-2\partial_{x_2} + 2\partial_{x_3} - l + x_3)\} - \chi_{6,\sigma,\nu,(l,l,l)}\Big\}\phi_{T,l} = 0. \end{aligned}$$

(ii) If σ is a character of $M_{\min}$ such that $(\sigma_1, \sigma_2, \sigma_3) \neq (0, 0, 0)$ or $(1, 1, 1)$, there exists multiplicity one K-types $\tau_{(l+1,l,l)}$, $\tau_{(l,l,l-1)}$ ($l \equiv \varepsilon_\sigma \mod 2$).

For $l \equiv \varepsilon_\sigma \mod 2$, Whittaker function $\Phi(T, S_{(l+1,l,l)}(M_l^{(\sigma;1)})) = \sum_{j=1}^{3} \phi_{T,l;j}^{(1)} \otimes f(M_{l;j}^{(1)})^*$ satisfy the following holonomic system of partial differential equations of rank 48:

$$\begin{aligned} &\{(\partial_{y_1} + l - 3)(\partial_{y_1} - l - 3) - y_1^2 - \tilde{\chi}_{2,\sigma,\nu,(l+1,l,l)}\}\phi_{T,l;1}^{(1)} \\ &+ \sqrt{-1}y_1(\partial_{y_2} - 4)\phi_{T,l;2}^{(1)} - y_1y_2\phi_{T,l;3}^{(1)} = 0, \end{aligned}$$

$$
\begin{aligned}
&\sqrt{-1}y_1(\partial_{y_2}-6)\phi^{(1)}_{T,l;1}\\
&+\{(-\partial_{y_1}+\partial_{y_2}+l-2)(-\partial_{y_1}+\partial_{y_2}-l-2)-y_1^2-y_2^2-\tilde{\chi}_{2,\sigma,\nu,(l+1,l,l)}\}\phi^{(1)}_{T,l;2}\\
&+\sqrt{-1}y_2(-\partial_{y_1}+2\partial_{y_3}-2+y_3)\phi^{(1)}_{T,l;3}=0,
\end{aligned}
$$

$$
\begin{aligned}
&-y_1y_2\phi^{(1)}_{T,l;1}+\sqrt{-1}y_2(-\partial_{y_1}+2\partial_{y_3}-4-y_3)\phi^{(1)}_{T,l;2}\\
&+\{(-\partial_{y_2}+2\partial_{y_3}+l-1-y_3)(-\partial_{y_2}+2\partial_{y_3}-l-1+y_3)-y_2^2-\tilde{\chi}_{2,\sigma,\nu,(l+1,l,l)}\}\phi^{(1)}_{T,l;3}=0,
\end{aligned}
$$

$$
\begin{aligned}
&\{(\partial_{y_1}+l-6+\delta_{1i})(\partial_{y_1}-l-\delta_{1i})+(-\partial_{y_1}+\partial_{y_2}+l-4+\delta_{2i})(-\partial_{y_1}+\partial_{y_2}-l-\delta_{2i})\\
&+(-\partial_{y_2}+2\partial_{y_3}+l-2+\delta_{3i}-y_3)(-\partial_{y_2}+2\partial_{y_3}-l-\delta_{3i}+y_3)\\
&-2y_1^2-2y_2^2-\chi_{2,\sigma,\nu,(l+1,l,l)}\}\phi^{(1)}_{T,l;i}-2\delta_{1i}\{2\phi^{(1)}_{T,l;1}-\sqrt{-1}y_1\phi^{(1)}_{T,l;2}\}\\
&-2\delta_{2i}\{\sqrt{-1}y_1\phi^{(1)}_{T,l;1}+\phi^{(1)}_{T,l;2}-\sqrt{-1}y_2\phi^{(1)}_{T,l;3}\}-2\delta_{3i}\sqrt{-1}y_2\phi^{(1)}_{T,l;2}=0,
\end{aligned}
$$

$$
\begin{aligned}
&\Big\{\{(-\partial_{y_1}+\partial_{y_2}+l-3+\delta_{2i})(-\partial_{y_2}+2\partial_{y_3}+l-2+\delta_{3i}-y_3)+y_2^2\}\\
&\qquad\times\{(-\partial_{y_1}+\partial_{y_2}-l-1-\delta_{2i})(-\partial_{y_2}+2\partial_{y_3}-l-\delta_{3i}+y_3)+y_2^2\}\\
&\quad+(\partial_{y_1}+l-5+\delta_{1i})(-\partial_{y_2}+2\partial_{y_3}+l-2+\delta_{3i}-y_3)\\
&\qquad\times(\partial_{y_1}-l-1-\delta_{1i})(-\partial_{y_2}+2\partial_{y_3}-l-\delta_{3i}+y_3)\\
&\quad+\{(\partial_{y_1}+l-5+\delta_{1i})(-\partial_{y_1}+\partial_{y_2}+l-4+\delta_{2i})+y_1^2\}\\
&\qquad\times\{(\partial_{y_1}-l-1-\delta_{1i})(-\partial_{y_1}+\partial_{y_2}-l-\delta_{2i})+y_1^2\}\\
&\quad-2y_1^2(-\partial_{y_2}+2\partial_{y_3}+l-2+\delta_{3i}-y_3)(-\partial_{y_2}+2\partial_{y_3}-l-\delta_{3i}+y_3)\\
&\quad+2y_1^2y_2^2-2y_2^2(\partial_{y_1}+l-5+\delta_{1i})(\partial_{y_1}-l-1-\delta_{1i})-\chi_{4,\sigma,\nu,(l+1,l,l)}\Big\}\phi^{(1)}_{T,l;i}\\
&+2\delta_{1i}\Big\{-\{(-\partial_{y_1}+\partial_{y_2}+l-4)(-\partial_{y_1}+\partial_{y_2}-l)\\
&\quad+(-\partial_{y_2}+2\partial_{y_3}+l-2-y_3)(-\partial_{y_2}+2\partial_{y_3}-l+y_3)-3y_1^2-2y_2^2\}\phi^{(1)}_{T,l;1}\\
&\quad+\sqrt{-1}y_1\{(-\partial_{y_2}+2\partial_{y_3}+l-2-y_3)(-\partial_{y_2}+2\partial_{y_3}-l+y_3)\\
&\quad-(\partial_{y_1}-l-1)(-\partial_{y_1}+\partial_{y_2}-l-1)+\partial_{y_2}-6-y_1^2-y_2^2\}\phi^{(1)}_{T,l;2}\\
&\quad+y_1y_2(2\partial_{y_3}-l-6+y_3)\phi^{(1)}_{T,l;3}\Big\}\\
&+2\delta_{2i}\Big\{-\sqrt{-1}y_1\{(-\partial_{y_2}+2\partial_{y_3}+l-2-y_3)(-\partial_{y_2}+2\partial_{y_3}-l+y_3)\\
&\quad-(\partial_{y_1}+l-4)(-\partial_{y_1}+\partial_{y_2}+l-4)-(\partial_{y_2}-4)-y_1^2-y_2^2\}\phi^{(1)}_{T,l;1}\\
&\quad-\{(\partial_{y_1}+l-5)(\partial_{y_1}-l-1)-y_1^2-2y_2^2\}\phi^{(1)}_{T,l;2}\\
&\quad+\sqrt{-1}y_2\{(\partial_{y_1}+l-5)(\partial_{y_1}-l-1)\\
&\quad-(-\partial_{y_1}+\partial_{y_2}-l-1)(-\partial_{y_2}+2\partial_{y_3}-l-1+y_3)-y_1^2-y_2^2\}\phi^{(1)}_{T,l;3}\Big\}\\
&+2\delta_{3i}\Big\{-y_1y_2(2\partial_{y_3}+l-7-y_3)\phi^{(1)}_{T,l;1}\\
&\quad+\sqrt{-1}y_2\{(-\partial_{y_1}+\partial_{y_2}+l-2)(-\partial_{y_2}+2\partial_{y_3}+l-2-y_3)\\
&\quad-(\partial_{y_1}+l-5)(\partial_{y_1}-l-1)+y_1^2+y_2^2\}\phi^{(1)}_{T,l;2}\Big\}=0,
\end{aligned}
$$

$$\begin{aligned}
&\Big\{\{(\partial_{y_1}+l-4+\delta_{1i})(-\partial_{y_1}+\partial_{y_2}+l-3+\delta_{2i})(-\partial_{y_2}+2\partial_{y_3}+l-2+\delta_{3i}-y_3)\\
&\quad +y_2^2(\partial_{y_1}+l-4+\delta_{1i})+y_1^2(-\partial_{y_2}+2\partial_{y_3}+l-2+\delta_{3i}-y_3)\}\\
&\quad \times\{(\partial_{y_1}-l-2-\delta_{1i})(-\partial_{y_1}+\partial_{y_2}-l-1-\delta_{2i})(-\partial_{y_2}+2\partial_{y_3}-l-\delta_{3i}+y_3)\\
&\quad +y_2^2(\partial_{y_1}-l-2-\delta_{1i})+y_1^2(-\partial_{y_2}+2\partial_{y_3}-l-\delta_{3i}+y_3)\}-\chi_{6,\sigma,\nu,(l+1,l,l)}\Big\}\phi^{(1)}_{T,l;i}\\
&+2\delta_{1i}\Big\{2y_1^2\{(-\partial_{y_2}+2\partial_{y_3}+l-2-y_3)(-\partial_{y_2}+2\partial_{y_3}-l+y_3)-y_2^2\}\phi^{(1)}_{T,l;1}\\
&\quad -\sqrt{-1}y_1\{(-\partial_{y_2}+2\partial_{y_3}+l-2-y_3)(\partial_{y_1}-l-2)(-\partial_{y_1}+\partial_{y_2}-l-2)(-\partial_{y_2}+2\partial_{y_3}-l+y_3)\\
&\quad +y_2^2(-\partial_{y_2}+2\partial_{y_3}+l-2-y_3)(\partial_{y_1}-l-2)\\
&\quad +y_1^2(-\partial_{y_2}+2\partial_{y_3}+l-2-y_3)(-\partial_{y_2}+2\partial_{y_3}-l+y_3)\}\phi^{(1)}_{T,l;2}\\
&\quad -y_1y_2\{(\partial_{y_1}-l-2)(-\partial_{y_1}+\partial_{y_2}-l-1)(-\partial_{y_2}+2\partial_{y_3}-l-1+y_3)\\
&\quad +y_2^2(\partial_{y_1}-l-2)+y_1^2(-\partial_{y_2}+2\partial_{y_3}-l-1+y_3)\}\phi^{(1)}_{T,l;3}\Big\}\\
&+2\delta_{2i}\Big\{\sqrt{-1}y_1\{(\partial_{y_1}+l-3)(-\partial_{y_1}+\partial_{y_2}+l-3)\\
&\qquad \times(-\partial_{y_2}+2\partial_{y_3}+l-2-y_3)(-\partial_{y_2}+2\partial_{y_3}-l+y_3)\\
&\quad +y_2^2(\partial_{y_1}+l-3)(-\partial_{y_2}+2\partial_{y_3}-l-2+y_3)\\
&\quad +y_1^2(-\partial_{y_2}+2\partial_{y_3}+l-2-y_3)(-\partial_{y_2}+2\partial_{y_3}-l+y_3)\}\phi^{(1)}_{T,l;1}\\
&\quad +2y_2^2(\partial_{y_1}+l-4)(\partial_{y_1}-l-2)\phi^{(1)}_{T,l;2}\\
&\quad -\sqrt{-1}y_2(\partial_{y_1}+l-4)\{(\partial_{y_1}-l-2)(-\partial_{y_1}+\partial_{y_2}-l-1)(-\partial_{y_2}+2\partial_{y_3}-l-1+y_3)\\
&\quad +y_2^2(\partial_{y_1}-l-2)+y_1^2(-\partial_{y_2}+2\partial_{y_3}-l-1+y_3)\}\phi^{(1)}_{T,l;3}\Big\}\\
&+2\delta_{3i}\Big\{y_1y_2\{(\partial_{y_1}+l-3)(-\partial_{y_1}+\partial_{y_2}+l-3)(-\partial_{y_2}+2\partial_{y_3}+l-2-y_3)\\
&\quad +y_2^2(\partial_{y_1}+l-3)+y_1^2(-\partial_{y_2}+2\partial_{y_3}+l-2-y_3)\}\phi^{(1)}_{T,l;1}\\
&\quad +\sqrt{-1}y_2\{(\partial_{y_1}+l-4)(-\partial_{y_1}+\partial_{y_2}+l-2)(-\partial_{y_2}+2\partial_{y_3}+l-2-y_3)\\
&\quad +y_2^2(\partial_{y_1}+l-4)+y_1^2(-\partial_{y_2}+2\partial_{y_3}+l-2-y_3)\}(\partial_{y_1}-l-2)\phi^{(1)}_{T,l;2}\Big\}=0,
\end{aligned}$$

for $i=1,2,3$.

For $l\equiv\varepsilon_\sigma \mod 2$, Whittaker function $\Phi(T,S_{(l,l,l-1)}(M_l^{(\sigma;2)}))=\sum_{j=1}^3\phi^{(2)}_{T,l;j}\otimes f(M^{(2)}_{l;j})^*$ satisfy the following holonomic system of partial differential equations of rank 48:

$$\begin{aligned}
&y_1y_2\phi^{(2)}_{T,l;1}+\sqrt{-1}y_1(\partial_{y_2}-4)\phi^{(2)}_{T,l;2}\\
&-\{(\partial_{y_1}-l-3)(\partial_{y_1}+l-3)-y_1^2-\tilde{\chi}_{2,\sigma,\nu,(l,l,l-1)}\}\phi^{(2)}_{T,l;3}=0,
\end{aligned}$$

$$\begin{aligned}
&-\sqrt{-1}y_2(-\partial_{y_1}+2\partial_{y_3}-2-y_3)\phi^{(2)}_{T,l;1}\\
&+\{(-\partial_{y_1}+\partial_{y_2}-l-2)(-\partial_{y_1}+\partial_{y_2}+l-2)-y_1^2-y_2^2-\tilde{\chi}_{2,\sigma,\nu,(l,l,l-1)}\}\phi^{(2)}_{T,l;2}\\
&-\sqrt{-1}y_1(\partial_{y_2}-6)\phi^{(2)}_{T,l;3}=0,
\end{aligned}$$

$$\begin{aligned}
&-\{(-\partial_{y_2}+2\partial_{y_3}-l-1+y_3)(-\partial_{y_2}+2\partial_{y_3}+l-1-y_3)-y_2^2-\tilde{\chi}_{2,\sigma,\nu,(l,l,l-1)}\}\phi^{(2)}_{T,l;1}\\
&+\sqrt{-1}y_2(-\partial_{y_1}+2\partial_{y_3}-4+y_3)\phi^{(2)}_{T,l;2}+y_1y_2\phi^{(2)}_{T,l;3},
\end{aligned}$$

$$\begin{aligned}
&\{(\partial_{y_1}+l-6-\delta_{3i})(\partial_{y_1}-l+\delta_{3i})\\
&\quad+(-\partial_{y_1}+\partial_{y_2}+l-4-\delta_{2i})(-\partial_{y_1}+\partial_{y_2}-l+\delta_{2i})\\
&\quad+(-\partial_{y_2}+2\partial_{y_3}+l-2-\delta_{1i}-y_3)(-\partial_{y_2}+2\partial_{y_3}-l+\delta_{1i}+y_3)\\
&\quad-2y_1^2-2y_2^2-\chi_{2,\sigma,\nu,(l,l,l-1)}\}\phi^{(2)}_{T,l;i}\\
&-2\delta_{1i}\{2\phi^{(2)}_{T,l;1}-\sqrt{-1}y_2\phi^{(2)}_{T,l;2}\}\\
&-2\delta_{2i}\{\sqrt{-1}y_2\phi^{(2)}_{T,l;1}+\phi^{(2)}_{T,l;2}-\sqrt{-1}y_1\phi^{(2)}_{T,l;3}\}-2\delta_{3i}\sqrt{-1}y_1\phi^{(2)}_{T,l;2}=0,
\end{aligned}$$

$$\begin{aligned}
&\Big\{\{(-\partial_{y_1}+\partial_{y_2}+l-3-\delta_{2i})(-\partial_{y_2}+2\partial_{y_3}+l-2-\delta_{1i}-y_3)+y_2^2\}\\
&\qquad\times\{(-\partial_{y_1}+\partial_{y_2}-l-1+\delta_{2i})(-\partial_{y_2}+2\partial_{y_3}-l+\delta_{1i}+y_3)+y_2^2\}\\
&\quad+(\partial_{y_1}+l-5-\delta_{3i})(-\partial_{y_2}+2\partial_{y_3}+l-2-\delta_{1i}-y_3)\\
&\qquad\times(\partial_{y_1}-l-1+\delta_{3i})(-\partial_{y_2}+2\partial_{y_3}-l+\delta_{1i}+y_3)\\
&\quad+\{(\partial_{y_1}+l-5-\delta_{3i})(-\partial_{y_1}+\partial_{y_2}+l-4-\delta_{2i})+y_1^2\}\\
&\qquad\times\{(\partial_{y_1}-l-1+\delta_{3i})(-\partial_{y_1}+\partial_{y_2}-l+\delta_{2i})+y_1^2\}\\
&\quad-2y_1^2(-\partial_{y_2}+2\partial_{y_3}+l-2-\delta_{1i}-y_3)(-\partial_{y_2}+2\partial_{y_3}-l+\delta_{1i}+y_3)\\
&\quad-2y_2^2\{(\partial_{y_1}+l-5-\delta_{3i})(\partial_{y_1}-l-1+\delta_{3i})-y_1^2\}-\chi_{4,\sigma,\nu,(l,l,l-1)}\Big\}\phi^{(2)}_{T,l;i}\\
&+2\delta_{1i}\Big\{-\{(\partial_{y_1}+l-5)(\partial_{y_1}-l-1)+(-\partial_{y_1}+\partial_{y_2}+l-3)(-\partial_{y_1}+\partial_{y_2}-l-1)\\
&\qquad-2y_1^2-3y_2^2\}\phi^{(2)}_{T,l;1}\\
&\quad+\sqrt{-1}y_2\{(\partial_{y_1}+l-5)(\partial_{y_1}-l-1)-(-\partial_{y_1}+\partial_{y_2}-l)(-\partial_{y_2}+2\partial_{y_3}-l+y_3)\\
&\qquad-(-\partial_{y_1}+2\partial_{y_3}-2+y_3)-y_1^2-y_2^2\}\phi^{(2)}_{T,l;2}\\
&\quad-y_1y_2(2\partial_{y_3}-l-3+y_3)\phi^{(2)}_{T,l;3}\Big\}\\
&+2\delta_{2i}\Big\{-\sqrt{-1}y_2\{(\partial_{y_1}+l-5)(\partial_{y_1}-l-1)-(-\partial_{y_1}+\partial_{y_2}+l-3)(-\partial_{y_2}+2\partial_{y_3}+l-3-y_3)\\
&\qquad-\partial_{y_1}+2\partial_{y_3}-4-y_3-y_1^2-y_2^2\}\phi^{(2)}_{T,l;1}\\
&\quad-\{(-\partial_{y_2}+2\partial_{y_3}+l-2-y_3)(-\partial_{y_2}+2\partial_{y_3}-l+y_3)-2y_1^2-y_2^2\}\phi^{(2)}_{T,l;2}\\
&\quad+\sqrt{-1}y_1\{-(\partial_{y_1}-l)(-\partial_{y_1}+\partial_{y_2}-l)\\
&\qquad+(-\partial_{y_2}+2\partial_{y_3}+l-2-y_3)(-\partial_{y_2}+2\partial_{y_3}-l+y_3)-y_1^2-y_2^2\}\phi^{(2)}_{T,l;3}\Big\}\\
&+2\delta_{3i}\Big\{y_1y_2(2\partial_{y_3}+l-9-y_3)\phi^{(2)}_{T,l;1}\\
&\quad+\sqrt{-1}y_1\{(\partial_{y_1}+l-5)(-\partial_{y_1}+\partial_{y_2}+l-5)\\
&\qquad-(-\partial_{y_2}+2\partial_{y_3}+l-2-y_3)(-\partial_{y_2}+2\partial_{y_3}-l+y_3)+y_1^2+y_2^2\}\phi^{(2)}_{T,l;2}\Big\}=0,
\end{aligned}$$

$$\begin{aligned}
&\Big\{\{(\partial_{y_1}+l-4-\delta_{3i})(-\partial_{y_1}+\partial_{y_2}+l-3-\delta_{2i})(-\partial_{y_2}+2\partial_{y_3}+l-2-\delta_{1i}-y_3)\\
&\quad+y_2^2(\partial_{y_1}+l-4-\delta_{3i})+y_1^2(-\partial_{y_2}+2\partial_{y_3}+l-2-\delta_{1i}-y_3)\}\\
&\quad\times\{(\partial_{y_1}-l-2+\delta_{3i})(-\partial_{y_1}+\partial_{y_2}-l-1+\delta_{2i})(-\partial_{y_2}+2\partial_{y_3}-l+\delta_{1i}+y_3)\\
&\qquad+y_1^2(-\partial_{y_2}+2\partial_{y_3}-l+\delta_{1i}+y_3)+y_2^2(\partial_{y_1}-l-2+\delta_{3i})\}-\chi_{6,\sigma,\nu,(l,l,l-1)}\Big\}\phi^{(2)}_{T,l;i}
\end{aligned}$$

$$
\begin{aligned}
&+2\delta_{1i}\Big\{2y_2^2\{(\partial_{y_1}+l-4)(\partial_{y_1}-l-2)-y_1^2\}\phi^{(2)}_{T,l;1}\\
&\quad-\sqrt{-1}y_2\{(\partial_{y_1}+l-4)(\partial_{y_1}-l-2)\\
&\qquad\times(-\partial_{y_1}+\partial_{y_2}-l)(-\partial_{y_2}+2\partial_{y_3}-l+y_3)\\
&\qquad+y_1^2(\partial_{y_1}+l-4)(-\partial_{y_2}+2\partial_{y_3}-l+y_3)\\
&\qquad+y_2^2(\partial_{y_1}+l-4)(\partial_{y_1}-l-2)\}\phi^{(2)}_{T,l;2}\\
&\quad+y_1y_2\{(\partial_{y_1}-l-1)(-\partial_{y_1}+\partial_{y_2}-l-1)(-\partial_{y_2}+2\partial_{y_3}-l+y_3)\\
&\qquad+y_1^2(-\partial_{y_2}+2\partial_{y_3}-l+y_3)+y_2^2(\partial_{y_1}-l-1)\}\phi^{(2)}_{T,l;3}\Big\}\\
&+2\delta_{2i}\Big\{\sqrt{-1}y_2\{(\partial_{y_1}+l-4)(-\partial_{y_1}+\partial_{y_2}+l-3)\\
&\qquad\times(-\partial_{y_2}+2\partial_{y_3}+l-3-y_3)(\partial_{y_1}-l-2)\\
&\qquad+y_2^2(\partial_{y_1}+l-4)(\partial_{y_1}-l-2)\\
&\qquad+y_1^2(-\partial_{y_2}+2\partial_{y_3}+l-3-y_3)(\partial_{y_1}-l)\}\phi^{(2)}_{T,l;1}\\
&\quad+2y_1^2(-\partial_{y_2}+2\partial_{y_3}+l-2-y_3)(-\partial_{y_2}+2\partial_{y_3}-l+y_3)\phi^{(2)}_{T,l;2}\\
&\quad-\sqrt{-1}y_1(-\partial_{y_2}+2\partial_{y_3}+l-2-y_3)\\
&\qquad\times\{(\partial_{y_1}-l-1)(-\partial_{y_1}+\partial_{y_2}-l-1)(-\partial_{y_2}+2\partial_{y_3}-l+y_3)\\
&\qquad+y_1^2(-\partial_{y_2}+2\partial_{y_3}-l+y_3)+y_2^2(\partial_{y_1}-l-1)\}\phi^{(2)}_{T,l;3}\Big\}\\
&+2\delta_{3i}\Big\{-y_1y_2\{(\partial_{y_1}+l-4)(-\partial_{y_1}+\partial_{y_2}+l-3)(-\partial_{y_2}+2\partial_{y_3}+l-3-y_3)\\
&\qquad+y_2^2(\partial_{y_1}+l-4)+y_1^2(-\partial_{y_2}+2\partial_{y_3}+l-3-y_3)\}\phi^{(2)}_{T,l;1}\\
&\quad+\sqrt{-1}y_1\{(\partial_{y_1}+l-4)(-\partial_{y_1}+\partial_{y_2}+l-4)(-\partial_{y_2}+2\partial_{y_3}+l-2-y_3)\\
&\qquad+y_2^2(\partial_{y_1}+l-4)+y_1^2(-\partial_{y_2}+2\partial_{y_3}+l-2-y_3)\}(-\partial_{y_2}+2\partial_{y_3}-l+y_3)\phi^{(2)}_{T,l;2}\Big\}=0,
\end{aligned}
$$

for $i=1,2,3$.

Proof. We obtain the assertion from Proposition 6.5, 6.7 and 6.11. □

From the results of Kostant ([6] Theorem 6.8.1), it follows that the dimension of the intertwining space $\mathcal{I}_{\xi,\pi}$ is 48. Hence, for a multiplicity one K-type τ, the dimension of the space $\mathrm{Wh}(\pi,\xi,\tau)$ of Whittaker functions is also 48. Therefore, every solution of the holonomic systems in the Theorem 6.12 gives an element of in $\mathrm{Wh}(\pi,\xi,\tau)|_{A_{\min}}$.

References

[1] I. M. Gel′fand and A. V. Zelevinskiĭ. A canonical basis in irreducible representations of gl_3 and its applications. In *Group-theoretic methods in physics, Vol. 2 (Russian) (Jūrmala, 1985)*, pp. 31–45. “Nauka”, Moscow, 1986.

[2] Miki Hirano. Fourier-Jacobi type spherical functions for principal series representations of $\mathrm{Sp}(2,\mathbf{R})$. *Indag. Math. (N.S.)*, Vol. 15, No. 1, pp. 43–53, 2004.

[3] Miki Hirano, Taku Ishii, and Takayuki Oda. Confluence from Siegel-Whittaker functions to Whittaker functions on $\mathrm{Sp}(2,\mathbb{R})$. *Math. Proc. Cambridge Philos. Soc.*, Vol. 141, No. 1, pp. 15–31, 2006.

[4] Miki Hirano and Takayuki Oda. Integral switching engine for special clebsch-gordan coefficients for the representations of $\mathfrak{gl}_3$ with respect to gelfand-zelevinsky basis, 2003, preprint.

[5] Taku Ishii. On principal series Whittaker functions on $\mathrm{Sp}(2,\mathbf{R})$. *J. Funct. Anal.*, Vol. 225, No. 1, pp. 1–32, 2005.

[6] Bertram Kostant. On Whittaker vectors and representation theory. *Invent. Math.*, Vol. 48, No. 2, pp. 101–184, 1978.

[7] Hrvoje Kraljević. Representations of the universal convering group of the group SU(n, 1). *Glasnik Mat. Ser. III*, Vol. 8(28), pp. 23–72, 1973.

[8] Soo Teck Lee, Kyo Nishiyama, and Akihito Wachi. Intersection of harmonics and capelli identities for symmetric pairs, preprint.

[9] Hans Maass. *Siegel's modular forms and Dirichlet series*. Springer-Verlag, Berlin, 1971. Dedicated to the last great representative of a passing epoch. Carl Ludwig Siegel on the occasion of his seventy-fifth birthday, Lecture Notes in Mathematics, Vol. 216.

[10] Tadashi Miyazaki. The $(\mathfrak{g}, k)$-module structures of principal series representations of $\mathrm{Sp}(2, \mathbf{R})$, 2007, Master's thesis, (University of Tokyo).

[11] Takuya Miyazaki. The generalized Whittaker functions for $\mathrm{Sp}(2, \mathbf{R})$ and the gamma factor of the Andrianov L-function. *J. Math. Sci. Univ. Tokyo*, Vol. 7, No. 2, pp. 241–295, 2000.

[12] Takuya Miyazaki and Takayuki Oda. Principal series Whittaker functions on $\mathrm{Sp}(2; \mathbf{R})$. Explicit formulae of differential equations. In *Automorphic forms and related topics (Seoul, 1993)*, pp. 59–92. Pyungsan Inst. Math. Sci., Seoul, 199?

[13] Shinji Niwa. Commutation relations of differential operators and Whittaker functions on $\mathrm{Sp}_2(\mathbf{R})$. *Proc. Japan Acad. Ser. A Math. Sci.*, Vol. 71, No. 8, pp. 189–191, 1995.

[14] Takayuki Oda. An explicit integral representation of Whittaker functions on $\mathrm{Sp}(2; \mathbf{R})$ for the large discrete series representations. *Tohoku Math. J. (2)*, Vol. 46, No. 2, pp. 261–279, 1994.

[15] Takayuki Oda. The standard $(\mathfrak{g}, k)$-modules of $\mathrm{Sp}(2, \mathbf{R})$ I, preprint.

[16] Takayuki Oda. The standard $(\mathfrak{g}, k)$-modules of $\mathrm{Sp}(2, \mathbf{R})$ II, preprint.

[17] Ernest Thieleker. On the integrable and square-integrable representations of $\mathrm{Spin}(1, 2m)$. *Trans. Amer. Math. Soc.*, Vol. 230, pp. 1–40, 1977.

Department of Mathematical Sciences, University of Tokyo
E-mail address: `miyaza@ms.u-tokyo.ac.jp`